

Product Datasheet

HIF-1 alpha Antibody (HA111) - BSA Free NB100-296

Unit Size: 0.1 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 9

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB100-296

Updated 9/9/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB100-296



NB100-296


HIF-1 alpha Antibody (HA111) - BSA Free

Product Information	
Unit Size	0.1 ml
Concentration	1.0 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	HA111
Preservative	0.02% Sodium Azide
Isotype	IgG2 Alpha
Purity	Protein G purified
Buffer	PBS
Target Molecular Weight	93 kDa
Product Description	
Description	Novus Biologicals Mouse HIF-1 alpha Antibody (HA111) - BSA Free (NB100-296) is a monoclonal antibody validated for use in IHC, WB and IP. Anti-HIF-1 alpha Antibody: Cited in 9 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Mouse
Gene ID	3091
Gene Symbol	HIF1A
Species	Human, Mouse, Rabbit
Reactivity Notes	Mouse reactivity reported in literature (PMID: 16738327). Rabbit reactivity reported in scientific literature (PMID: 16738327).
Immunogen	This HIF-1 alpha Antibody (HA111) was developed against Human HIF-1 alpha, corresponding to amino acids 329 - 530 [Uniprot# Q16665].
Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, Immunohistochemistry, Immunoprecipitation
Recommended Dilutions	Western Blot 0.5 - 2.0 ug/ml, Immunohistochemistry 1:100, Immunoprecipitation 1:10 - 1:500. Use reported in scientific literature (PMID 18222538), Immunohistochemistry-Paraffin 1:100
Application Notes	In IHC-P, staining was observed in the nucleus of human ovarian cancer tumor. Prior to immunostaining paraffin tissues, antigen retrieval with sodium citrate buffer (pH 6.0) is recommended. Nuclear extracts should be used for Western blot analysis.




Images

Immunohistochemistry-Paraffin: HIF-1 alpha Antibody (HA111) [NB100-296] - HIF-1 antibody was tested in human ovarian cancer tumor xenograft using DAB with hematoxylin counterstain.



Western Blot: HIF-1 alpha Antibody (HA111) [NB100-296] - Detection of HIF-1 alpha (125-130 kDa) from human placental villous explant total protein.



Publications

Iorio J, Lastraioli E, Tofani L et al. hERG1 and HIF-2 alpha Behave as Biomarkers of Positive Response to Bevacizumab in Metastatic Colorectal Cancer Patients *Transl Oncol* 2020-02-24 [PMID: 32105990] (IHC-P, Human)

Lee KE, Spata M, Maduka R et al. Hif1a Deletion Limits Tissue Regeneration via Aberrant B Cell Accumulation in Experimental Pancreatitis. *Cell Rep* 2018-06-19 [PMID: 29924990] (IF/IHC, Mouse)

Li F, Lee KE, Simon MC. Detection of Hypoxia and HIF in Paraffin-Embedded Tumor Tissues *Methods Mol. Biol.* 2018-01-14 [PMID: 29330808] (IHC-P)

Jensen RL, Ragel BT, Whang K, Gillespie D Inhibition of hypoxia inducible factor-1alpha (HIF-1alpha) decreases vascular endothelial growth factor (VEGF) secretion and tumor growth in malignant gliomas. *J Neurooncol* 2006-07-01 [PMID: 16612574]

Lee KE, Spata M, Bayne LJ et al. Hif1alpha deletion reveals pro-neoplastic function of B cells in pancreatic neoplasia. *Cancer Discov.* 2015-12-29 [PMID: 26715642] (IHC-P, Human)

Zheng X, Ruas JL, Cao R et al. Cell-type-specific regulation of degradation of hypoxia-inducible factor 1a: Role of subcellular compartmentalization. *Mol Cell Biol* 2006-06-01 [PMID: 16738327] (IP, WB, Rabbit)

Rajakumar A, Michael HM, Daftary A et al. Proteasomal activity in placentas from women with preeclampsia and intrauterine growth restriction: implications for expression of HIF-alpha proteins. *Placenta* 2008-03-01 [PMID: 18222538] (IP, Human)

Lauzier, MC et al. (2R)-[(4-Biphenylsulfonyl)amino]-N-hydroxy-3-phenylpropionamide (BiPS), a matrix metalloprotease inhibitor, is a novel and potent activator of hypoxia-inducible factors. *Mol Pharmacol*;74(1):282-8. 2008-07-01 [PMID: 18424552]

Gillespie, DL et al. Silencing of hypoxia inducible factor-1alpha by RNA interference attenuates human glioma cell growth in vivo. *Clin Cancer Res.* 2007-01-01 [PMID: 17438103]

Procedures

Immunohistochemistry-Paraffin protocol for HIF-1 alpha Antibody (NB100-296)

HIF-1 alpha Antibody (HA111):

Immunohistochemistry-Paraffin Embedded Sections

Antigen Unmasking:

Bring slides to a boil in 10 mM sodium citrate buffer (pH 6.0) then maintain at a sub-boiling temperature for 10 minutes. Cool slides on bench-top for 30 minutes.

Staining:

1. Wash sections in deionized water three times for 5 minutes each.
2. Wash sections in wash buffer for 5 minutes.
3. Block each section with 100-400 ul blocking solution for 1 hour at room temperature.
4. Remove blocking solution and add 100-400 ul diluted primary antibody. Incubate overnight at 4 degrees C.
5. Remove antibody solution and wash sections in wash buffer three times for 5 minutes each.
6. Add 100-400 ul biotinylated diluted secondary antibody. Incubate 30 minutes at room temperature.
7. Remove secondary antibody solution and wash sections three times with wash buffer for 5 minutes each.
8. Add 100-400 ul Streptavidin-HRP reagent to each section and incubate for 30 minutes at room temperature.
9. Wash sections three times in wash buffer for 5 minutes each.
10. Add 100-400 ul DAB substrate to each section and monitor staining closely.
11. As soon as the sections develop, immerse slides in deionized water.
12. Counterstain sections in hematoxylin.
13. Wash sections in deionized water two times for 5 minutes each.
14. Dehydrate sections.
15. Mount coverslips.





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NB100-296

NB800-PC26	COS-7 Nuclear Hypoxic Induced Cell Lysate
NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB7539	Goat anti-Mouse IgG (H+L) Secondary Antibody [HRP]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-296

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

