

Product Datasheet

ATR Antibody - BSA Free NB100-323

Unit Size: 0.1 ml

Store at 4C. Do not freeze.

www.novusbio.com



technical@novusbio.com

Reviews: 1 Publications: 6

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB100-323

Updated 9/9/2025 v.20.1

**Earn rewards for product
reviews and publications.**

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB100-323



NB100-323

ATR Antibody - BSA Free

Product Information	
Unit Size	0.1 ml
Concentration	1.0 mg/ml
Storage	Store at 4C. Do not freeze.
Clonality	Polyclonal
Preservative	0.09% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	Tris-Citrate/Phosphate (pH 7.0 - 8.0)
Target Molecular Weight	301 kDa


Product Description	
Description	Novus Biologicals Rabbit ATR Antibody - BSA Free (NB100-323) is a polyclonal antibody validated for use in WB and IP. Anti-ATR Antibody: Cited in 6 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	545
Gene Symbol	ATR
Species	Human, Mouse
Reactivity Notes	Mouse reactivity reported in scientific literature (PMID:24356582).
Immunogen	The immunogen recognized by this antibody maps to a region between residues 400 and 450 of human Ataxia Telangiectasia and Rad3-related using the numbering given in entry NP_001175.1 (GeneID 545).

Product Application Details	
Applications	Western Blot, Immunoprecipitation
Recommended Dilutions	Western Blot 1:2000- 1:10000, Immunoprecipitation 2-5 ug/mg lysate
Application Notes	ATR antibody validated for WB from a verified customer review.




Images


Western Blot: ATR Antibody [NB100-323] - Whole cell lysate (15 ug) from HeLa, HEK293T, and Jurkat cells prepared using NETN lysis buffer. Antibody: Affinity purified rabbit anti-ATR antibody used for WB at 0.4 ug/ml. Detection: Chemiluminescence with an exposure time of 3 minutes.



Western Blot: ATR Antibody [NB100-323] - Whole cell lysate from HeLa (5, 15 and 50 ug for WB; 1 mg for IP, 20% of IP loaded) and 293T (T; 50 ug) cells. Antibodies: Affinity purified rabbit anti-ATR antibody NB100-323 lot 3 used for WB at 0.1 ug/ml (A) and 1 ug/ml (B) and used for IP at 3 ug/mg lysate (B). ATR was also immunoprecipitated by a previous lot (lot 2) of this antibody. Detection: Chemiluminescence with exposure times of 30 seconds (A) and 10 seconds (B).



Western Blot: ATR Antibody [NB100-323] - H1299 (human lung cancer cells) and HCC 44 (human liver cancer cells). Antibody at 1:500. Image submitted by a verified customer review.



Publications

Hung PJ, Johnson B, Chen BR et al. MRI Is a DNA Damage Response Adaptor during Classical Non-homologous End Joining. *Mol. Cell.* 2018-07-19 [PMID: 30017584]

Morales AJ, Carrero JA, Hung PJ et al. A type I IFN-dependent DNA damage response regulates the genetic program and inflammasome activation in macrophages. *Elife* 2017-03-31 [PMID: 28362262] (WB)

Boichuk S, Hu L, Hein J, Gjoerup OV. Multiple DNA damage signaling and repair pathways deregulated by simian virus 40 large T antigen. *J Virol* 2010-08-01 [PMID: 20519379]

Min W, Bruhn C, Grigaravicius et al. Poly(ADP-ribose) binding to Chk1 at stalled replication forks is required for S-phase checkpoint activation. *Nat Commun* 2013-12-20 [PMID: 24356582] (WB, Mouse)

Shao L, Fujii H, Colmegna I et al. Deficiency of the DNA repair enzyme ATM in rheumatoid arthritis. *J Exp Med*;206 (6):1435-1449. 2009-01-01 [PMID: 19451263]

Agarwal, C, Tyagi, A, Agarwal, R. Gallic acid causes inactivating phosphorylation of cdc25A/cdc25C-cdc2 via ATM-Chk2 activation, leading to cell cycle arrest, and induces apoptosis in human prostate carcinoma DU145 cells. *Mol Cancer Ther*5(12):3294-302. 2006-12-01 [PMID: 17172433]



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NB100-323

NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-323

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

