

Product Datasheet

TLR4 Antibody (HTA125) [FITC] NB100-56059

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com



technical@novusbio.com

Publications: 26

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB100-56059

Updated 2/5/2026 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB100-56059



NB100-56059**TLR4 Antibody (HTA125) [FITC]**


Product Information	
Unit Size	0.1 ml
Concentration	Please see the vial label for concentration. If unlisted please contact technical services.
Storage	Store at 4C in the dark.
Clonality	Monoclonal
Clone	HTA125
Preservative	0.05% Sodium Azide
Isotype	IgG2a
Conjugate	FITC
Purity	Protein G purified
Buffer	50 mM Sodium Borate
Target Molecular Weight	95.7 kDa

Product Description	
Host	Mouse
Gene ID	7099
Gene Symbol	TLR4
Species	Human, Mouse, Canine
Specificity/Sensitivity	The antibody has been shown to block the activation of monocytes with LPS (Paik, et al, 2003).
Immunogen	This TLR4 Antibody (HTA125) [FITC] was developed by immunizing mice with Ba/F3 cell line expressing human TLR4 cell surface antigen.

Product Application Details	
Applications	Flow Cytometry, Flow (Cell Surface), Flow (Intracellular), Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Western Blot (Negative)
Recommended Dilutions	Flow Cytometry 1uL / 1 million cells, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Flow (Cell Surface), Flow (Intracellular), Western Blot (Negative)
Application Notes	Optimal dilution of this antibody should be experimentally determined.

Images

Flow Cytometry: TLR4 Antibody (HTA125) [FITC] [NB100-56059] - An intracellular stain was performed on MG-63 cells with TLR4 Antibody [HTA125] NB100-56059 (blue) and a matched isotype control (orange). Cells were fixed with 4% PFA and then permeabilized with 0.1% saponin. Cells were incubated in an antibody dilution of 10 ug/mL for 30 minutes at room temperature. Both antibodies were conjugated to FITC.



Copyright © 2018 Novus Biologicals

Publications

Akhter N, Madhoun A, Arefanian H et al Oxidative Stress Induces Expression of the Toll-Like Receptors (TLRs) 2 and 4 in the Human Peripheral Blood Mononuclear Cells: Implications for Metabolic Inflammation Cell. Cell Physiol Biochem. 2019-01-01 [PMID: 31162913] (FLOW, Human)

Details:

Citation using the PE version of this antibody.

Ahmad Rasheed, Al-Roub Areej, Kochumon Shihab et al. The Synergy between Palmitate and TNF- α for CCL2 Production Is Dependent on the TRIF/IRF3 Pathway: Implications for Metabolic Inflammation J Immunol. 2018-04-09 [PMID: 29632147] (FLOW, Human)

Details:

Citation using the FITC form of this antibody.

Heftrig D, Sturm R, Oppermann E et al. Impaired Surface Expression of HLA-DR, TLR2, TLR4, and TLR9 in Ex Vivo-In Vitro Stimulated Monocytes from Severely Injured Trauma Patients. Mediators Inflamm. [PMID: 28255201]

Details:

Citation using the FITC form of this antibody.

Larsson O, Tengroth L, Xu Y et al. Substance P represents a novel first-line defense mechanism in the nose. J Allergy Clin Immunol. [PMID: 28219705] (FLOW, Mouse)

Details:

Citation using the FITC form of this antibody.

Jamin A, Dehoux L, Dossier C et al. Toll-like receptor 3 expression and function in childhood idiopathic nephrotic syndrome Clin Exp Immunol. [PMID: 26123900] (FLOW, Human)

Zanoni G, Navone R, Lunardi C et al. In Celiac Disease, a Subset of Autoantibodies Against Transglutaminase Binds Toll-Like Receptor 4 and Induces Activation of Monocytes PLoS Med 2006-09-01 [PMID: 16984219]

Cognasse F, Hamzeh H, Chavarin P et al. Evidence of Toll-like receptor molecules on human platelets. Immunol Cell Biol. 2005-04-01 [PMID: 15748217]

Details:

TLR2-PE (IMG-416D), TLR4-PE (IMG-417D), TLR6 (IMG-304A), TLR8-PE (IMG-321D), TLR9-PE (IMG-305D). Applications: Intracellular Flow Cytometry and Cell Surface Flow Cytometry: Figs 1 and 2. A comparison of staining results, intracellular versus cell surface flow cytometry is shown. Cell type: Human platelets.

Mempel M, Voelcker V, Kollisch G et al. Toll-like receptor expression in human keratinocytes: nuclear factor kappaB controlled gene activation by Staphylococcus aureus is toll-like receptor 2 but not toll-like receptor 4 or platelet activating factor receptor dependent. J Invest Dermatol. 2003-12-01 [PMID: 14675188] (ICC/IF, Human)

Details:

TLR2 (IMG-416) 2. TLR4 (IMG-417) [IF/ICC, Fig.2A and 2D (human keratinocytes)].

Pietschmann K, Beetz S, Welte S et al. Toll-like receptor expression and function in subsets of human gammadelta T lymphocytes. Scand J Immunol. 2009-09-01 [PMID: 19703014]

Wu CY, Chi PL, Hsieh HL et al. TLR4-dependent induction of vascular adhesion molecule-1 in rheumatoid arthritis synovial fibroblasts: Roles of cytosolic phospholipase A(2) α /cyclooxygenase-2. J Cell Physiol. 2010-05-01 [PMID: 20112284]



Prabha C, Rajashree P, Sulochana DD. TLR2 and TLR4 expression on the immune cells of tuberculous pleural fluid. Immunol Lett. 2008-04-15 [PMID: 18295348] (Flow Cytometry Control, Human)

Details:

TLR2- FITC (IMG-416C): Flow (cell surface): Figs. 1A, B (human CD4+T cells, CD8+T cells, B cells, CD16+56+ cells and monocytes); 2(CD4+T cells); 4A, B (human Treg cells). Flow (intracellular): Fig. 3A, B (CD4+T cells) 2. TLR4-FITC (IMG-417C).Flow (cell surface): Figs. 1B, C (human CD4+T cells, CD8+T cells, B cells, CD16+56+ cells and monocytes); 2(CD4+T cells); 4A, B (human Treg cells). Flow (intracellular): Fig. 3A, B (CD4+T cells).

Matsunaga N, Tsuchimori N, Matsumoto T, li M. TAK-242 (resatorvid), a small-molecule inhibitor of Toll-like receptor (TLR) 4 signaling, binds selectively to TLR4 and interferes with interactions between TLR4 and its adaptor molecules. Mol Pharmacol. 2011-01-01 [PMID: 20881006]

More publications at <http://www.novusbio.com/NB100-56059>





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-
techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-56059

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

