

Product Datasheet

Bad Antibody - BSA Free

NB100-56080

Unit Size: 0.05 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Reviews: 1

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB100-56080

Updated 9/9/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB100-56080



NB100-56080

Bad Antibody - BSA Free

Product Information

Unit Size	0.05 ml
Concentration	This product is unpurified. The exact concentration of antibody is not quantifiable.
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.02% Sodium Azide
Isotype	IgG
Purity	Unpurified
Buffer	Whole antisera

Product Description


Description	Novus Biologicals Rabbit Bad Antibody - BSA Free (NB100-56080) is a polyclonal antibody validated for use in IHC, WB, ICC/IF and IP. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	572
Gene Symbol	BAD
Species	Human
Immunogen	A full length recombinant protein corresponding to human Bad was used as immunogen.

Product Application Details


Applications	Western Blot, Immunohistochemistry-Paraffin, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunoprecipitation
Recommended Dilutions	Western Blot 1:1000-1:2000, Immunohistochemistry 1:2000, Immunocytochemistry/ Immunofluorescence 1:500-1:2000, Immunoprecipitation 1:50-1:200, Immunohistochemistry-Paraffin 1:2000

Images


Western Blot: Bad Antibody [NB100-56080] - Analysis of Bad in Daoy whole cell lysate using anti-Bad antibody. Image from verified customer review.



Immunocytochemistry/Immunofluorescence: Bad Antibody [NB100-56080] - Immunofluorescence microscopy of BAD using NB100-56080 at 1:2000. Du145 human prostate carcinoma cells were cultured without (A) or with (B) and (C) 1 μ M of the triphosphatase inhibitor thapsigargin (THG) for 12 hr. A and B, staining with BAD antibody, followed by a FITC-conjugated secondary antibody. C, staining with a mitochondrial marker (antibody to mitochondrial Hsp60), followed by a rhodamine-conjugated secondary antibody. THG induces Ca^{2+} release from internal stores which can promote apoptosis. BAD staining was located diffusely throughout the cytoplasm of untreated cells (A), and localized to the mitochondria in treated cells (B).



Immunohistochemistry-Paraffin: Bad Antibody [NB100-56080] - IHC analysis of formalin fixed paraffin-embedded (FFPE) human liver using 1:2000 conc. of Bad antibody on a Bond Rx autostainer (Leica Biosystems). The assay involved 20 minutes of heat induced antigen retrieval (HIER) using 10mM sodium citrate buffer (pH 6.0) and endogenous peroxidase quenching with peroxide block. The sections were incubated with primary antibody for 30 minutes and Bond Polymer Refine Detection (Leica Biosystems) with DAB was used for signal development followed by counterstaining with hematoxylin. Whole slide scanning and capturing of representative images was performed using Aperio AT2 (Leica Biosystems). Cytoplasmic staining of Bad was observed. Staining was performed by Histowiz.





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NB100-56080

NBL1-07900	Bad Overexpression Lysate
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-56080

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications