

# Product Datasheet

## Caspase-3 Antibody - (active/cleaved) - BSA Free NB100-56113

Unit Size: 0.05 ml

Store at -20C. Avoid freeze-thaw cycles.

[www.novusbio.com](http://www.novusbio.com)



[technical@novusbio.com](mailto:technical@novusbio.com)

**Publications: 81**

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:  
[www.novusbio.com/NB100-56113](http://www.novusbio.com/NB100-56113)

Updated 9/9/2025 v.20.1

Earn rewards for product  
reviews and publications.

Submit a publication at [www.novusbio.com/publications](http://www.novusbio.com/publications)

Submit a review at [www.novusbio.com/reviews/destination/NB100-56113](http://www.novusbio.com/reviews/destination/NB100-56113)



**NB100-56113**


Caspase-3 Antibody - (active/cleaved) - BSA Free

<b>Product Information</b>	
<b>Unit Size</b>	0.05 ml
<b>Concentration</b>	This product is unpurified. The exact concentration of antibody is not quantifiable.
<b>Storage</b>	Store at -20C. Avoid freeze-thaw cycles.
<b>Clonality</b>	Polyclonal
<b>Preservative</b>	0.05% Sodium Azide
<b>Isotype</b>	IgG
<b>Purity</b>	Unpurified
<b>Buffer</b>	Whole antisera
<b>Target Molecular Weight</b>	31.7 kDa
<b>Product Description</b>	
<b>Description</b>	Novus Biologicals Rabbit Caspase-3 Antibody - (active/cleaved) - BSA Free (NB100-56113) is a polyclonal antibody validated for use in IHC, WB, Flow, ICC/IF and IP. Anti-Caspase-3 Antibody: Cited in 77 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
<b>Host</b>	Rabbit
<b>Gene ID</b>	836
<b>Gene Symbol</b>	CASP3
<b>Species</b>	Human, Mouse, Rat, Gerbil
<b>Reactivity Notes</b>	Use in Rat reported in scientific literature (PMID:34597692). Rat reactivity reported in scientific literature (PMID:32818590).
<b>Immunogen</b>	This Caspase-3 Antibody - (active/cleaved) was developed against catalytically active human caspase-3 protein.
<b>Product Application Details</b>	
<b>Applications</b>	Western Blot, Immunohistochemistry-Paraffin, Flow Cytometry, Flow (Intracellular), Immunoblotting, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Frozen, Immunoprecipitation
<b>Recommended Dilutions</b>	Western Blot 1:1000-1:2000, Flow Cytometry reported in scientific literature (PMID 28287161), Immunohistochemistry, Immunocytochemistry/ Immunofluorescence reported in scientific literature (PMID 29963272), Immunoprecipitation 1:50-1:200, Immunohistochemistry-Paraffin 1:1000-1:5000, Immunohistochemistry-Frozen reported in scientific literature (PMID 29864441), Immunoblotting reported in scientific literature (PMID 27912032), Flow (Intracellular) reported in scientific literature (PMID 24804954)
<b>Application Notes</b>	Preferentially detects active caspase-3 (large subunit: ~14-21 kDa, and small subunit: ~10 kDa). However, it may also detect pro-caspase-3 (~32 kDa) in some cell or tissue systems. Nuclear immunostaining of caspase-3 is considered be an indication of active/cleaved caspase-3.




## Images


**Immunohistochemistry: Caspase-3 Antibody - (active/cleaved) [NB100-56113]** - Immunohistochemistry analysis of sectioned tumor tissues from the MDA-MB-231 study. Each section was subjected to the specified antibody followed by a biotinylated secondary antibody. Detection was done using a DAB Peroxidase HRP Substrate Kit (brown) followed by Hematoxylin counterstaining (purple). Images were obtained using inverted bright field microscopy. Sectioning results are representative of three individual tumors. Scale bar is 50 microns. Statistical analysis using One-Way ANOVA. \* $p < 0.05$  vs tumor volume of the control. Image collected and cropped by CiteAb from the following publication ([nature.com/articles/s41598-017-01230-4](https://www.nature.com/articles/s41598-017-01230-4)), licensed under a CC-BY license.




**Western Blot: Caspase-3 Antibody - (active/cleaved) [NB100-56113]** - The protein expression levels of Bcl-2, Bax and cleaved caspase-3 were detected in the different groups using Western blotting and were normalized to the housekeeping gene GAPDH. Image collected and cropped by CiteAb from the following publication ([onlinelibrary.wiley.com/doi/abs/10.1111/jcmm.14135](https://onlinelibrary.wiley.com/doi/abs/10.1111/jcmm.14135)), licensed under a CC-BY license.




**Western Blot: Caspase-3 Antibody - (active/cleaved) [NB100-56113]** - Cell apoptosis was detected by flow cytometric analysis, and the percentage of apoptotic cells transfected with miR-200c inhibitor or NC inhibitor and treated with AngII or PBS for 48 h was determined. Image collected and cropped by CiteAb from the following publication ([onlinelibrary.wiley.com/doi/abs/10.1111/jcmm.14135](https://onlinelibrary.wiley.com/doi/abs/10.1111/jcmm.14135)), licensed under a CC-BY license.




**Immunohistochemistry-Paraffin: Caspase-3 Antibody - (active/cleaved) [NB100-56113]** - Dog ischemic brain stained for Active/Cleaved Caspase-3 expression using Caspase-3 Antibody - (active/cleaved) (NB100-56113) at 1:2000. Staining is seen in the nuclei of dying neurons (black arrow) but not in the morphologically normal nuclei (open arrows). Caspase-3 expression in the nucleus is considered to be a marker of active/caspase-3 expression and apoptosis. Hematoxylin-eosin counterstain.




Immunohistochemistry-Paraffin: Caspase-3 Antibody - (active/cleaved) [NB100-56113] - Irradiated mouse spleen stained for Active/Cleaved Caspase-3 expression using Caspase-3 Antibody - (active/cleaved) (NB100-56113) at 1:2000. Staining is seen in the nuclei of a subset of the cell population. Caspase-3 expression in the nucleus is considered to be a marker of active/caspase-3 expression and apoptosis. Hematoxylin-eosin counterstain.




Antidepressants-mediated effect on cellular protein content. HT-22 cells were treated with antidepressants for 48 and 96 h and densitometry analysis of NF- $\kappa$ B (b), p16 (c), p21 (d), p27 (e), p53 (f), TRF1 (g), TRF2 (h), calnexin (i), NuMa (j), cleaved caspase 3 (k), Bcl-2 (l) was evaluated. Representative Western Blots are presented (a). Bars indicate SD,  $n = 3$ ,  $***/^{^^}p < 0.001$ ,  $**/^{^^}p < 0.01$ ,  $*/^{^^}p < 0.05$ , no indication—no statistical significance (one-way ANOVA and Dunnett's a posteriori test)




Immunohistochemistry: Caspase-3 Antibody - (active/cleaved) [NB100-56113] - Pathology in DOX treated acute & recovery phase mice. Regions of myofiber loss & frank replacement fibrosis were noted, most commonly in atria (A, acute phase), & rarely in ventricles (D, recovery phase). These areas were accompanied by macrophage infiltration (B, E) & myofibroblast proliferation (C) consistent with fibroplasia. Rare myofibers were matrix metalloproteinase 2 (F, recovery phase animal) or caspase-3 positive (G, acute phase animal). Reticulin staining (A, D); Immunohistochemistry: Iba 1 (B, E; macrophages), alpha SMA (C), MMP-2 (F) & cleaved caspase -3 (G) Bar = 100 $\mu$ m (A-C); 50 $\mu$ m (D-F); 20 $\mu$ m (G). Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/31263061>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Western Blot: Caspase-3 Antibody - (active/cleaved) [NB100-56113] - Antidepressants-mediated effect on cellular protein content. HT-22 cells were treated with antidepressants for 48 & 96 h & densitometry analysis of NF- $\kappa$ B (b), p16 (c), p21 (d), p27 (e), p53 (f), TRF1 (g), TRF2 (h), calnexin (i), NuMa (j), cleaved caspase 3 (k), Bcl-2 (l) was evaluated. Representative Western Blots are presented (a). Bars indicate SD,  $n = 3$ ,  $***/^{^^}p < 0.001$ ,  $**/^{^^}p < 0.01$ ,  $*/^{^^}p < 0.05$ , no indication—no statistical significance (one-way ANOVA & Dunnett's a posteriori test) Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/31278507>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Immunocytochemistry/ Immunofluorescence: Caspase-3 Antibody - (active/cleaved) [NB100-56113] - Morphological changes & expression of key apoptosis-related molecules in 4T1 cells after DE-EDCP treatment (A) Morphological changes of 4T1 cells exposed to various concentrations of DE-EDCP for 24h. (B) Immunofluorescence staining for Bcl-2 (green), Bax (green) & cleaved caspase-3 (green) together with DNA staining with DAPI (blue) in 4T1 cells incubated with DE-EDCP or cisplatin (31.25  $\mu$ M) for 24h, as well as in untreated cells (magnification at x200). (C) mRNA expression of Bcl-2, Bax & caspase-3 quantified by RT-PCR in 4T1 cells after DE-EDCP 24h treatment. DE-EDCP treatment markedly increased the expression of Bax & caspase-3 mRNA & decreased the expression of Bcl-2 mRNA in 4T1 cells.  $\beta$ -actin mRNA was used as an internal control. Data points are represented by the expression ratio & mean $\pm$ SD fold of control in 4T1 cells. (\* Bcl-2-: DE-EDCP vs. untreated  $p=0.03$ ; DE-EDCP vs. cisplatin  $p=0.006$ ; cisplatin vs. untreated  $p=0.001$ ; Bax-: DE-EDCP vs. untreated  $p=0.011$ ; cisplatin vs. untreated  $p=0.009$ ; caspase-3-: DE-EDCP vs. untreated  $p=0.015$ ; DE-EDCP vs. cisplatin  $p=0.021$ ) Image collected & cropped by CiteAb from the following publication (<https://www.oncotarget.com/lookup/doi/10.18632/oncotarget.25610>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



## Publications

Corsetti G, Romano C, Codenotti S et al. A Diet Rich in Essential Amino Acids Inhibits the Growth of HCT116 Human Colon Cancer Cell In Vitro and In Vivo. *International Journal of Molecular Sciences* 2025-07-29 [PMID: 40725262]

Liu D, Tang X, Huang Z et al. Histone deacetylase HDAC2 regulates microRNA-125a expression in neuroblastoma Brain and behavior 2022-01-21 [PMID: 35060363] (Immunohistochemistry-Paraffin)

Shaalán AK, Teshima THN, Tucker AS, Proctor GB. Inhibition of Aurora Kinase B activity disrupts development and differentiation of salivary glands *Cell Death Discovery* 2021-01-18 [PMID: 33462217] (Immunohistochemistry-Paraffin)

Haushalter C, Schuhbaur B, Doll P, Rhinn M. Meningeal retinoic acid contributes to neocortical lamination and radial migration during mouse brain development *Biology Open* 2017-02-15 [PMID: 28011626] (Immunohistochemistry-Paraffin)

Albadawy R, Hasanin AH, Agwa SHA et al. Rosavin Ameliorates Hepatic Inflammation and Fibrosis in the NASH Rat Model via Targeting Hepatic Cell Death *International Journal of Molecular Sciences* 2022-09-05 [PMID: 36077546] (Immunohistochemistry-Paraffin)

Martin KK, Parvin S, Garraway SM. Peripheral inflammation accelerates the onset of mechanical hypersensitivity after spinal cord injury and engages TNF $\alpha$  signaling mechanisms *J. Neurotrauma* 2018-12-06 [PMID: 30520675] (Immunohistochemistry-Paraffin)

Silconi ZB, Rosic V, Benazic S et al. The Pt(S-pr-thiosal)<sub>2</sub> and BCL1 Leukemia Lymphoma: Antitumor Activity In Vitro and In Vivo *International Journal of Molecular Sciences* 2022-07-24 [PMID: 35897737] (Immunohistochemistry-Paraffin)

Wang S, Liu A, Xu C, Hou J et Al. GLP-1(7-36) protected against oxidative damage and neuronal apoptosis in the hippocampal CA region after traumatic brain injury by regulating ERK5/CREB *Mol Biol Rep* 2024-02-20 [PMID: 38374452]

Elkattawy HA, Mahmoud SM, Hassan AE et Al. Vagal Stimulation Ameliorates Non-Alcoholic Fatty Liver Disease in Rats *Biomedicines* 2023-12-08 [PMID: 38137476]

Dragojević T, Ivković E, Diklić M et Al. Hydroxyurea inhibits proliferation and stimulates apoptosis through inducible nitric oxide synthase in erythroid cells *Biomed Pharmacother* 2024-11-29 [PMID: 39615166]

Kaur, B;Miglioranza Scavuzzi, B;Yang, M;Yao, J;Jia, L;Abcouwer, SF;Zacks, DN; ER Stress and Mitochondrial Perturbations Regulate Cell Death in Retinal Detachment: Exploring the Role of HIF1? *Investigative ophthalmology & visual science* 2024-09-03 [PMID: 39325470]

Enes Akkaya, Şevket Evran, Fatih Çalış, Serdar Çevik, Salim Katar, Ersin Karataş, Abdurrahim Koçyiğit, Mustafa Yasin Sağlam, Mustafa Aziz Hatiboğlu, Hakan Hanımoğlu, Mehmet Yaşar Kaynar Thymoquinone ameliorates delayed cerebral injury and cerebral vasospasm secondary to experimental subarachnoid haemorrhage. *Neurologia i neurochirurgia polska* 2021-01-07 [PMID: 33252137]

More publications at <http://www.novusbio.com/NB100-56113>



### **Novus Biologicals USA**

10730 E. Briarwood Avenue  
Centennial, CO 80112  
USA  
Phone: 303.730.1950  
Toll Free: 1.888.506.6887  
Fax: 303.730.1966  
nb-customerservice@bio-techne.com

### **Bio-Techne Canada**

21 Canmotor Ave  
Toronto, ON M8Z 4E6  
Canada  
Phone: 905.827.6400  
Toll Free: 855.668.8722  
Fax: 905.827.6402  
canada.inquires@bio-techne.com

### **Bio-Techne Ltd**

19 Barton Lane  
Abingdon Science Park  
Abingdon, OX14 3NB, United Kingdom  
Phone: (44) (0) 1235 529449  
Free Phone: 0800 37 34 15  
Fax: (44) (0) 1235 533420  
info.EMEA@bio-techne.com

### **General Contact Information**

www.novusbio.com  
Technical Support: nb-technical@bio-techne.com  
Orders: nb-customerservice@bio-techne.com  
General: novus@novusbio.com

### **Products Related to NB100-56113**

---

NBP3-11853	Jurkat Staurosporine Treated / Untreated Cell Lysate
NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

---

### **Limitations**

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit [www.novusbio.com/guarantee](http://www.novusbio.com/guarantee)

Earn gift cards/discounts by submitting a review: [www.novusbio.com/reviews/submit/NB100-56113](http://www.novusbio.com/reviews/submit/NB100-56113)

Earn gift cards/discounts by submitting a publication using this product:  
[www.novusbio.com/publications](http://www.novusbio.com/publications)

