

Product Datasheet

CCR10/GPR2 Antibody NB100-56319

Unit Size: 0.1 mg

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 7

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB100-56319

Updated 9/9/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB100-56319




NB100-56319

CCR10/GPR2 Antibody

Product Information	
Unit Size	0.1 mg
Concentration	0.5 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.05% Sodium Azide
Isotype	IgG
Purity	Protein G purified
Buffer	PBS with 0.2% BSA
Product Description	
Description	Novus Biologicals Rabbit CCR10/GPR2 Antibody (NB100-56319) is a polyclonal antibody validated for use in IHC, WB, Flow and Simple Western. Anti-CCR10/GPR2 Antibody: Cited in 7 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	2826
Gene Symbol	CCR10
Species	Human
Reactivity Notes	Predicted to react with Mouse.
Immunogen	Rabbit anti-CCR10 polyclonal antibody was raised against a peptide corresponding to amino acids 350-362 of human CCR10. In this region, mouse and human amino acid sequences are 100% identical.
Product Application Details	
Applications	Western Blot, Simple Western, Flow Cytometry, Flow (Cell Surface), Flow (Intracellular), Immunohistochemistry, Immunohistochemistry-Frozen
Recommended Dilutions	Western Blot 1-2 ug/ml, Simple Western 1:50, Flow Cytometry 1:10-1:1000, Immunohistochemistry, Immunohistochemistry-Frozen 1:10-1:2000. Use reported in scientific literature (Vestergaard et al (2003)), Flow (Cell Surface), Flow (Intracellular) reported in scientific literature (PMID 15663561)
Application Notes	<p>In A375, a ~50 kDa band is observed. The CCR10 antibody is made against an epitope in the cytoplasmic domain of the receptor (Moed et al, 2004). For flow cytometry (intracellular) Moed et al fixed cells with 2% formaldehyde and permeabilized with 0.1% saponin.</p> <p>In Simple Western only 10 - 15 uL of the recommended dilution is used per data point. See Simple Western Antibody Database for Simple Western validation: Tested in MCF-7 lysate 0.5 mg/mL, separated by Size, antibody dilution of 1:50, apparent MW was 39 kDa. Separated by Size-Wes, Sally Sue/Peggy Sue.</p>

Images

Western Blot: CCR10/GPR2 Antibody [NB100-56319] - Analysis for GPR2/CCR10 using NB100-56319 at 2 ug/ml dilution against 10 ug of MCF-7 (a human breast cancer cell line) cell lysate.



Simple Western: CCR10/GPR2 Antibody [NB100-56319] - Simple Western lane view shows a specific band for CCR10 in 0.5 mg/ml of MCF-7 lysate. This experiment was performed under reducing conditions using the 12-230 kDa separation system.



Publications

T Karnezis, R Farnsworth, NC Harris, SP Williams, C Caesar, DJ Byrne, P Herle, ML Macheda, R Shayan, YF Zhang, S Yazar, SJ Takouridis, C Gerard, S Fox, MG Achen, SA Stacker CCL27/CCL28-CCR10 CHEMOKINE SIGNALING MEDIATES MIGRATION OF LYMPHATIC ENDOTHELIAL CELLS *Cancer Res.*, 2019-02-01;0(0):. 2019-02-01 [PMID: 30709930]

Moed H, Boorsma DM, Stoof TJ et al. Nickel-responding T cells are CD4+ CLA+ CD45RO+ and express chemokine receptors CXCR3, CCR4 and CCR10. *Br J Dermatol.* 2004-07-01 [PMID: 15270870] (Flow - IC, Human)

Details:

Flow (Intracellular): human PMBC from nickel-allergic patients (Fig 6).

Sell H, Dietze-Schroeder D, Kaiser U, Eckel J. Monocyte chemotactic protein-1 is a potential player in the negative cross-talk between adipose tissue and skeletal muscle. *Endocrinology.* 2006-05-01 [PMID: 16439461] (WB, Human)

Details:

WB: CCR4 (IMG-332) and CCR10 (IMG-286): human skeletal muscle cells and adipocytes (Fig 3).

Moed H, Boorsma DM, Tensen CP et al. Increased CCL27-CCR10 expression in allergic contact dermatitis: implications for local skin memory. *J Pathol.* 2004-09-01 [PMID: 15307136] (IHC-Fr, Human)

Details:

IHC-F (human skin biopsies from patients with allergic contact dermatitis): Tables 2,3; Fig. 5.

Rodrigues-Lisoni FC, Mehet DK, Peitl P Jr et al. In vitro and in vivo studies on CCR10 regulation by Annexin A1. *FEBS Lett.* 2006-02-20 [PMID: 16460738] (WB)

Details:

WB (human Hep-2 cells): Fig 3a.

Moed H, Stoof TJ, Boorsma DM et al. Identification of anti-inflammatory drugs according to their capacity to suppress type-1 and type-2 T cell profiles. *Clin Exp Allergy.* 2004-12-01 [PMID: 15663561] (Flow - IC, Human)

Details:

Flow (Intracellular): human PBMC from nickel-allergic patients (Figs 1B, 2B, 3B).

Vestergaard C, Deleuran M, Gesser B et al. Expression of the T-helper 2-specific chemokine receptor CCR4 on CCR10-positive lymphocytes in atopic dermatitis skin but not in psoriasis skin. *Br J Dermatol.* 2003-09-01 [PMID: 14510975] (IHC-Fr, Human)

Details:

IHC-F (human skin biopsies from patients with atopic dermatitis and psoriasis): Figs 1 and 2, Table 1.



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NB100-56319

NBL1-08888	CCR10/GPR2 Overexpression Lysate
NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-56319

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

