

Product Datasheet

TRAILR1/TNFRSF10A Antibody (32A242) - BSA Free NB100-56528

Unit Size: 0.1 mg

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 8

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB100-56528

Updated 9/9/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB100-56528



NB100-56528


TRAILR1/TNFRSF10A Antibody (32A242) - BSA Free

Product Information	
Unit Size	0.1 mg
Concentration	1.0 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	32A242
Preservative	0.05% Sodium Azide
Isotype	IgG1
Purity	Protein G purified
Buffer	PBS
Product Description	
Description	Novus Biologicals Mouse TRAILR1/TNFRSF10A Antibody (32A242) - BSA Free (NB100-56528) is a monoclonal antibody validated for use in WB and ICC/IF. Anti-TRAILR1/TNFRSF10A Antibody: Cited in 8 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Mouse
Gene ID	8797
Gene Symbol	TNFRSF10A
Species	Human, Mouse, Rat
Immunogen	This monoclonal antibody was raised against a peptide corresponding to amino acid 2 to 21 of human DR4.
Product Application Details	
Applications	Western Blot, Immunoblotting, Immunocytochemistry/ Immunofluorescence
Recommended Dilutions	Western Blot 1 - 3 ug/ml, Immunocytochemistry/ Immunofluorescence reported in scientific literature (PMID 23988408), Immunoblotting reported in scientific literature (PMID 31534206)
Application Notes	In HeLa, a 57 kDa band should be observed.




Images

Western Blot: TRAIL R1/TNFRSF10A Antibody (32A242) [NB100-56528]
- Analysis using Azide/BSA FREE version of NB100-56528. Analysis of 20 ug of total cell lysate from HeLa cells with DR4 antibody at 2 ug/ml. Goat anti-mouse Ig HRP secondary antibody and PicoTect ECL substrate solution were used for this test.



Western Blot: TRAIL R1/TNFRSF10A Antibody (32A242) [NB100-56528]
- Analysis of 20 ug of total cell lysate from HeLa cells with DR4 antibody at 2 ug/ml. Goat anti-mouse Ig HRP secondary antibody and PicoTect ECL substrate solution were used for this test.



Publications

Lee SR, Quan KT, Byun HS et al. Accelerated degradation of cFLIPL and sensitization of the TRAIL DISC-mediated apoptotic cascade by pinoresinol, a lignan isolated from *Rubia philippinensis* Sci Rep 2019-09-18 [PMID: 31534206] (IB, Human)

Wen Q, Wang W, Luo J et al. CGP57380 enhances efficacy of RAD001 in non-small cell lung cancer through abrogating mTOR inhibition-induced phosphorylation of eIF4E and activating mitochondrial apoptotic pathway. Oncotarget. 2016-03-30 [PMID: 27050281] (WB, Human)

Di X, Zhang G, Zhang Y et al. Accumulation of autophagosomes in breast cancer cells induces TRAIL resistance through downregulation of surface expression of death receptors 4 and 5. Oncotarget 2013-09-01 [PMID: 23988408] (ICC/IF)

Brooks AD, Jacobsen KM, Li W et al. Bortezomib sensitizes human renal cell carcinomas to TRAIL apoptosis through increased activation of caspase-8 in the death-inducing signaling complex. Mol Cancer Res. 2010-05-01 [PMID: 20442297]

Xu J, Zhou JY, Wu GS. Tumor necrosis factor-related apoptosis-inducing ligand is required for tumor necrosis factor alpha-mediated sensitization of human breast cancer cells to chemotherapy. Cancer Res. 2006-10-15 [PMID: 17047073]

Laguinge LM, Samara RN, Wang W et al. DR5 receptor mediates anoikis in human colorectal carcinoma cell lines. Cancer Res. 2008-02-01 [PMID: 18245494]

Details:

WB [human colorectal cancer (CRC) cells], Fig. 4A.

Cisternas P, Moreno RD. Comparative analysis of apoptotic pathways in rat, mouse, and hamster spermatozoa. Mol Reprod Dev. 2006-10-01 [PMID: 16868928] (WB)

Details:

DR4/TRAILR1 (NT) (IMG-275). WB: Mouse, rat, and hamster spermatozoa, Fig 3A. Note: The DR4/TRAIL1 antibody was detected in mouse and rat spermatozoa but not in hamster, Fig 3A.

Jin F, Liu X, Zhou Z et al. Activation of nuclear factor-kappaB contributes to induction of death receptors and apoptosis by the synthetic retinoid CD437 in DU145 human prostate cancer cells. Cancer Res. 2005-07-15 [PMID: 16024638] (WB)

Details:

IMGENEX antibodies cited for WB: 1. Caspase-3 mAb, clone 31A1067 (IMG-144A): Fig 3C, DU145 prostate adenocarcinoma cells. 2. DcR1 pAb (IMG-245-1/IMG-245-2): Fig 6C, DcR1 overexpressing DU145 cells. Note: The specificity of the DcR1 pAb was validated in Dc



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NB100-56528

NB800-PC1	HeLa Whole Cell Lysate
NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB7539	Goat anti-Mouse IgG (H+L) Secondary Antibody [HRP]
NBP1-97005-0.5mg	Mouse IgG1 Isotype Control (MG1)

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-56528

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

