

Product Datasheet

I κ B-alpha [p Ser32, p Ser36] Antibody (39A1413) - BSA Free **NB100-56724**

Unit Size: 0.1 mg

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 13

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB100-56724

Updated 9/9/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB100-56724



NB100-56724I κ B-alpha [p Ser32, p Ser36] Antibody (39A1413) - BSA Free


Product Information	
Unit Size	0.1 mg
Concentration	1.0 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	39A1413
Preservative	0.05% Sodium Azide
Isotype	IgG1 Kappa
Purity	Protein G purified
Buffer	PBS
Product Description	
Description	Novus Biologicals Mouse I κ B-alpha [p Ser32, p Ser36] Antibody (39A1413) - BSA Free (NB100-56724) is a monoclonal antibody validated for use in WB, ICC/IF, Simple Western and IP. Anti-I κ B-alpha Antibody: Cited in 13 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Mouse
Gene ID	4792
Gene Symbol	NFKBIA
Species	Human, Mouse, Rat, Porcine, Bovine, Canine
Immunogen	This monoclonal antibody was raised against a synthetic peptide containing phosphorylated serines at amino acid residues 32 and 36 of human I κ B α .
Product Application Details	
Applications	Western Blot, Simple Western, Immunoprecipitation
Recommended Dilutions	Western Blot 1-2 ug/ml, Simple Western 1:50, Immunoprecipitation 1-2 ug/ml
Application Notes	See Simple Western Antibody Database for Simple Western validation: tested in MCF10A-/+EGF, Jurkat-/+TNF, HEK293-/+UV; antibody dilution of 1:50; separated by charge.

Images

Western Blot: I κ B-alpha [p Ser32, p Ser36] Antibody (39A1413) [NB100-56724] - Analysis using the Biotin conjugate of NB100-56724. Detection of Jurkat cells were treated for 30 min with 100 ug/ml ALLN (N-Acetyl-Leu-Leu-Norleucinal; a Calpain inhibitor and also proteasome inhibitor that prevents I κ Ba dephosphorylation) followed by incubation with (lanes 2 & 4) or without 1 nM TNF-a (1 & 3). The membranes were blotted with this antibody.



Western Blot: I κ B-alpha [p Ser32, p Ser36] Antibody (39A1413) [NB100-56724] - Jurkat cells were treated for 30 min with 100 ug/ml ALLN (N-Acetyl-Leu-Leu-Norleucinal) - a Calpain inhibitor and also proteasome inhibitor that prevents I κ Ba dephosphorylation) followed by incubation with (lane 2) or without 1 nM TNF-a (lane 1). The membrane was blotted with NB100-56724 and immunoreactivity was detected by ECL. The data shows that NB100-56724 detects specifically the phosphorylated form of I κ Ba.



Publications

Shahbazi S, Zakerali T The inhibitory role of benzo-dioxole-piperamide on the phosphorylation process as an NF-Kappa B silencer *Biomedicine & pharmacotherapy = Biomedecine & pharmacotherapie* 2021-11-28 [PMID: 34852990] (ICC/IF, WB, Human)

Wen ZS, Tang Z, Gu LX et al. Immunomodulatory effect of low molecular-weight seleno-aminopolysaccharide on immunosuppressive mice. *Int. J. Biol. Macromol.* 2018-10-17 [PMID: 30342138] (WB, Mouse)

Kanno Y, Maruyama C, Matsuda A, Ishisaki A. uPA-derived peptide, A6 is involved in the suppression of lipopolysaccharide-promoted inflammatory osteoclastogenesis and the resultant bone loss. *Immun Inflamm Dis* 2017-05-11 [PMID: 28493442] (WB, Mouse)

Gu LX, Wen ZS, Xiang XW et al. Immunomodulatory effect of low molecular-weight seleno-aminopolysaccharides in intestinal epithelial cells. *Int. J. Biol. Macromol.* 2017-03-04 [PMID: 28267616] (WB, Porcine)

Lotz Sonja, Aga Eresso, Wilde Inga et al. Highly purified lipoteichoic acid activates neutrophil granulocytes and delays their spontaneous apoptosis via CD14 and TLR2. *J Leukoc Biol.* 2004-03-01 [PMID: 14673018]

Shakibaei Mehdi, Allaway David, Nebrich Simone, Mobasheri Ali. Botanical Extracts from Rosehip (*Rosa canina*), Willow Bark (*Salix alba*), and Nettle Leaf (*Urtica dioica*) Suppress IL-1beta-Induced NF-kappaB Activation in Canine Articular Chondrocytes. *Evid Based Complement Alternat Med.* 2012-01-01 [PMID: 22474508]

Samanta D, Gilkes D, Chaturvedi P, Xiang L, Semenza G. Hypoxia-inducible factors are required for chemotherapy resistance of breast cancer stem cells. *Proc Natl Acad Sci U S A.* 2014-12-01 [PMID: 25453096] (WB, Human)

Das U, Manna K, Sinha M et al. Role of ferulic Acid in the amelioration of ionizing radiation induced inflammation: a murine model. *PLoS One* 2014-05-24 [PMID: 24854039] (WB, Mouse)

Jazirehi AR, Huerta-Yepez S, Cheng G et al. Rituximab (chimeric anti-CD20 monoclonal antibody) inhibits the constitutive nuclear factor- κ B signaling pathway in non-Hodgkin's lymphoma B-cell lines: role in sensitization to chemotherapeutic drug-induced apoptosis. *Cancer Res.* 2005-01-01 [PMID: 15665303] (WB)

Details:

WB (Ramos and Daudi cells growing in RPMI 1640 and 1% fetal bovine serum and rituximab), Fig. 4a

Ikezoe T, Yang Y, Heber D et al. PC-SPES: a potent inhibitor of nuclear factor-kappa B rescues mice from lipopolysaccharide-induced septic shock. *Mol Pharmacol.* 2003-12-01 [PMID: 14645683]

Han HS, Karabiyikoglu M, Kelly S et al. Mild hypothermia inhibits nuclear factor-kappaB translocation in experimental stroke. *J Cereb Blood Flow Metab.* 2003-05-01 [PMID: 12771574]

Details:

Products cited for western blot in male Sprague-Dawley rats ischemic cortex, ischemic subcortex, contralateral cortex, and contralateral subcortex tissue. The rats were subject to mild and delayed hypothermia. 1. IKK-gamma (IMG-324A): Fig 4A & 4B. 2. I κ B-

Li Q, Guo Y, Tan W et al. Cardioprotection afforded by inducible nitric oxide synthase gene therapy is mediated by cyclooxygenase-2 via a nuclear factor-kappaB dependent pathway. *Circulation.* 2007-10-02 [PMID: 17785622] (WB, Mouse)

Details:

WB (WT, COX-2^{-/-}, I κ B α (Ser 32A, Ser 36A) Tg mice injected with iNOS gene), Fig. 5.

More publications at <http://www.novusbio.com/NB100-56724>



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NB100-56724

NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB7539	Goat anti-Mouse IgG (H+L) Secondary Antibody [HRP]
NBP1-43319-0.5mg	Mouse IgG1 Kappa Isotype Control (P3.6.2.8.1)

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-56724

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

