

Product Datasheet

D4-GDI/RhoGDI2 Antibody (97A1015) - (cleavage specific) - BSA Free NB100-56725

Unit Size: 0.1 mg

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Reviews: 1 Publications: 16

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB100-56725

Updated 9/9/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB100-56725



NB100-56725

D4-GDI/RhoGDI2 Antibody (97A1015) - (cleavage specific) - BSA Free


Product Information	
Unit Size	0.1 mg
Concentration	1.0 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	97A1015
Preservative	0.05% Sodium Azide
Isotype	IgG2b Kappa
Purity	Protein G purified
Buffer	PBS

Product Description	
Description	Novus Biologicals Mouse D4-GDI/RhoGDI2 Antibody (97A1015) - (cleavage specific) - BSA Free (NB100-56725) is a monoclonal antibody validated for use in IHC, WB, Flow and IP. Anti-D4-GDI/RhoGDI2 Antibody: Cited in 16 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Mouse
Gene ID	397
Gene Symbol	ARHGDI2
Species	Human, Mouse
Immunogen	A synthetic peptide corresponding to the region of the Fas-induced cleavage site of human D4-GDI/RhoGDI2 was used as the immunogen.


Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, Flow Cytometry, Flow (Intracellular), Immunohistochemistry, Immunoprecipitation
Recommended Dilutions	Western Blot 2 ug/ml, Flow Cytometry 0.5-1 ug/10 ⁶ cells, Immunohistochemistry 1:10-1:500, Immunoprecipitation 1-2 ug/ml, Immunohistochemistry-Paraffin 1:10-1:500, Flow (Intracellular) 0.5-1 ug/10 ⁶ cells
Application Notes	In anti-Fas treated Jurkat, an approx. 23 kDa band should be observed.

Images


Western Blot: D4-GDI/RhoGDI2 Antibody (97A1015) - (cleavage specific) [NB100-56725] - WB analysis: HL60 cells were treated for different time periods with 10 ng/ml of anti-Fas mAb (NB100-56725, Clone DX2). Harvested cells were lysed and resolved by SDS-PAGE and transferred onto membrane. D4-GDI/RhoGDI2 cleavage was detected.




Immunohistochemistry-Paraffin: D4-GDI/RhoGDI2 Antibody (97A1015) - (cleavage specific) [NB100-56725] - analysis of D4GD1-cv in formalin-fixed, paraffin-embedded human breast tumor tissue using an isotype control (top) and this antibody.




Flow (Intracellular): D4-GDI/RhoGDI2 Antibody (97A1015) - (cleavage specific) [NB100-56725] - Intracellular analysis of cleaved D4-GDI in 10^6 untreated (left) and 3 hour staurosporine-treated (right) Jurkat cells using 0.5 ug of this antibody was used for this test. Green histogram represents isotype control; red represents anti-D4-GDI antibody. Image using the Alexa Fluor 647 format of this antibody.




Western Blot: D4-GDI/RhoGDI2 Antibody (97A1015) - (cleavage specific) [NB100-56725] - Western blot analysis of D4GDI in Jurkat cells using NB100-56556 (IMG-396A) at 2 ug/ml and NB100-56725 (IMG-160A) at 0.1 ug/ml. Cells were treated with 2 uM staurosporine for different time periods. NB100-56556 (IMG-396A) detects both the full-length and cleaved forms of D4GDI. While (NB100-56725) IMG-160A specifically detects only the cleaved form of the protein.




Flow (Intracellular): D4-GDI/RhoGDI2 Antibody (97A1015) - (cleavage specific) [NB100-56725] - HL60 cells were left untreated (green) or treated with camptothecin to induce apoptosis (red). Cells were analyzed by intracellular flow cytometry using either the (0.5 ug antibody/test/1 million cells) and ((as seen in B). this is because (cells not undergoing apoptosis) and cleaved (apoptotic cells) forms of D4-GDI.



Flow (Intracellular): D4-GDI/RhoGDI2 Antibody (97A1015) - (cleavage specific) [NB100-56725] - Analysis using the FITC conjugate of NB100-56725. Staining of cleaved D4-GDI in 10^6 camptothecin-treated (red) and untreated (green) HL60 cells using 0.5 ug of NB100-56060. The shaded histogram represents cells without antibody.



Flow (Intracellular): D4-GDI/RhoGDI2 Antibody (97A1015) - (cleavage specific) [NB100-56725] - Analysis using the FITC conjugate of NB100-56725. Staining of cleaved D4-GDI in untreated (left) and staurosporine-treated (right) Ramos cells using 0.5 ug of was used for this test.



Publications

Yamaguchi H, Wang HG. The protein kinase PKB/Akt regulates cell survival and apoptosis by inhibiting Bax conformational change. *Oncogene*. 2001-11-22 [PMID: 11753656]

Chen YC, Shen SC, Lin HY. Rutinoside at C7 attenuates the apoptosis-inducing activity of flavonoids. *Biochem Pharmacol*. 2003-10-01 [PMID: 14505793]

Gildea JJ, Seraj MJ, Oxford G et al. RhoGDI2 is an invasion and metastasis suppressor gene in human cancer. *Cancer Res*. 2002-11-15 [PMID: 12438227]

Jenkins CE, Swiatoniowski A, Power MR, Lin TJ. *Pseudomonas aeruginosa*-induced human mast cell apoptosis is associated with up-regulation of endogenous Bcl-xS and down-regulation of Bcl-xL. *J Immunol*. 2006-12-01 [PMID: 17114473]

Jenkins CE, Swiatoniowski A, Issekutz AC, Lin TJ. *Pseudomonas aeruginosa* exotoxin A induces human mast cell apoptosis by a caspase-8 and -3-dependent mechanism. *J Biol Chem*. 2004-08-27 [PMID: 15205454]

Cuddeback SM, Yamaguchi H, Komatsu K et al. Molecular cloning and characterization of Bif-1. A novel Src homology 3 domain-containing protein that associates with Bax. *J Biol Chem*. 2001-06-08 [PMID: 11259440]

Liu CY, Lee CF, Hong CH, Wei YH. Mitochondrial DNA mutation and depletion increase the susceptibility of human cells to apoptosis. *Ann N Y Acad Sci*. 2004-04-01 [PMID: 15126291]

Sato S, Fujita N, Tsuruo T. Interference with PDK1-Akt survival signaling pathway by UCN-01 (7-hydroxystaurosporine). *Oncogene*. 2002-03-07 [PMID: 11896604]

Ko CH, Shen SC, Chen YC. Hydroxylation at C4' or C6 is essential for apoptosis-inducing activity of flavanone through activation of the caspase-3 cascade and production of reactive oxygen species. *Free Radic Biol Med*. 2004-04-01 [PMID: 15019974]

Yamaguchi H, Paranawithana SR, Lee MW et al. Epothilone B analogue (BMS-247550)-mediated cytotoxicity through induction of Bax conformational change in human breast cancer cells. *Cancer Res*. 2002-01-15 [PMID: 11809697]

Chen YC, Shen SC, Lee WR et al. Emodin induces apoptosis in human promyeloleukemic HL-60 cells accompanied by activation of caspase 3 cascade but independent of reactive oxygen species production. *Biochem Pharmacol*. 2002-12-15 [PMID: 12445860]

Okamoto M, Koga S, Tatsuka M. Differential regulation of caspase-9 by ionizing radiation- and UV-induced apoptotic pathways in thymic cells. *Mutat Res*. 2010-06-01 [PMID: 20346366] (WB, Mouse)

Details:

D4-GDI (IMG-160A). WB: Mouse lymphoma 3SB cell line subject to either ionizing radiation or UV light, Fig 2D.

More publications at <http://www.novusbio.com/NB100-56725>





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NB100-56725

NBL1-07674	D4-GDI/RhoGDI2 Overexpression Lysate
NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB7539	Goat anti-Mouse IgG (H+L) Secondary Antibody [HRP]
NBP1-43317-0.5mg	Mouse IgG2b Kappa Light Chain Isotype Control (MG2b)

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-56725

Earn gift cards/discounts by submitting a publication using this product:

www.novusbio.com/publications

