

# Product Datasheet

## CTLA-4 Antibody (BNI3) - BSA Free NB100-64849-0.1mg

Unit Size: 0.1 mg

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

[www.novusbio.com](http://www.novusbio.com)



[technical@novusbio.com](mailto:technical@novusbio.com)

**Reviews: 1 Publications: 4**

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:  
[www.novusbio.com/NB100-64849](http://www.novusbio.com/NB100-64849)

Updated 9/9/2025 v.20.1

**Earn rewards for product  
reviews and publications.**

Submit a publication at [www.novusbio.com/publications](http://www.novusbio.com/publications)

Submit a review at [www.novusbio.com/reviews/destination/NB100-64849](http://www.novusbio.com/reviews/destination/NB100-64849)



**NB100-64849-0.1mg**


CTLA-4 Antibody (BNI3) - BSA Free

Product Information	
Unit Size	0.1 mg
Concentration	1.0 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	BNI3
Preservative	0.09% Sodium Azide
Isotype	IgG2a
Purity	Protein A purified
Buffer	PBS
Product Description	
Description	Novus Biologicals Mouse CTLA-4 Antibody (BNI3) - BSA Free (NB100-64849) is a monoclonal antibody validated for use in IHC, ELISA, Flow and IP. Anti-CTLA-4 Antibody: Cited in 4 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Mouse
Gene ID	1493
Gene Symbol	CTLA4
Species	Human
Specificity/Sensitivity	NB100-64849 recognizes the human CD152 cell surface antigen, also known as CTLA-4 (cytotoxic T-lymphocyte-associated antigen 4). CD152 is a 45kD glycoprotein, expressed normally as a disulphide-linked homodimer. It is expressed transiently on the surface of T cells following activation (peaking at 3 days), but is often present with the cytoplasm. CD152 is a ligand for CD80 and CD86, and is important in cellular interactions during the immune response.
Immunogen	Human CTLA-4/human IgG heavy chain fusion protein
Product Application Details	
Applications	Immunohistochemistry-Paraffin, ELISA, Flow Cytometry, Immunohistochemistry, Immunohistochemistry-Frozen, Immunoprecipitation, Immunofluorescence
Recommended Dilutions	Flow Cytometry Neat - 1:10, ELISA, Immunohistochemistry 1:10 - 1:500, Immunoprecipitation 1:10 - 1:500, Immunohistochemistry-Paraffin, Immunohistochemistry-Frozen 5 ug/mL - 50 ug/mL, Immunofluorescence
Application Notes	Use in IHC-P reported in scientific literature (PMID:34513744). Membrane permeabilization is required for Flow Cytometry. The epitope recognised by this antibody is reported to be sensitive to formaldehyde fixation and tissue processing. The use of acetone fixation for frozen sections is recommended.

## Images

### ELISA: CTLA-4 Antibody (BNI3) [NB100-64849] - Human Anti-Abatacept Antibody bridging ELISA for Pharmacokinetic (PK) assay development

A microtiter plate was coated overnight with Mouse Anti-Human CD152 Antibody, clone BNI3 at a concentration of 1 µg/ml. After washing and blocking with PBST + 5% BSA, 10% human serum was added spiked with increasing concentrations of abatacept. Detection was performed using HRP conjugated Human Anti-Abatacept Antibody, clone AbD37058ia at a concentration of 2 µg/ml in HISPEC Assay Diluent followed by QuantaBlu Fluorogenic Peroxidase Substrate. Data are shown as the mean of three measurements. HRP conjugation of the antibody used in this assay was performed using a LYNX Rapid HRP Antibody Conjugation Kit.



## Publications

Lu B, Lu T, Shi J et al. Basic Transcription Factor 3 Like 4 Enhances Malignant Phenotypes through Modulating Tumor Cell Function and Immune Microenvironment in Glioma. *The American journal of pathology* 2024-04-22 [PMID: 38320629]

Dai T, Ye L, Yu H et al. Regulation Network and Prognostic Significance of Aldo-Keto Reductase (AKR) Superfamily Genes in Hepatocellular Carcinoma *Journal of hepatocellular carcinoma* 2021-08-30 [PMID: 34513744] (IHC-P, Human)

Jiang S, Zhang Y, Zhang X Et al. GARP Correlates With Tumor-Infiltrating T-Cells and Predicts the Outcome of Gastric Cancer *Frontiers in immunology* 2021-08-06 [PMID: 34421887] (IHC-P, Human)

Santoni G, Amantini C, Morelli MB et al. High CTLA-4 expression correlates with poor prognosis in thymoma patients. *Oncotarget* 2018-03-30 [PMID: 29682176] (IF/IHC, Human)



### **Novus Biologicals USA**

10730 E. Briarwood Avenue  
Centennial, CO 80112  
USA  
Phone: 303.730.1950  
Toll Free: 1.888.506.6887  
Fax: 303.730.1966  
nb-customerservice@bio-techne.com

### **Bio-Techne Canada**

21 Canmotor Ave  
Toronto, ON M8Z 4E6  
Canada  
Phone: 905.827.6400  
Toll Free: 855.668.8722  
Fax: 905.827.6402  
canada.inquires@bio-techne.com

### **Bio-Techne Ltd**

19 Barton Lane  
Abingdon Science Park  
Abingdon, OX14 3NB, United Kingdom  
Phone: (44) (0) 1235 529449  
Free Phone: 0800 37 34 15  
Fax: (44) (0) 1235 533420  
info.EMEA@bio-techne.com

### **General Contact Information**

www.novusbio.com  
Technical Support: nb-technical@bio-techne.com  
Orders: nb-customerservice@bio-techne.com  
General: novus@novusbio.com

### **Products Related to NB100-64849-0.1mg**

---

HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB7539	Goat anti-Mouse IgG (H+L) Secondary Antibody [HRP]
NBP1-96778	Mouse IgG2a Isotype Control (M2A)
NBP2-22987	Recombinant Human CTLA-4 His Protein

---

### **Limitations**

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit [www.novusbio.com/guarantee](http://www.novusbio.com/guarantee)

Earn gift cards/discounts by submitting a review: [www.novusbio.com/reviews/submit/NB100-64849](http://www.novusbio.com/reviews/submit/NB100-64849)

Earn gift cards/discounts by submitting a publication using this product:  
[www.novusbio.com/publications](http://www.novusbio.com/publications)

