

# Product Datasheet

## Alix Antibody (3A9) - BSA Free NB100-65678

Unit Size: 0.1 mg

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

[www.novusbio.com](http://www.novusbio.com)



[technical@novusbio.com](mailto:technical@novusbio.com)

### Publications: 12

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:  
[www.novusbio.com/NB100-65678](http://www.novusbio.com/NB100-65678)

Updated 9/9/2025 v.20.1

Earn rewards for product  
reviews and publications.

Submit a publication at [www.novusbio.com/publications](http://www.novusbio.com/publications)

Submit a review at [www.novusbio.com/reviews/destination/NB100-65678](http://www.novusbio.com/reviews/destination/NB100-65678)



**NB100-65678**


Alix Antibody (3A9) - BSA Free

<b>Product Information</b>	
<b>Unit Size</b>	0.1 mg
<b>Concentration</b>	1.0 mg/ml
<b>Storage</b>	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
<b>Clonality</b>	Monoclonal
<b>Clone</b>	3A9
<b>Preservative</b>	0.09% Sodium Azide
<b>Isotype</b>	IgG1
<b>Purity</b>	Protein A purified
<b>Buffer</b>	PBS
<b>Product Description</b>	
<b>Description</b>	Novus Biologicals Mouse Alix Antibody (3A9) - BSA Free (NB100-65678) is a monoclonal antibody validated for use in IHC, WB, ELISA, Flow and IP. Anti-Alix Antibody: Cited in 11 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
<b>Host</b>	Mouse
<b>Gene ID</b>	10015
<b>Gene Symbol</b>	PDCD6IP
<b>Species</b>	Human
<b>Reactivity Notes</b>	Predicted cross-reactivities: Rat, Xenopus, Mouse, Human Please note that this antibody is reactive to Mouse and derived from the same host, Mouse. Additional Mouse on Mouse blocking steps may be required for IHC and ICC experiments. Please contact Technical Support for more information.
<b>Specificity/Sensitivity</b>	NB100-65678 recognizes apoptotic-linked-gene-product 2 (ALG-2) interacting protein X (ALIX), a conserved adaptor protein which is ubiquitously expressed. Alix was originally reported to play a role in apoptosis but has recently been shown to be involved in other cellular mechanisms including endosomal sorting, endocytosis, viral budding and actin cytoskeleton assembly.
<b>Immunogen</b>	Alix-GST fusion protein
<b>Product Application Details</b>	
<b>Applications</b>	Western Blot, Immunohistochemistry-Paraffin, ELISA, Flow Cytometry, Immunohistochemistry, Immunoprecipitation
<b>Recommended Dilutions</b>	Western Blot 1:100-1:2000, Flow Cytometry, ELISA 1:100-1:2000, Immunohistochemistry 1:10-1:500, Immunoprecipitation 1:10-1:500, Immunohistochemistry-Paraffin 1:10-1:500
<b>Application Notes</b>	Use in Flow Cytometry reported in scientific literature (PMID:34298673). For IHC - This product does not require protein digestion pre-treatment of paraffin sections.




## Images


Western Blot: Alix Antibody (3A9) [NB100-65678] - B. Western blot for the exosomal marker Alix and Nef in cells and exosomes (exNef). Image collected and cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/31344124>) licensed under a CC-BY license.



Western Blot: Alix Antibody (3A9) [NB100-65678] - Western blot analysis of HeLa whole cell lysate probed with MOUSE ANTI ALIX F(ab')<sub>2</sub> Rabbit anti Mouse IgG:HRP (STAR13B)



Western Blot: Alix Antibody (3A9) [NB100-65678] - Nef-containing exosomes deliver Nef to macrophages. A—Size distribution of the extracellular vesicles secreted by HEK293 cells determined by EM; Inset—EM micrograph of the vesicles; bar—200 nm. B—Western blot for the exosomal marker Alix & Nef in cells & exosomes (exNef); C—Western blot for the indicated amounts of rNef & in a typical preparation of exNef (10 µg of exosomal protein); D, E—Time-course of exosome uptake quantitated by confocal microscopy; F—Time-course of exosome uptake quantitated by fluorimetry; percentage of added exosomes that was taken up is shown; G—Cells were incubated with exosomes for 48 h, excess exosomes was washed out & cells incubated for the indicated periods of time in exosome-free medium; retained fluorescence of the exosome stain PKH67 was assessed using confocal microscopy; H—Visualisation of Nef-GFP inside the cells after exposure to exNef-GFP (5 µg/ml of exosomal protein) after staining with anti-GFP antibody. Scale bars—10 µm. Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/31344124>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



## Publications

Han, G;Kim, H;Jang, H;Kim, ES;Kim, SH;Yang, Y; Oral TNF- $\alpha$  siRNA delivery via milk-derived exosomes for effective treatment of inflammatory bowel disease *Bioactive materials* 2024-04-01 [PMID: 38223538]

Singh R, Santos M, Herndon C et al. Detection by super-resolution microscopy of viral proteins inside bloodborne extracellular vesicles *Extracellular Vesicles and Circulating Nucleic Acids* 2023-11-08

Eslami-S Z, Cortés-Hernández LE, Sinoquet L et al. Circulating tumour cells and PD-L1-positive small extracellular vesicles: the liquid biopsy combination for prognostic information in patients with metastatic non-small cell lung cancer *British journal of cancer* 2023-11-16 [PMID: 37973956]

Han G, Cho H, Kim H et al. Bovine colostrum derived-exosomes prevent dextran sulfate sodium-induced intestinal colitis via suppression of inflammation and oxidative stress *Biomaterials science* 2022-04-12 [PMID: 35315847]

Kim H, Jang Y, Kim E et al. Potential of Colostrum-Derived Exosomes for Promoting Hair Regeneration Through the Transition From Telogen to Anagen Phase *Frontiers in Cell and Developmental Biology* 2022-03-10 [PMID: 35359449] (WB, Bovine)

Kim H, Kim DE, Han G et al. Harnessing the Natural Healing Power of Colostrum: Bovine Milk-Derived Extracellular Vesicles from Colostrum Facilitating the Transition from Inflammation to Tissue Regeneration for Accelerating Cutaneous Wound Healing *Advanced healthcare materials* 2021-12-05 [PMID: 34865307] (WB, Mouse)

Yin GN, Shin TY, Ock J et al. Pericyte derived extracellular vesicles mimetic nanovesicles improves peripheral nerve regeneration in mouse models of sciatic nerve transection *International journal of molecular medicine* 2022-02-01 [PMID: 34935051] (WB, Mouse)

Sadri M, Hirose N, Le J et al. Tumor necrosis factor receptor-1 is selectively sequestered into Schwann cell extracellular vesicles where it functions as a TNF  $\alpha$  decoy *Glia* 2021-09-24 [PMID: 34559433]

Rabe Dc, Walker Nd, Rustandy Fd Et Al. Tumor Extracellular Vesicles Regulate Macrophage-Driven Metastasis through CCL5 *Cancers* 2021-07-10 [PMID: 34298673] (FLOW)

Mukhamedova N, Hoang A, Dragoljevic D et al. Exosomes containing HIV protein Nef reorganize lipid rafts potentiating inflammatory response in bystander cells *PLoS Pathog.* 2019-07-01 [PMID: 31344124] (WB, Mouse, Human)

Mariscal J, Fernandez-Puente P, Calamia V et al. Proteomic characterization of epithelial-like extracellular vesicles in advanced endometrial cancer. *J. Proteome Res.* 2018-12-26 [PMID: 30585730] (WB, Human)

Haga H, Yan IK, Takahashi K et al. Extracellular Vesicles from Bone Marrow-Derived Mesenchymal Stem Cells Improve Survival from Lethal Hepatic Failure in Mice. *Stem Cells Transl Med.* 2017-02-18 [PMID: 28213967] (Mouse)



### **Novus Biologicals USA**

10730 E. Briarwood Avenue  
Centennial, CO 80112  
USA  
Phone: 303.730.1950  
Toll Free: 1.888.506.6887  
Fax: 303.730.1966  
nb-customerservice@bio-techne.com

### **Bio-Techne Canada**

21 Canmotor Ave  
Toronto, ON M8Z 4E6  
Canada  
Phone: 905.827.6400  
Toll Free: 855.668.8722  
Fax: 905.827.6402  
canada.inquires@bio-techne.com

### **Bio-Techne Ltd**

19 Barton Lane  
Abingdon Science Park  
Abingdon, OX14 3NB, United Kingdom  
Phone: (44) (0) 1235 529449  
Free Phone: 0800 37 34 15  
Fax: (44) (0) 1235 533420  
info.EMEA@bio-techne.com

### **General Contact Information**

www.novusbio.com  
Technical Support: nb-technical@bio-techne.com  
Orders: nb-customerservice@bio-techne.com  
General: novus@novusbio.com

### **Products Related to NB100-65678**

---

NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB7539	Goat anti-Mouse IgG (H+L) Secondary Antibody [HRP]
NBP1-97005-0.5mg	Mouse IgG1 Isotype Control (MG1)

---

### **Limitations**

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit [www.novusbio.com/guarantee](http://www.novusbio.com/guarantee)

Earn gift cards/discounts by submitting a review: [www.novusbio.com/reviews/submit/NB100-65678](http://www.novusbio.com/reviews/submit/NB100-65678)

Earn gift cards/discounts by submitting a publication using this product:  
[www.novusbio.com/publications](http://www.novusbio.com/publications)

