

Product Datasheet

Actin Antibody (mAbGEa) - BSA Free NB100-74340

Unit Size: 0.1 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Reviews: 8 Publications: 34

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB100-74340

Updated 9/9/2025 v.20.1

**Earn rewards for product
reviews and publications.**

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB100-74340




NB100-74340

Actin Antibody (mAbGEa) - BSA Free


Product Information	
Unit Size	0.1 ml
Concentration	1.0 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	mAbGEa
Preservative	0.05% Sodium Azide
Isotype	IgM Kappa
Purity	IgM purified
Buffer	PBS
Product Description	
Description	Novus Biologicals Mouse Actin Antibody (mAbGEa) - BSA Free (NB100-74340) is a monoclonal antibody validated for use in IHC, WB, ELISA, Flow, ICC/IF and Simple Western. Anti-Actin Antibody: Cited in 31 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Mouse
Gene ID	58
Gene Symbol	ACTA1
Species	Human, Mouse, Rat, Bovine, Chinese Hamster, Drosophila, Fungi, Plant, Protozoa, Rabbit, Sheep, Xenopus, Yeast, Zebrafish
Reactivity Notes	Chinese hamster data from customer review. Sordaria macrospora reactivity reported in scientific literature (PMID: 24720701).
Specificity/Sensitivity	Reacts with Actin 1, 2, 3, 4, 7, 8, 11 and 12.
Immunogen	Purified recombinant Arabidopsis Actin protein [UniProt# P0CJ46]
Product Application Details	
Applications	Western Blot, Simple Western, Immunohistochemistry-Paraffin, ELISA, Flow Cytometry, Flow (Intracellular), Immunocytochemistry/ Immunofluorescence, Immunohistochemistry
Recommended Dilutions	Western Blot 1:100-1:1000, Simple Western 1:25, Flow Cytometry 1 ug/ml, ELISA 1:100 - 1:2000, Immunohistochemistry 1:200, Immunocytochemistry/ Immunofluorescence 1:20-1:100, Immunohistochemistry-Paraffin 1:200, Flow (Intracellular) 1 ug/ml
Application Notes	In Western blot, a band is seen at ~45 kDa representing Actin. The observed molecular weight of the protein may vary from the listed predicted molecular weight due to post translational modifications, post translation cleavages, relative charges, and other experimental factors. In IHC-P and ICC/IF, cytoplasmic staining is observed. Prior to immunostaining paraffin tissues, antigen retrieval with sodium citrate buffer (pH 6.0) is recommended. In Simple Western only 10 - 15 uL of the recommended dilution is used per data point. See Simple Western Antibody Database for Simple Western validation: Tested in HeLa lysate 0.1 mg/mL, separated by Size, antibody dilution of 1:25, apparent MW was 50 kDa. Separated by Size-Wes, Sally Sue/Peggy Sue.

Images


Western Blot: Actin Antibody (mAbGEa) [NB100-74340] - Total protein from HeLa, 3T3, PC12 and Bovine normal tissue was separated on a 12% gel by SDS-PAGE, transferred to PVDF membrane and blocked in 5% non-fat milk in TBST. The membrane was probed with 1.0 ug/mL anti-Actin in 5% blocking buffer and detected with an anti-mouse IgM secondary antibody using chemiluminescence.




Immunocytochemistry/Immunofluorescence: Actin Antibody (mAbGEa) [NB100-74340] - NIH-3T3 cells were fixed for 10 minutes using 10% formalin and then permeabilized for 5 minutes using 1X PBS + 0.05% Triton X-100. The cells were incubated with anti-Actin (mAbGEa) at 5 ug/mL overnight at 4C and detected with an anti-Mouse IgM DyLight 488 (Green) at a 1:500 dilution. Nuclei were counterstained with DAPI (Blue). Cells were imaged using a 40X objective.




Simple Western: Actin Antibody (mAbGEa) [NB100-74340] - Image shows a specific band for Actin in 0.1 mg/mL of HeLa lysate. This experiment was performed under reducing conditions using the 12-230 kDa separation system.




Immunohistochemistry-Paraffin: Actin Antibody (mAbGEa) [NB100-74340] - Both normal and cancer biopsies of deparaffinized Human tonsil tissues.




Flow Cytometry: Actin Antibody (mAbGEa) [NB100-74340] - Analysis of HeLa cells using mouse Actin antibody (Orange) and Isotype control Antibody (Blue).




Western Blot: Actin Antibody (mAbGEa) [NB100-74340] - Increased relative abundance of Park7 in Lp(a) mice. Western blots were used to validate the comparative proteomic analysis between the Lp(a) and wildtype mice for the Park7 protein. Pooled liver protein extracts (n = 4 livers per pool) were separated by SDS PAGE in multiple replicates (n = 7) and transferred onto nitrocellulose membrane. Membranes were probed with an anti-Park7 antibody using an anti-actin antibody as a loading control. Liver protein extracts were the same as those used for Figure 4(a) as well as fresh liver protein extracts from new mice of the same age, sex, and genotype. Image collected and cropped by CiteAb from the following publication (null), licensed under a CC-BY license.




Immunocytochemistry/Immunofluorescence: Actin Antibody (mAbGEa) [NB100-74340] - HeLa cells were fixed for 10 minutes using 10% formalin and then permeabilized for 5 minutes using 1X PBS + 0.05% Triton-X100. The cells were incubated with anti-Actin Antibody (mAbGEa) at 2 ug/ml overnight at 4C and detected with an anti-mouse Dylight 488 (Green) at a 1:500 dilution. Actin was detected with Phalloidin 568 (Red) at a 1:200 dilution. Nuclei were counterstained with DAPI (Blue). Cells were imaged using a 40X objective.




Flow (Intracellular): Actin Antibody (mAbGEa) [NB100-74340] - An intracellular stain was performed on A549 cells with NB100-74340 (blue) and a matched isotype control (orange). Cells were fixed with 4% PFA and then permeabilized with 0.1% saponin. Cells were incubated in an antibody dilution of 1 ug/mL for 30 minutes at room temperature, followed by mouse IgM Alexa Fluor 488-conjugated secondary antibody.




Western Blot: Actin Antibody (mAbGEa) [NB100-74340] - Analysis of Actin expression in 2) HeLa, 3) NTERA-2, 4) A431, 5) HepG2, 6) MCF7, 7) NIH-3T3, 8) PC-12 and 9) COS-7 whole cell lysates.




Western Blot: Actin Antibody (mAbGEa) [NB100-74340] - Analysis of Jurkat and CHO cell lysates using actin antibody [NB100-74340] at 1:100.



Immunocytochemistry/Immunofluorescence: Actin Antibody (mAbGEa) [NB100-74340] - Actin was detected in NIH-3T3 cells fixed with methanol using mouse anti-mouse beta-Actin monoclonal antibody (NB100-74340) at 1:1800. Cells were stained using Northern Lights 557 conjugated anti-mouse secondary antibody (NL007) and counterstained with DAPI.




Immunocytochemistry/Immunofluorescence: Actin Antibody (mAbGEa) [NB100-74340] - HeLa cells were fixed for 10 minutes using 10% formalin and then permeabilized for 5 minutes using 1X PBS + 0.05% Triton X-100. The cells were incubated with anti-Actin (mAbGEa) at 2 ug/mL overnight at 4C and detected with an anti-mouse IgM DyLight 488 (Green) at 1:500. Nuclei were counterstained with DAPI (Blue). Cells were imaged using a 40X objective.




NOVUS
BIOLOGICALS
a biotechne brand


Immunohistochemistry-Paraffin: Actin Antibody (mAbGEa) [NB100-74340] - Analysis of Actin on human breast cancer tissue using DAB with hematoxylin counterstain.




Immunohistochemistry-Paraffin: Actin Antibody (mAbGEa) [NB100-74340] - Both normal and cancer biopsies of deparaffinized Human colon carcinoma tissues.




Immunohistochemistry-Paraffin: Actin Antibody (mAbGEa) [NB100-74340] - Both normal and cancer biopsies of deparaffinized Human skeletal muscle tissues.




Western Blot: Actin Antibody (mAbGEa) [NB100-74340] - Lp(a) upregulates GPx1 & Prdx6 expression in human HepG2 cells. HepG2 cells were treated with 5 $\mu\text{g}/\text{mL}$ of Lp(a) or LDL for 6 hours at 37°C. Western blots of cell lysates were performed with an anti-Gpx1 antibody (a), an anti-Prdx6 antibody (b), an anti-Sod1 antibody (c), & an anti-Park7 antibody (d) using an anti-actin antibody as a loading control. Representative blots are shown. Protein levels were normalized against actin & expressed relative to that of untreated cells. Results are expressed as mean \pm SEM for pooled triplicate incubations run in quadruplicate. \square $P < 0.05$, relative to untreated HepG2 cells. Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/30596094>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.




Western Blot: Actin Antibody (mAbGEa) [NB100-74340] - DenA deneddylase activity. (A) Recombinant human DEN1 & fungal DenA deneddylate a human CUL1-Nedd8 substrate in vitro. SDS-PAGE & subsequent western analysis show cleavage of the substrate (\square 60 kDa) producing the C-terminal CUL1 (\square 50 kDa) as outlined in experimental procedures. (B) Deneddylation test in a heterologous yeast system. *A. nidulans* DenA can remove Rub1 from CulD in heterologous expression experiments in *S. cerevisiae*. DenA expressed as native protein or C-terminally fused w/ a V5/His6 epitope tag. Both variants driven by the inducible GAL1 promoter. CulD, N-terminal-fused w/ the LexA activation domain, expressed under control of the constitutive ADH promoter. *A. nidulans* proteins expressed in *S. cerevisiae* wild type & Δ csn5 background. Western analysis w/ antibodies against Rub1 (α -Rub1), the LexA epitope (α -LexA) & the V5 epitope (α -V5) performed. Detection w/ α -Rub1 generated two additional signals upon culD expression, representing LexA-CulD & a second CulD pool where LexA unspecifically cleaved off. Both signals disappeared upon co-expression of DenA indicating deneddylation activity (red arrows). The slower migrating signal of α -LexA western experiments corresponded to Rub1 modified LexA-CulD. This signal absent when DenA co-expressed. Detection of the V5 tag applied to monitor DenA expression. The neddylated yeast cullin migrating at around 100 kDa not affected by DenA activity. (C) Deneddylation of fungal CulA by CSN & DenA. Whole cell lysates of *A. nidulans* wild type, Δ csnE & Δ denA probed w/ α -CulA. The ratio of neddylated CulA (CulA*N8; \square 106 kDa*) to non-neddylated CulA (\square 96 kDa**) calculated from three independent experiments. Membranes reprobed w/ α -Actin (α -ACT) for normalization. Image collected & cropped by CiteAb from the following publication (<https://dx.plos.org/10.1371/journal.pgen.1003275>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.




Western Blot: Actin Antibody (mAbGEa) [NB100-74340] - Fibrillar adhesions & FN fibrils are dynamic structures governed by Rab21 & PPFIA1. (a) Living siCTL & siRAB21 ECs incubated w/ IST9 mAb for 30 min, fixed, acid washed & stained. RAB21 silencing impairs ED-A FN endocytosis as revealed by decrease of IST9 punctae co-localizing w/ endosome marker EEA1. Right panels are magnifications of boxed areas in left panels. Data are mean value \pm s.e.m., n=70 cells per condition pooled from two independent experiments. ***P<0.001. (b) Left panel, WB analysis of PPFIA1, RAB21 & actin on total lysates of siCTL, siPPFIA1 & siPPFIA1+siRAB21 ECs. Right panel, WB analysis of insoluble matrix fraction of ECs extracted w/ DOC buffer. PPFIA1 silencing dramatically reduces amount of DOC-insoluble fraction of endogenous ED-A FN. Of note, simultaneous silencing of Rab21 GTPase (siPPFIA1+siRAB21), which drives integrin endocytosis, rescues defective incorporation of endogenous ED-A FN in DOC-insoluble fraction of siPPFIA1 ECs. (c) Confocal microscopy analysis of patterning of endogenous cellular ED-A FN (green) in fixed confluent ECs. Before fixation, living ECs incubated for 20 min w/ exogenous SNAKA51 (red). ED-A FN polymerizes into a fibrillar network in siCTL, but not in siPPFIA1 ECs. Simultaneous silencing of Rab21 GTPase (siPPFIA1+siRAB21) fully restores ED-A FN polymerization in siPPFIA1 ECs. SNAKA51+ active α 5 β 1 integrin localizes in fibrillar adhesion in siCTL, but not in siPPFIA1 ECs. Notably, simultaneous Rab21 (siPPFIA1+siRAB21) silencing promotes localization of SNAKA51 in fibrillar adhesion of siPPFIA1 ECs. Data are mean \pm s.e.m., n=20 cells per condition pooled from two independent experiments. Scale bar, 20 μ m (a,c, right), 50 μ m (c, left). ***P<0.001; Student's t-test. Image collected & cropped by CiteAb from following publication (<https://www.nature.com/articles/ncomms13546>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.




Western Blot: Actin Antibody (mAbGEa) [NB100-74340] - Effect of the four most common herbal formulas & single herbs on the phosphorylation of myosin light chain (MLC) protein. Briefly, A10 cells were treated with herbal formulas (A) or single herbs (B). Y27632 (Y10; 10 μ M) & calyculin A (A50; 50 μ g/ml) were used as negative & positive controls. Western blot analysis & staining with anti-phospho-MLC, anti-total-MLC, & anti-beta actin antibodies was then performed. Phospho-MLC, total-MLC, & beta actin were all obtained with their appropriate protein size bands. The relative Phospho-MLC intensity (%) was expressed as [(Phospho-MLC/total-MLC)drug treated/ (Phospho-MLC/total-MLC)cell only x 100%]. The Mean \pm SEM values for at least three independent experiments along with the representative western blot were performed. Image collected & cropped by CiteAb from the following publication (<https://dx.plos.org/10.1371/journal.pone.0145109>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.




Western Blot: Actin Antibody (mAbGEa) [NB100-74340] - CSN targets DEN1/DenA for degradation in fungal & human cells. (A) Quantitative analysis of repeated western blot experiments displayed the differences in DenA abundance in fungal wild type & Δ csnG cells. DenA levels of the different asexual developmental time points are shown relative to the vegetative (veg.) control for each strain. Anti-Actin was applied as loading control (statistics: 2-way ANOVA; $n=3$; * $p<0.05$, ** $p<0.01$). (B) Xpress-DEN1 was overexpressed in siGFP & siCSN1 human cells & steady state Xpress-DEN1 levels were estimated by western analysis with the α -Xpress antibody. Xpress-CSN1 was overexpressed in siCSN1 cells & DEN1 & CSN8 were probed with appropriate antibodies. (C) Xpress-DEN1 was overexpressed in siGFP human cells & the proteasome inhibitor MG132 was added 6 h before cell lysis at a final concentration of 10 μ M. Cyclohexamide (CHX) was added in a final concentration of 10 μ g/ml (D) Xpress-DEN1 was co-expressed in HeLa cells together with Xpress-CSN1wt, Xpress-CSN1(1–221) or Xpress-CSN1(222–527) in the absence or in the presence of MG132 (right panel), which was added 6 h before cell lysis. Cells were lysed 24 h after co-transfection & lysates were analyzed by western blot using the α -DEN1 antibody (0=only Xpress-DEN1). Image collected & cropped by CiteAb from the following publication (<https://dx.plos.org/10.1371/journal.pgen.1003275>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.




Western Blot: Actin Antibody (mAbGEa) [NB100-74340] - Analysis of Ia production by engineered *S. cerevisiae*. (A) Analysis of the synthesis of Ia in *S. cerevisiae* strains, producing actin-R177K, E270D & E270Q. Yeast strains, producing actin-R177K, E270D or E270Q & transformed with the Ia-containing plasmid (Ia) or the vector alone (Vector), were cultivated in SGal for 20 h at 30°C. Cells were broken by glass beads treatment & analyzed by ³²P-ADP-ribosylation in the presence of additionally added purified wild type yeast actin (1 μ g). Labeled bands represent modified yeast actin & confirm intracellular production of functionally active Ia by the *S. cerevisiae* strains. (B) Production of Ia by the wild type *S. cerevisiae* strain. Wild type yeast strains harboring the Ia-containing plasmid (Ia) or the control vector (vector) were cultivated in glucose-containing liquid medium until OD₅₉₅ = 0.5. Afterwards, glucose was replaced by galactose & cultivation continued for 9 h at 30°C. Cells were lysed & the resulting extract preparations were ADP-ribosylated in the presence of Ia (+ Ia), TccC3 toxin of *P. luminescens* [42] (+ TccC3), purified muscle actin (+ α -actin) or tested in Western blotting with the anti-actin serum to show equal actin concentrations in the samples. (C, D) Mass spectrometry of actin variants. MALDI-TOF MS of wild type (C) & actin-R177K (D) protein variants isolated from *S. cerevisiae*. Spectra demonstrate disappearance of R177- & appearance of K177-containing peptide in mass analysis (substituted amino acid residue within identified peptides is shown in red). Image collected & cropped by CiteAb from the following publication (<https://dx.plos.org/10.1371/journal.pone.0145708>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Western Blot: Actin Antibody (mAbGEa) [NB100-74340] - $\alpha 5\beta 1$ regulates ED-A FN secretion & polymerization. (a) Western blot analysis of lysates of ECs control (siCTL) & $\alpha 5$ integrin subunit (siITGA5) silenced ECs. Cells were lysed 24 hours after the second siRNA oligofection & proteins were separated by SDS-PAGE & probed for $\alpha 5$ integrin subunit or actin (for control purposes). (b) Confocal microscopy analysis of IST9 mAb-labelled endogenous ED-A FN (green) in confluent ECs. ED-A FN polymerizes into a fibrillar network in siCTL, but not in siITGA5 ECs in which it accumulates in the TGN46+ (red) TGN cisternae. The relative amount of fibrillar ED-A FN area was calculated in siCTL & siITGA5 ECs. Data are mean \pm s.e.m., n=20 cells per condition pooled from two independent experiments. ***P<0.001; Student's t-test. (c) Western blot analysis of soluble ED-A FN released by confluent ECs seeded on Transwell inserts. An equal percentage of apical & basolateral volumes of medium were collected after 72 h of culture, from different wells of siCTL or siITGA5 ECs. Equal amounts of exogenous rabbit IgG were added to samples (spike normalization) for loading control purposes. Quantification of the ratio between apical or basolateral amount of ED-A FN released by siCTL over siITGA5 ECs. $\alpha 5$ integrin subunit silencing impairs basolateral, but not apical ED-A FN secretion. Data are mean \pm s.e.m., n=8 wells per condition pooled from four independent experiments. **P<0.01; Student's t-test. Scale bar, 50 μ m (b). Image collected & cropped by CiteAb from the following publication (<https://www.nature.com/articles/ncomms13546>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Western Blot: Actin Antibody (mAbGEa) [NB100-74340] - Effect of the four most common herbal formulas & single herbs on the phosphorylation of myosin light chain (MLC) protein. Briefly, A10 cells were treated with herbal formulas (A) or single herbs (B). Y27632 (Y10; 10 μ M) & calyculin A (A50; 50 μ g/ml) were used as negative & positive controls. Western blot analysis & staining with anti-phospho-MLC, anti-total-MLC, & anti-beta actin antibodies was then performed. Phospho-MLC, total-MLC, & beta actin were all obtained with their appropriate protein size bands. The relative Phospho-MLC intensity (%) was expressed as [(Phospho-MLC/total-MLC)drug treated/ (Phospho-MLC/total-MLC)cell only x 100%]. The Mean \pm SEM values for at least three independent experiments along with the representative western blot were performed. Image collected & cropped by CiteAb from the following publication (<https://dx.plos.org/10.1371/journal.pone.0145109>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Publications

Hamann A, Osiewacz HD, Teichert I. *Sordaria macrospora* Sterile Mutant pro34 Is Impaired in Respiratory Complex I Assembly *Journal of Fungi* 2022-09-27 [PMID: 36294581] (Western Blot, Plant)

Parra Bravo C, Krukowski K, Barker S et Al. Anti-acetylated-tau immunotherapy is neuroprotective in tauopathy and brain injury *Mol Neurodegener* 2024-06-24 [PMID: 38915105]

Wang C, Zhang Y, Zhou B et al. Modifications of Two ESCRT-I Subunits with Distinct Ubiquitin Chains Regulate Plant Immunity *bioRxiv* 2023-08-16 (WB, Plant)

Details:
Arabidopsis thaliana

Biehler C, Rothenberg Ke, Jette A Et Al. Pak1 and PP2A antagonize aPKC function to support cortical tension induced by the Crumbs-Yurt complex *eLife* 2021-07-02 [PMID: 34212861] (WB)

Biehler C, Rothenberg K, Jetté A, et al. Functional plasticity of polarity proteins controls epithelial tissue architecture *bioRxiv* 2021-01-06 (WB, Drosophila)

Frey S, Lahmann Y et al. Deletion of Smgpi1 encoding a GPI-anchored protein suppresses sterility of the STRIPAK mutant delta Smmob3 in the filamentous ascomycete *Sordaria macrospora*. *Mol Microbiol* 2015-01-08 [PMID: 25989468] (WB, Fungus)

Details:
Sordaria macrospora

Belyy A, Tabakova I et al. Roles of Asp179 and Glu270 in ADP-Ribosylation of Actin by *Clostridium perfringens* Iota Toxin. *PLoS One* 2015-12-30 [PMID: 26713879] (WB, Yeast)

Details:
Saccharomyces cerevisiae (Yeast)

Rodger E J, Porteous C M et al. Proteomic Analysis of Liver from Human Lipoprotein(a) Transgenic Mice Shows an Oxidative Stress and Lipid Export Response. *Biomed Res Int* 2019-01-01 [PMID: 30596094] (WB, Mouse, Human)

Redhu AK, Bhat JP Mitochondrial glucose 6-phosphate dehydrogenase and 6-phosphogluconate dehydrogenase abrogate p53 induced apoptosis in a yeast model: Possible implications for apoptosis resistance in cancer cells *Biochim Biophys Acta Gen Subj* 2019-12-17 [PMID: 31862471] (WB, Yeast)

Redhu AK, Bhat JP Novel Proteomic changes in Yeast Mitochondria provide insights into mitochondrial functioning upon over-expression of human p53 *bioRxiv* 2019-08-24 (WB, Yeast)

Taylor SM, Giuffre E, Moseley P, Hitchcock PF The MicroRNA, miR-18a, Regulates NeuroD and Photoreceptor Differentiation in the Retina of Zebrafish *Dev Neurobiol* 2019-02-01 [PMID: 30615274] (WB, Zebrafish)

Audesse, AJ;Dhakal, S;Hassell, LA;Gardell, Z;Nemtsova, Y;Webb, AE; FOXO3 directly regulates an autophagy network to functionally regulate proteostasis in adult neural stem cells *PLoS Genet.* 2019-04-01 [PMID: 30973875] (WB, Mouse)

More publications at <http://www.novusbio.com/NB100-74340>

Procedures

Western Blot Protocol Specific for Actin Antibody (mAbGEa) [NB100-74340]

Western Blot Protocol

1. Perform SDS-PAGE on samples to be analyzed, loading 40 ug of total protein per lane.
 2. Transfer proteins to membrane according to the instructions provided by the manufacturer of the membrane and transfer apparatus.
 3. Stain according to standard Ponceau S procedure (or similar product) to assess transfer success, and mark molecular weight standards where appropriate.
 4. Rinse the blot.
 5. Block the membrane using standard blocking buffer for at least 1 hour.
 6. Wash the membrane in wash buffer three times for 10 minutes each.
 7. Dilute primary antibody in blocking buffer and incubate 1 hour at room temperature.
 8. Wash the membrane in wash buffer three times for 10 minutes each.
 9. Apply the diluted HRP conjugated secondary antibody in blocking buffer (as per manufacturers instructions) and incubate 1 hour at room temperature.
 10. Wash the blot in wash buffer three times for 10 minutes each (this step can be repeated as required to reduce background).
 11. Apply the detection reagent of choice in accordance with the manufacturers instructions.
- Note: Tween-20 can be added to the blocking or antibody dilution buffer at a final concentration of 0.05-0.2%.

*The above information is only intended as a guide. The researcher should determine what protocol best meets their needs. Please follow safe laboratory procedures.

Immunohistochemistry protocol for Actin Antibody (NB100-74340)

Immunohistochemistry-Paraffin Embedded Sections

Antigen Unmasking:

Bring slides to a boil in 10 mM sodium citrate buffer (pH 6.0) then maintain at a sub-boiling temperature for 10 minutes. Cool slides on bench-top for 30 minutes.

Staining:

1. Wash sections in deionized water three times for 5 minutes each.
2. Wash sections in wash buffer for 5 minutes.
3. Block each section with 100-400 ul blocking solution for 1 hour at room temperature.
4. Remove blocking solution and add 100-400 ul diluted primary antibody. Incubate overnight at 4 C.
5. Remove antibody solution and wash sections in wash buffer three times for 5 minutes each.
6. Add 100-400 ul biotinylated diluted secondary antibody. Incubate 30 minutes at room temperature.
7. Remove secondary antibody solution and wash sections three times with wash buffer for 5 minutes each.
8. Add 100-400 ul Streptavidin-HRP reagent to each section and incubate for 30 minutes at room temperature.
9. Wash sections three times in wash buffer for 5 minutes each.
10. Add 100-400 ul DAB substrate to each section and monitor staining closely.
11. As soon as the sections develop, immerse slides in deionized water.
12. Counterstain sections in hematoxylin.
13. Wash sections in deionized water two times for 5 minutes each.
14. Dehydrate sections.
15. Mount coverslips.

*The above information is only intended as a guide. The researcher should determine what protocol best meets their needs. Please follow safe laboratory procedures.

Immunocytochemistry/Immunofluorescence protocol for Actin Antibody (NB100-74340)

Immunocytochemistry Protocol

Culture cells to appropriate density in 35 mm culture dishes or 6-well plates.

1. Remove culture medium and add 10% formalin to the dish. Fix at room temperature for 30 minutes.
2. Remove the formalin and add ice cold methanol. Incubate for 5-10 minutes.
3. Remove methanol and add washing solution (i.e. PBS). Be sure to not let the specimen dry out. Wash three times for 10 minutes.
4. To block nonspecific antibody binding incubate in 10% normal goat serum from 1 hour to overnight at room temperature.
5. Add primary antibody at appropriate dilution and incubate at room temperature from 2 hours to overnight at room temperature.
6. Remove primary antibody and replace with washing solution. Wash three times for 10 minutes.
7. Add secondary antibody at appropriate dilution. Incubate for 1 hour at room temperature.
8. Remove antibody and replace with wash solution, then wash for 10 minutes. Add Hoechst 33258 to wash solution at 1:25,000 and incubate for 10 minutes. Wash a third time for 10 minutes.
9. Cells can be viewed directly after washing. The plates can also be stored in PBS containing Azide covered in Parafilm (TM). Cells can also be cover-slipped using Fluoromount, with appropriate sealing.

*The above information is only intended as a guide. The researcher should determine what protocol best meets their needs. Please follow safe laboratory procedures.



Flow (Intracellular) Protocol for Actin Antibody (NB100-74340)

Protocol for Flow Cytometry Intracellular Staining

Sample Preparation.

1. Grow cells to 60-85% confluency. Flow cytometry requires between 2×10^5 and 1×10^6 cells for optimal performance.
2. If cells are adherent, harvest gently by washing once with staining buffer and then scraping. Avoid using trypsin as this can disrupt certain epitopes of interest. If enzymatic harvest is required, use Accutase, Collagenase, or TrypLE Express for a less damaging option.
3. Reserve 100 μ L for counting, then transfer cell volume into a 50 mL conical tube and centrifuge for 8 minutes at 400 RCF.
 - a. Count cells using a hemocytometer and a 1:1 trypan blue exclusion stain to determine cell viability before starting the flow protocol. If cells appear blue, do not proceed.
4. Re-suspend cells to a concentration of 1×10^6 cells/mL in staining buffer (NBP2-26247).
5. Aliquot out 100 μ L samples in accordance with your experimental samples.

Tip: When cell surface and intracellular staining are required in the same sample, it is advisable that the cell surface staining be performed first since the fixation and permeabilization steps might reduce the availability of surface antigens.

Intracellular Staining.

Tip: When performing intracellular staining, it is important to use appropriate fixation and permeabilization reagents based upon the target and its subcellular location. Generally, our Intracellular Flow Assay Kit (NBP2-29450) is a good place to start as it contains an optimized combination of reagents for intracellular staining as well as an inhibitor of intracellular protein transport (necessary if staining secreted proteins). Certain targets may require more gentle or transient permeabilization protocols such as the commonly employed methanol or saponin-based methods.

Protocol for Cytoplasmic Targets:

1. Fix the cells by adding 100 μ L fixation solution (such as 4% PFA) to each sample for 10-15 minutes.
2. Permeabilize cells by adding 100 μ L of a permeabilization buffer to every 1×10^6 cells present in the sample. Mix well and incubate at room temperature for 15 minutes.
 - a. For cytoplasmic targets, use a gentle permeabilization solution such as 1X PBS + 0.5% Saponin or 1X PBS + 0.5% Tween-20.
 - b. To maintain the permeabilized state throughout your experiment, use staining buffer + 0.1% of the permeabilization reagent (i.e. 0.1% Tween-20 or 0.1% Saponin).
3. Following the 15 minute incubation, add 2 mL of the staining buffer + 0.1% permeabilizer to each sample.
4. Centrifuge for 1 minute at 400 RCF.
5. Discard supernatant and re-suspend in 100 μ L of staining buffer + 0.1% permeabilizer.
6. Add appropriate amount of each antibody (eg. 1 test or 1 μ g per sample, as experimentally determined).
7. Mix well and incubate at room temperature for 30 minutes- 1 hour. Gently mix samples every 10-15 minutes.
8. Following the primary/conjugate incubation, add 1-2 mL/sample of staining buffer + 0.1% permeabilizer and centrifuge for 1 minute at 400 RCF.
9. Wash twice by re-suspending cells in staining buffer (2 mL for tubes or 200 μ L for wells) and centrifuging at 400 RCF for 5 minutes. Discard supernatant.
10. Add appropriate amount of secondary antibody (as experimentally determined) to each sample.
11. Incubate at room temperature in dark for 20 minutes.
12. Add 1-2 mL of staining buffer and centrifuge at 400 RCF for 1 minute and discard supernatant.
13. Wash twice by re-suspending cells in staining buffer (2 mL for tubes or 200 μ L for wells) and centrifuging at 400 RCF for 5 minutes. Discard supernatant.
14. Resuspend in an appropriate volume of staining buffer (usually 500 μ L per sample) and proceed with analysis on your flow cytometer.



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NB100-74340

NBP1-42569	HepG2 Whole Cell Lysate
NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB7539	Goat anti-Mouse IgG (H+L) Secondary Antibody [HRP]
NBP1-96975-0.5mg	Mouse IgM Kappa Light Chain Isotype Control (MMK)

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-74340

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

