

Product Datasheet

Presenilin-1 Antibody (APS 18) NB100-74510

Unit Size: 200ug

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Reviews: 1 Publications: 5

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB100-74510

Updated 9/9/2025 v.20.1

**Earn rewards for product
reviews and publications.**

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB100-74510



NB100-74510

Presenilin-1 Antibody (APS 18)

Product Information	
Unit Size	200ug
Concentration	1 mg/ml
Storage	Store at -20C. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	APS 18
Preservative	0.05% Sodium Azide
Isotype	IgG1
Purity	Protein G purified
Buffer	PBS with 1 mg/ml BSA


Product Description	
Description	Novus Biologicals Mouse Presenilin-1 Antibody (APS 18) (NB100-74510) is a monoclonal antibody validated for use in IHC, WB, ELISA, ICC/IF and IP. Anti-Presenilin-1 Antibody: Cited in 5 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Mouse
Gene ID	5663
Gene Symbol	PSEN1
Species	Human, Mouse, Rat, Primate
Reactivity Notes	Please note that this antibody is reactive to Mouse and derived from the same host, Mouse. Additional Mouse on Mouse blocking steps may be required for IHC and ICC experiments. Please contact Technical Support for more information.
Specificity/Sensitivity	Presenilin 1 (APS 18)
Immunogen	Synthetic peptide corresponding to residues C S(313) K Y N A E S T E R E S Q D T V A E N D D G(334) of human PS1.

Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, ELISA, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunoprecipitation
Recommended Dilutions	Western Blot 1:50 - 1:250, ELISA 1:1000, Immunohistochemistry 1:20 - 1:200, Immunocytochemistry/ Immunofluorescence 1:50 - 1:200, Immunoprecipitation 1:10 - 1:500, Immunohistochemistry-Paraffin 1:20 - 1:200
Application Notes	IP usage was reported in scientific literatures (PMID: 9852041 and 10092585).




Images


Western Blot: Presenilin-1 Antibody (APS 18) [NB100-74510] - Analysis of THP-1 cell lysate.



Immunocytochemistry/Immunofluorescence: Presenilin-1 Antibody (APS 18) [NB100-74510] - Analysis of Presenilin 1 using Presenilin 1 Monoclonal antibody (APS 18) shows staining in A2058 melanoma cells. Presenilin 1 staining (green), F-Actin staining with Phalloidin (red) and nuclei with DAPI (blue) is shown. Cells were grown on chamber slides and fixed with formaldehyde prior to staining. Cells were probed without (control) or with or an antibody recognizing Presenilin 1 at a dilution of 1:20-1:100 over night at 4C, washed with PBS and incubated with a DyLight-488 conjugated.




Immunohistochemistry-Paraffin: Presenilin-1 Antibody (APS 18) [NB100-74510] - Normal biopsies of deparaffinized Human liver tissue.




Immunocytochemistry/Immunofluorescence: Presenilin-1 Antibody (APS 18) [NB100-74510] - Staining of PS1 in mouse fibroblasts.


Fig. 1




Immunocytochemistry/Immunofluorescence: Presenilin-1 Antibody (APS 18) [NB100-74510] - Analysis of Presenilin 1 using Presenilin 1 Monoclonal antibody (APS 18) shows staining in MCF-7 cells. Presenilin 1 staining (green), F-Actin staining with Phalloidin (red) and nuclei with DAPI (blue) is shown. Cells were grown on chamber slides and fixed with formaldehyde prior to staining. Cells were probed without (control) or with or an antibody recognizing Presenilin 1 at a dilution of 1:20-1:100 over night at 4C, washed with PBS and incubated with a DyLight-488 conjugated.



Immunocytochemistry/Immunofluorescence: Presenilin-1 Antibody (APS 18) [NB100-74510] - Analysis of Presenilin 1 using Presenilin 1 Monoclonal antibody (APS 18) shows staining in HeLa cells. Presenilin 1 staining (green), F-Actin staining with Phalloidin (red) and nuclei with DAPI (blue) is shown. Cells were grown on chamber slides and fixed with formaldehyde prior to staining. Cells were probed without (control) or with or an antibody recognizing Presenilin 1 at a dilution of 1:20-1:100 over night at 4C, washed with PBS and incubated with a DyLight-488 conjugated.



Immunohistochemistry-Paraffin: Presenilin-1 Antibody (APS 18) [NB100-74510] - Normal biopsies of deparaffinized Human tonsil tissue.



Publications

Zhang H, Jiang R, Zhou J et al. CTL Attenuation Regulated by PS1 in Cancer-Associated Fibroblast Front Immunol 2020-06-10 [PMID: 32587587] (IF/IHC, Mouse)

McCombs JE, Gibson EA, Palmer AE. Using a genetically targeted sensor to investigate the role of presenilin-1 in ER Ca²⁺ levels and dynamics. Mol Biosyst. 2010-04-08 [PMID: 20379593] (WB, Mouse)

Stahl B et al. Direct interaction of Alzheimer's disease-related presenilin 1 with armadillo protein p0071. J Biol Chem274(14):9141-8. 1999-04-02 [PMID: 10092585] (IP)

Tesco G et al. Abrogation of the presenilin 1/beta-catenin interaction and preservation of the heterodimeric presenilin 1 complex following caspase activation. J Biol Chem273(51):33909-14. 1998-12-18 [PMID: 9852041] (IP)

Diehlmann A et al. Analysis of presenilin 1 and presenilin 2 expression and processing by newly developed monoclonal antibodies. J Neurosci Res 56(4):405-19. 1999-05-15 [PMID: 10340748] (ELISA, WB, ICC/IF)



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NB100-74510

NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB7539	Goat anti-Mouse IgG (H+L) Secondary Antibody [HRP]
NBP1-97005-0.5mg	Mouse IgG1 Isotype Control (MG1)

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-74510

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

