

Product Datasheet

NCF1 Antibody NB100-790

Unit Size: 0.1 mg

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 5

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB100-790

Updated 9/9/2025 v.20.1

**Earn rewards for product
reviews and publications.**

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB100-790



NB100-790

NCF1 Antibody


| Product Information | |
|---------------------|--|
| Unit Size | 0.1 mg |
| Concentration | 0.5 mg/ml |
| Storage | Store at -20C. Avoid freeze-thaw cycles. |
| Clonality | Polyclonal |
| Preservative | 0.02% Sodium Azide |
| Isotype | IgG |
| Purity | Immunogen affinity purified |
| Buffer | Tris saline (20 mM Tris pH 7.3, 150 mM NaCl), 0.5% BSA |

| Product Description | |
|---------------------|--|
| Description | Novus Biologicals Goat NCF1 Antibody (NB100-790) is a polyclonal antibody validated for use in IHC, WB, ELISA, Flow, ICC/IF and IP. Anti-NCF1 Antibody: Cited in 5 publications. All Novus Biologicals antibodies are covered by our 100% guarantee. |
| Host | Goat |
| Gene ID | 653361 |
| Gene Symbol | NCF1 |
| Species | Human, Mouse, Porcine, Bacteria |
| Reactivity Notes | Bacteria reactivity reported in scientific literature (PMID: 28725637). |
| Immunogen | Peptide with sequence C-SESTKRKLASAV corresponding to C-Terminus according to NP_000256.3. |


| Product Application Details | |
|-----------------------------|---|
| Applications | Western Blot, Immunohistochemistry-Paraffin, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunoprecipitation, Peptide ELISA |
| Recommended Dilutions | Western Blot 0.01 - 1 ug/mL, Flow Cytometry 10 ug/mL, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunoprecipitation, Immunohistochemistry-Paraffin 5 ug/mL, Peptide ELISA Detection limit 1:20000 |
| Application Notes | WB: Approx 48 kDa band observed in lysates of cell lines Daudi, U251 and U937 and in Mouse Thymus, and approx, 50 kDa in Pig Spleen lysates (calculated MW of 44.7 kDa according to Human NP_000256.4, 46.0 kDa according to Mouse NP_001272966.1 and 45.3kDa according to Pig NP_001106691.1). Use in immunoprecipitation reported in scientific literature (PMID:28725637). |

Images


Western Blot: NCF1 Antibody [NB100-790] - Staining of U937 (A), Daudi (B), U251 (C) and negative control A431 (D) cell lysate (35 ug protein in RIPA buffer). Antibody at 0.2, 0.01, and 0.3 ug/mL respectively. Detected by chemiluminescence.




Immunohistochemistry-Paraffin: NCF1 Antibody [NB100-790] - Staining of paraffin embedded Human Colon. Antibody at 5 ug/mL. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.




Flow Cytometry: NCF1 Antibody [NB100-790] - Analysis of paraformaldehyde fixed HeLa cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10 ug/mL) followed by Alexa Fluor 488 secondary antibody (1 ug/mL). IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.



Western Blot: NCF1 Antibody [NB100-790] - Staining of Mouse Thymus (A) and Pig Spleen (B) lysate (35 ug protein in RIPA buffer). Antibody at 1 ug/mL. Primary incubation was 1 hour. Detected by chemiluminescence.



Immunofluorescence: NCF1 Antibody [NB100-790] - Analysis of paraformaldehyde fixed HeLa cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10 ug/mL) followed by Alexa Fluor 488 secondary antibody (2 ug/mL), showing cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10 ug/mL) followed by Alexa Fluor 488 secondary antibody (2 ug/mL).



Publications

Huang F, Ren X, Yuan B et al. Systemic mutation of Ncf1 ameliorates obstruction-induced renal fibrosis while macrophage-rescued NCF1 further alleviates renal fibrosis *Antioxidants & redox signaling* 2023-06-30 [PMID: 37392014]

Johnson JL, Ramadass M, Rahman F et al. The atypical small GTPase GEM/Kir is a negative regulator of the NADPH oxidase and NETs production through macroautophagy *Journal of leukocyte biology* 2021-06-04 [PMID: 34085299] (IF/IHC)

Zhou L, Tang J et al. Five Constituents in *Psoralea corylifolia* L. Attenuate Palmitic Acid-Induced Hepatocyte Injury via Inhibiting the Protein Kinase C-alpha/Nicotinamide-Adenine Dinucleotide Phosphate Oxidase Pathway. *Front Pharmacol* 2020-03-03 [PMID: 32116659] (ICC/IF, Mouse)

Roberts JoAnn S, Atanasova Kalina R, Lee Junnam et al. Opportunistic Pathogen *Porphyromonas gingivalis* Modulates Danger Signal ATP-Mediated Antibacterial NOX2 Pathways in Primary Epithelial Cells. *Frontiers in Cellular and Infection Microbiology* 2017-07-04 [PMID: 28725637] (IP, Bacteria)

Davidson-Moncada JK, Lopez-Lluch G, Segal AW, Dekker LV Involvement of protein kinase D in Fc gamma-receptor activation of the NADPH oxidase in neutrophils. *Biochem J* 2002-04-01 [PMID: 11903052]



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NB100-790

| | |
|-----------------|--|
| NBL1-13509 | NCF1 Overexpression Lysate |
| NBP2-33376H | Blue Marker Antibody (6F4-F6) [HRP] |
| HAF017 | Rabbit anti-Goat IgG Secondary Antibody [HRP (Horseradish Peroxidase)] |
| HAF109 | Donkey anti-Goat IgG Secondary Antibody [HRP (Horseradish Peroxidase)] |
| NB410-28088-1mg | Goat IgG Isotype Control |

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-790

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications



