



**NB110-40424**

Sin1/MAPKAP1 Antibody - BSA Free

**Product Information**

|                                |                                       |
|--------------------------------|---------------------------------------|
| <b>Unit Size</b>               | 100 ul                                |
| <b>Concentration</b>           | 1.0 mg/ml                             |
| <b>Storage</b>                 | Store at 4C. Do not freeze.           |
| <b>Clonality</b>               | Polyclonal                            |
| <b>Preservative</b>            | 0.09% Sodium Azide                    |
| <b>Isotype</b>                 | IgG                                   |
| <b>Purity</b>                  | Immunogen affinity purified           |
| <b>Buffer</b>                  | Tris-Citrate/Phosphate (pH 7.0 - 8.0) |
| <b>Target Molecular Weight</b> | 59 kDa                                |

**Product Description**

|                         |  |
|-------------------------|--|
| <b>Description</b>      | Novus Biologicals Rabbit Sin1/MAPKAP1 Antibody - BSA Free (NB110-40424) is a polyclonal antibody validated for use in IHC, WB and IP. Anti-Sin1/MAPKAP1 Antibody: Cited in 7 publications. All Novus Biologicals antibodies are covered by our 100% guarantee. |
| <b>Host</b>             | Rabbit   |
| <b>Gene ID</b>          | 79109  |
| <b>Gene Symbol</b>      | MAPKAP1  |
| <b>Species</b>          | Human  |
| <b>Reactivity Notes</b> | Orangutan (100%).  |
| <b>Immunogen</b>        | The immunogen recognized by this antibody maps to a region between residue 470 and the C-terminus (residue 522) of human stress-activated map kinase interacting protein 1 using the numbering given in entry NP_001006618.1 (GeneID 79109).                   |

**Product Application Details**


|                              |  |
|------------------------------|--|
| <b>Applications</b>          | Western Blot, Immunohistochemistry-Paraffin, Immunohistochemistry, Immunoprecipitation   |
| <b>Recommended Dilutions</b> | Western Blot 1:2000-1:10000, Immunohistochemistry 1:1000 - 1:5000, Immunoprecipitation 2-5 ug/mg lysate, Immunohistochemistry-Paraffin 1:1000-1:5000 |
| <b>Application Notes</b>     | Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE tissue sections.   |

## Images


Western Blot: Sin1/MAPKAP1 Antibody [NB110-40424] - Truncated Sin1/MAPKAP1 displaces endogenous Sin1/MAPKAP1 from mTORC2 in DLD1 colon cancer cells. Sin1/MAPKAP1 constructs incorporate into mTORC2 and displace endogenous Sin1/MAPKAP1. Constructs were induced for 72 hours prior to immune precipitation. mTORC2 subunits, mTOR and Rictor, only appear in myc immunoprecipitates after induction with doxycycline (Left panels); myc-deltaSin1/MAPKAP1 cannot be directly detected in precipitates due to secondary antibody cross reaction with precipitating IgG. Right panels indicate unchanging expression levels of Rictor and mTOR in immune precipitation input lysates, which is further quantified from 3 independent experiments. Image collected and cropped by CiteAb from the following publication (<https://www.oncotarget.com/fulltext/20086>), licensed under a CC-BY license.

Immunohistochemistry-Paraffin: Sin1/MAPKAP1 Antibody [NB110-40424] - Section of human prostate carcinoma. Antibody: Affinity purified rabbit anti- Sin1 used at a dilution of 1:1,000 (1ug/ml). Detection: DAB


Western Blot: Sin1/MAPKAP1 Antibody [NB110-40424] - Detection of Human Sin1 on HeLa whole cell lysate.



Truncated Sin1 displaces endogenous Sin1 from mTORC2 in DLD1 colon cancer cells. A. Schematic indicating the domain structure of Sin1 & the constructs used to displace endogenous Sin1 from mTORC2. B. Expression of myc tagged Sin1 constructs can be detected only after induction with Doxycycline (Dox). Cells were treated with 100nM of doxycycline (+) for 72 hours & expressed proteins were detected by immunoblot of whole cell lysates with anti-myc (9E10) antibodies. C. & D. Sin1 constructs incorporate into mTORC2 & displace endogenous Sin1. Constructs were induced for 72 hours prior to immune precipitation. (C) mTORC2 subunits, mTOR & Rictor, only appear in myc immunoprecipitates after induction with doxycycline (Left panels); myc-ΔSin1 cannot be directly detected in precipitates due to secondary antibody cross reaction with precipitating IgG. Right panels indicate unchanging expression levels of Rictor & mTOR in immune precipitation input lysates, which is further quantified from 3 independent experiments. E. Endogenous Sin1 & Rictor immunoprecipitates demonstrate displacement of endogenous Sin1 from mTORC2. Following induction, band shifted myc-tagged FL Sin1 can be detected in Sin1 & Rictor precipitates (Left panels). Truncated ΔSin1 can be detected in Rictor, but not Sin1, immunoprecipitates as the Sin1 antibody epitope is deleted from ΔSin1. F. Quantification of Sin1 levels detected in Rictor immunoprecipitates indicates the level of endogenous mTORC2 disruption following Sin1 construct induction (data are mean +/- S.D; n = 3). Myc-ΔSin1 displaces >80% of endogenous Sin1 while levels of myc-FL Sin1 associated with Rictor are comparable with endogenous Sin1 levels. Image collected & cropped by CiteAb from the following publication (<https://www.oncotarget.com/lookup/doi/10.18632/oncotarget.20086>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Truncated Sin1 displaces endogenous Sin1 from mTORC2 in DLD1 colon cancer cells. A. Schematic indicating the domain structure of Sin1 & the constructs used to displace endogenous Sin1 from mTORC2. B. Expression of myc tagged Sin1 constructs can be detected only after induction with Doxycycline (Dox). Cells were treated with 100nM of doxycycline (+) for 72 hours & expressed proteins were detected by immunoblot of whole cell lysates with anti-myc (9E10) antibodies. C. & D. Sin1 constructs incorporate into mTORC2 & displace endogenous Sin1. Constructs were induced for 72 hours prior to immune precipitation. (C) mTORC2 subunits, mTOR & Rictor, only appear in myc immunoprecipitates after induction with doxycycline (Left panels); myc-ΔSin1 cannot be directly detected in precipitates due to secondary antibody cross reaction with precipitating IgG. Right panels indicate unchanging expression levels of Rictor & mTOR in immune precipitation input lysates, which is further quantified from 3 independent experiments. E. Endogenous Sin1 & Rictor immunoprecipitates demonstrate displacement of endogenous Sin1 from mTORC2. Following induction, band shifted myc-tagged FL Sin1 can be detected in Sin1 & Rictor precipitates (Left panels). Truncated ΔSin1 can be detected in Rictor, but not Sin1, immunoprecipitates as the Sin1 antibody epitope is deleted from ΔSin1. F. Quantification of Sin1 levels detected in Rictor immunoprecipitates indicates the level of endogenous mTORC2 disruption following Sin1 construct induction (data are mean +/- S.D; n = 3). Myc-ΔSin1 displaces >80% of endogenous Sin1 while levels of myc-FL Sin1 associated with Rictor are comparable with endogenous Sin1 levels. Image collected & cropped by CiteAb from the following publication (<https://www.oncotarget.com/lookup/doi/10.18632/oncotarget.20086>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



## Publications

Wang Q, Zhu J et al. Tumor suppressor Pdcd4 attenuates Sin1 translation to inhibit invasion in colon carcinoma. *Oncogene* 2017-09-11 [PMID: 28692058] (WB, Human)

Huang Y, Feng G, Cai J et al. Sin1 promotes proliferation and invasion of prostate Cancer cells by modulating mTORC2-AKT and AR signaling cascades *Life Sci.* 2020-02-20 [PMID: 32088212] (IF/IHC)

Cameron AJM, Veeriah S, Marshall J JT et al. Uncoupling TORC2 from AGC kinases inhibits tumour growth *Oncotarget*. 2017-10-17 [PMID: 29156676] (WB, Human)

Das F, Ghosh-Choudhury N, Dey N et al. Unrestrained mammalian target of rapamycin complexes 1 and 2 increase expression of phosphatase and tensin homolog deleted on chromosome 10 to regulate phosphorylation of Akt kinase. *J Biol Chem* 2012-02-01 [PMID: 22184110]

Misra UK, Pizzo SV. Upregulation of mTORC2 activation by the selective agonist of EPAC, 8-CPT-2Me-cAMP, in prostate cancer cells: assembly of a multiprotein signaling complex. *J Cell Biochem* 2012-05-01 [PMID: 22173835]

Lu M, Wang J, Ives HE et al. mSin1 protein mediates SGK1 protein interaction with mTORC2 protein complex and is required for selective activation of the epithelial sodium channel. *J Biol Chem* 2011-09-01 [PMID: 21757730]

Huang J, Wu S, Wu CL et al. Signaling events downstream of mammalian target of rapamycin complex 2 are attenuated in cells and tumors deficient for the tuberous sclerosis complex tumor suppressors. *Cancer Res* 2009-08-01 [PMID: 19602587]



## Novus Biologicals USA

10730 E. Briarwood Avenue  
Centennial, CO 80112  
USA  
Phone: 303.730.1950  
Toll Free: 1.888.506.6887  
Fax: 303.730.1966  
nb-customerservice@bio-techne.com

## Bio-Techne Canada

21 Canmotor Ave  
Toronto, ON M8Z 4E6  
Canada  
Phone: 905.827.6400  
Toll Free: 855.668.8722  
Fax: 905.827.6402  
canada.inquires@bio-techne.com

## Bio-Techne Ltd

19 Barton Lane  
Abingdon Science Park  
Abingdon, OX14 3NB, United Kingdom  
Phone: (44) (0) 1235 529449  
Free Phone: 0800 37 34 15  
Fax: (44) (0) 1235 533420  
info.EMEA@bio-techne.com

## General Contact Information

[www.novusbio.com](http://www.novusbio.com)  
Technical Support: [nb-technical@bio-techne.com](mailto:nb-technical@bio-techne.com)  
Orders: [nb-customerservice@bio-techne.com](mailto:nb-customerservice@bio-techne.com)  
General: [novus@novusbio.com](mailto:novus@novusbio.com)

## Products Related to NB110-40424

|            |   |
|------------|---|
| NB800-PC1  | HeLa Whole Cell Lysate                              |
| HAF008     | Goat anti-Rabbit IgG Secondary Antibody [HRP]       |
| NB7160     | Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP] |
| NBP2-24891 | Rabbit IgG Isotype Control                          |

## Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit [www.novusbio.com/guarantee](http://www.novusbio.com/guarantee)

Earn gift cards/discounts by submitting a review: [www.novusbio.com/reviews/submit/NB110-40424](http://www.novusbio.com/reviews/submit/NB110-40424)

Earn gift cards/discounts by submitting a publication using this product:  
[www.novusbio.com/publications](http://www.novusbio.com/publications)