

Product Datasheet

Osteopontin/OPN Antibody (1B20) - BSA Free NB110-89062

Unit Size: 0.1 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Reviews: 2 Publications: 38

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB110-89062

Updated 9/9/2025 v.20.1

**Earn rewards for product
reviews and publications.**

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB110-89062



NB110-89062

Osteopontin/OPN Antibody (1B20) - BSA Free


Product Information	
Unit Size	0.1 ml
Concentration	1 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	1B20
Preservative	0.05% Sodium Azide
Isotype	IgG1
Purity	Protein G purified
Buffer	PBS

Product Description	
Description	Novus Biologicals Mouse Osteopontin/OPN Antibody (1B20) - BSA Free (NB110-89062) is a monoclonal antibody validated for use in IHC, WB, ICC/IF, Simple Western and IP. Anti-Osteopontin/OPN Antibody: Cited in 34 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Mouse
Gene ID	6696
Gene Symbol	SPP1
Species	Human, Mouse, Rat, Rabbit
Reactivity Notes	Human, Rat and Rabbit. Rabbit reactivity reported in scientific literature (PMID:33001866). Use in Mouse reported in scientific literature (PMID:33580126).
Immunogen	Synthetic peptide corresponding to the C-terminus of human Osteopontin [Swiss-Prot# P10451].


Product Application Details	
Applications	Western Blot, Simple Western, Immunohistochemistry-Paraffin, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Frozen, Immunoprecipitation
Recommended Dilutions	Western Blot 1:1000, Simple Western 1:20, Immunohistochemistry 1:100, Immunocytochemistry/ Immunofluorescence 1:50, Immunoprecipitation 1:10-1:500, Immunohistochemistry-Paraffin 1:100, Immunohistochemistry-Frozen

Images

Western Blot: Osteopontin/OPN Antibody (1B20) - BSA Free [NB110-89062] - Analysis of Osteopontin expression in U2OS whole cell lysate.




Immunocytochemistry/Immunofluorescence: Osteopontin/OPN Antibody (1B20) - BSA Free [NB110-89062] - Osteopontin antibody was tested at 1:50 in U2OS cells with FITC (green). Nuclei and actin were counterstained with Dapi (blue) and Phalloidin (red).




Immunohistochemistry: Osteopontin/OPN Antibody (1B20) - BSA Free [NB110-89062] - Immunohistological results at 12 weeks for (A) Group I, (B) Group II (C) Group III. The pictures are arranged by staining technique (columns) and by the investigated treatment (rows). Areas that stained positive for osteopontin are indicated by red arrowheads. Image collected and cropped by CiteAb from the following publication (<https://peerj.com/articles/3513>) licensed under a CC-BY license.




Immunohistochemistry: Osteopontin/OPN Antibody (1B20) - BSA Free [NB110-89062] - Osteogenic differentiation in 3D. Immunostaining for osteopontin (OPN) and nuclear DAPI in 3D cultures after 5 weeks. Scale bar = 50 um. Image collected and cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/34948393/>) licensed under a CC-BY license.




Immunohistochemistry: Osteopontin/OPN Antibody (1B20) - BSA Free [NB110-89062] - Analysis of Osteopontin on human lung adenocarcinoma.




Immunohistochemistry: Osteopontin/OPN Antibody (1B20) - BSA Free [NB110-89062] - Osteopontin in (A) Group I, (B) Group II (C) Group III at 6 weeks. The pictures are arranged by staining technique (columns) and the investigated treatment (rows). Areas that stained positive for osteocalcin and osteopontin are indicated by red arrowheads. NB, new bone; OS, osteoid; FM, fatty marrow. Image collected and cropped by CiteAb from the following publication (<https://peerj.com/articles/3513>) licensed under a CC-BY license.




Simple Western: Osteopontin/OPN Antibody (1B20) - BSA Free [NB110-89062] - Simple Western lane view shows a specific band for Osteopontin in 0.5 mg/ml of Human Breast Cancer lysate. This experiment was performed under reducing conditions using the 12-230 kDa separation system. * Non-specific interaction with the 230 kDa Simple Western standard may be seen with this antibody




Immunohistochemistry: Osteopontin/OPN Antibody (1B20) - BSA Free [NB110-89062] - Photomicrographs of immunostaining for osteocalcin & osteopontin in (A) Group I, (B) Group II (C) Group III at 6 weeks. The pictures are arranged by staining technique (columns) & the investigated treatment (rows). Areas that stained positive for osteocalcin & osteopontin are indicated by red arrowheads. NB, new bone; OS, osteoid; FM, fatty marrow. Image collected & cropped by CiteAb from the following publication (<https://peerj.com/articles/3513>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.




Immunohistochemistry: Osteopontin/OPN Antibody (1B20) - BSA Free [NB110-89062] - Photomicrographs of immunostaining for osteocalcin & osteopontin in (A) Group I, (B) Group II (C) Group III at 6 weeks. The pictures are arranged by staining technique (columns) & the investigated treatment (rows). Areas that stained positive for osteocalcin & osteopontin are indicated by red arrowheads. NB, new bone; OS, osteoid; FM, fatty marrow. Image collected & cropped by CiteAb from the following publication (<https://peerj.com/articles/3513>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Western Blot: Osteopontin/OPN Antibody (1B20) - BSA Free [NB110-89062] - Photomicrographs of immunostaining for osteocalcin & osteopontin in (A) Group I, (B) Group II (C) Group III at 6 weeks. The pictures are arranged by staining technique (columns) & the investigated treatment (rows). Areas that stained positive for osteocalcin & osteopontin are indicated by red arrowheads. NB, new bone; OS, osteoid; FM, fatty marrow. Image collected & cropped by CiteAb from the following publication (<https://peerj.com/articles/3513>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Osteopontin/OPN (1B20) was detected in immersion fixed HepG2 human hepatocellular carcinoma cell line using Rat anti-Osteopontin/OPN (1B20) Protein G Purified Monoclonal Antibody conjugated to Alexa Fluor® 488 (Catalog # NB110-89062AF488) (green) at 10 μg/mL overnight at 4C. Cells were counterstained with DAPI (dark blue). Cells were imaged using a 100X objective and digitally deconvolved.



Publications

Elgamily M, Denewar M, Emam A et al. Injectable GSK3 inhibitor (Tideglusib) hydrogel versus enhanced β -tricalcium phosphate in tibial bone healing (in vitro and in vivo study). *Odontology* 2025-09-21 [PMID: 40975852]

Anerillas, LO;Wiberg, M;Kingham, PJ;Kelk, P; Platelet lysate for expansion or osteogenic differentiation of bone marrow mesenchymal stem cells for 3D tissue constructs *Regenerative therapy* 2023-12-01 [PMID: 37588134] (Western Blot, Human)

G Feng, P Zhang, J Huang, Y Yu, F Yang, X Zhao, W Wang, D Li, S Sun, X Niu, L Chai, J Li Sequential Release of Panax Notoginseng Saponins and Osteopractic Total Flavone from Poly (L-Lactic Acid) Scaffold for Treating Glucocorticoid-Associated Osteonecrosis of Femoral Head *Journal of functional biomaterials*, 2023-01-04;14(1):. 2023-01-04 [PMID: 36662078] (Western Blot, Human)

Shen JF, Fan ZB, Wu CW et al. Sacubitril Valsartan Enhances Cardiac Function and Alleviates Myocardial Infarction in Rats through a SUV39H1/SPP1 Axis *Oxidative Medicine and Cellular Longevity* 2022-09-22 [PMID: 36193085] (Western Blot, Human)

V?ru? RM, Melinte PR, P?rvu AS et al. Calcium fructoborate coating of titanium-hydroxyapatite implants by chemisorption deposition improves implant osseointegration in the femur of New Zealand White rabbit experimental model *Romanian Journal of Morphology and Embryology* 2021-07-01 [PMID: 34171072] (Western Blot, Human)

Zhao, L;Wang, Z;Tan, Y;Ma, J;Huang, W;Zhang, X;Jin, C;Zhang, T;Liu, W;Yang, YG; IL-17A/CEBP?/OPN/LYVE-1 axis inhibits anti-tumor immunity by promoting tumor-associated tissue-resident macrophages *Cell reports* 2024-12-05 [PMID: 39643970]

Liao H, Zou Z, Liu W et Al. Osteopontin-integrin signaling positively regulates neuroplasticity through enhancing neural autophagy in the peri-infarct area after ischemic stroke *Am J Transl Res* 2022-11-15 [PMID: 36505285]

Abou-Rjeileh U, Dos Santos Neto JM, Chirivi M et al. Oleic acid abomasal infusion limits lipolysis and improves insulin sensitivity in adipose tissue from periparturient dairy cows *Journal of dairy science* 2023-04-25 [PMID: 37105874] (Simple Western, Bovine)

Details:

Dilutions 1:50

Li, Jiabin, J Self-healing hybrid hydrogels with sustained bioactive components release for guided bone regeneration *Research Square* 2022-11-29 [PMID: 36814282] (ICC/IF, Mouse)

Park S, Park M, Saleh T, Lee B Autologous stromal vascular fraction-loaded hyaluronic acid/gelatin-biphasic calcium phosphate scaffold for bone tissue regeneration *Mater Sci Eng C Mater Biol Appl* 2022-02-12 [PMID: 35148865]

Kang H, Makkar P, Padalhin A et al. Comparative study on biodegradation and biocompatibility of multichannel calcium phosphate based bone substitutes *Mater Sci Eng C Mater Biol Appl* 2020-03-25 [PMID: 32204008]

Singh A, Kumar S, Acharya T et al. Application of nanohydroxyapatite-polysaccharide based biomaterial for bone cell mineralization in tissue engineering *Materials Today Communications* 2022-06-01 (ICC/IF, Human)

More publications at <http://www.novusbio.com/NB110-89062>

Procedures

Western Blot Protocol Specific for Osteopontin Antibody (1B20)

Western Blot Protocol

1. Perform SDS-PAGE on samples to be analyzed, loading 30 ug of total protein per lane.
2. Transfer proteins to membrane according to the instructions provided by the manufacturer of the membrane and transfer apparatus.
3. Stain according to standard Ponceau S procedure (or similar product) to assess transfer success, and mark molecular weight standards where appropriate.
4. Rinse the blot.
5. Block the membrane using standard blocking buffer for at least 1 hour.
6. Wash the membrane in wash buffer three times for 10 minutes each.
7. Dilute primary antibody in blocking buffer and incubate 1 hour at room temperature.
8. Wash the membrane in wash buffer three times for 10 minutes each.
9. Apply the diluted HRP conjugated secondary antibody in blocking buffer (as per manufacturers instructions) and incubate 1 hour at room temperature.
10. Wash the blot in wash buffer three times for 10 minutes each (this step can be repeated as required to reduce background).
11. Apply the detection reagent of choice in accordance with the manufacturers instructions.

Note: Tween-20 can be added to the blocking or antibody dilution buffer at a final concentration of 0.05-0.2%.



Immunohistochemistry-Paraffin protocol for Osteopontin/OPN Antibody (NB110-89062)

Immunohistochemistry-Paraffin Embedded Sections

Antigen Unmasking:

Bring slides to a boil in 10 mM sodium citrate buffer (pH 6.0) then maintain at a sub-boiling temperature for 10 minutes. Cool slides on bench-top for 30 minutes.

Staining:

1. Wash sections in deionized water three times for 5 minutes each.
2. Wash sections in wash buffer for 5 minutes.
3. Block each section with 100-400 ul blocking solution for 1 hour at room temperature.
4. Remove blocking solution and add 100-400 ul diluted primary antibody. Incubate overnight at 4C.
5. Remove antibody solution and wash sections in wash buffer three times for 5 minutes each.
6. Add 100-400 ul biotinylated diluted secondary antibody. Incubate 30 minutes at room temperature.
7. Remove secondary antibody solution and wash sections three times with wash buffer for 5 minutes each.
8. Add 100-400 ul Streptavidin-HRP reagent to each section and incubate for 30 minutes at room temperature.
9. Wash sections three times in wash buffer for 5 minutes each.
10. Add 100-400 ul DAB substrate to each section and monitor staining closely.
11. As soon as the sections develop, immerse slides in deionized water.
12. Counterstain sections in hematoxylin.
13. Wash sections in deionized water two times for 5 minutes each.
14. Dehydrate sections.
15. Mount coverslips.

Immunocytochemistry Protocol

Culture cells to appropriate density in 35 mm culture dishes or 6-well plates.

1. Remove culture medium and add 10% formalin to the dish. Fix at room temperature for 30 minutes.
2. Remove the formalin and add ice cold methanol. Incubate for 5-10 minutes.
3. Remove methanol and add washing solution (i.e. PBS). Be sure to not let the specimen dry out. Wash three times for 10 minutes.
4. To block nonspecific antibody binding incubate in 10% normal goat serum from 1 hour to overnight at room temperature.
5. Add primary antibody at appropriate dilution and incubate at room temperature from 2 hours to overnight at room temperature.
6. Remove primary antibody and replace with washing solution. Wash three times for 10 minutes.
7. Add secondary antibody at appropriate dilution. Incubate for 1 hour at room temperature.
8. Remove antibody and replace with wash solution, then wash for 10 minutes. Add Hoechst 33258 to wash solution at 1:25,000 and incubate for 10 minutes. Wash a third time for 10 minutes.
9. Cells can be viewed directly after washing. The plates can also be stored in PBS containing Azide covered in Parafilm (TM). Cells can also be cover-slipped using Fluoromount, with appropriate sealing.

*The above information is only intended as a guide. The researcher should determine what protocol best meets their needs. Please follow safe laboratory procedures."



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NB110-89062

NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB7539	Goat anti-Mouse IgG (H+L) Secondary Antibody [HRP]
NBP1-97005-0.5mg	Mouse IgG1 Isotype Control (MG1)

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB110-89062

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

