

Product Datasheet

DDX6 Antibody NB200-191

Unit Size: 0.1 ml

Store at 4C. Do not freeze.

www.novusbio.com



technical@novusbio.com

Reviews: 1 **Publications: 45**

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB200-191

Updated 9/9/2025 v.20.1

**Earn rewards for product
reviews and publications.**

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB200-191



NB200-191

DDX6 Antibody


Product Information	
Unit Size	0.1 ml
Concentration	0.2 mg/ml
Storage	Store at 4C. Do not freeze.
Clonality	Polyclonal
Preservative	0.09% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	TBS, 0.1% BSA

Product Description	
Description	Novus Biologicals Rabbit DDX6 Antibody (NB200-191) is a polyclonal antibody validated for use in IHC, WB, ICC/IF and IP. Anti-DDX6 Antibody: Cited in 42 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	1656
Gene Symbol	DDX6
Species	Human, Mouse, Hamster
Reactivity Notes	Hamster reactivity reported from a verified customer review.
Immunogen	The immunogen recognized by NB200-191 maps to a region between residues 1 and 50 of human DEAD (Asp-Glu-Ala-Asp) box polypeptide 6 using the numbering given Swiss-Prot entry P26196 (GeneID 1656).


Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunoprecipitation
Recommended Dilutions	Western Blot 1:5000-1:15000, Immunohistochemistry 1:100-1:500, Immunocytochemistry/ Immunofluorescence, Immunoprecipitation 2-5 ug/mg lysate, Immunohistochemistry-Paraffin 1:100-1:500
Application Notes	ICC/IF was reported in scientific literature (PMID: 25800057) and a verified customer review. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use. Epitope retrieval with Tris-EDTA pH 9.0 is recommended for FFPE tissue sections.

Images


Western Blot: DDX6 Antibody [NB200-191] - Western blot analysis of LBCs show increased Staufen (STAU1) (NBP1-33202) levels compared with normal controls. DDX6 (NB200-191) levels are unchanged. HD and SCA3 patient (polyQ expanded) FBs were used as additional controls. Four normal and five SCA2 FBs, and two normal and three SCA2 LBCs were used. Image collected and cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/30194296/>) licensed under a CC-BY license.




Immunocytochemistry/Immunofluorescence: DDX6 Antibody [NB200-191] - In normal condition, cells were fixed and stained for DDX6, that usually punctuate in process bodies (PBs). Primary Antibody rabbit-anti-DDX6 diluted 1:1000. Image submitted by a verified customer review.




Immunohistochemistry-Paraffin: DDX6 Antibody [NB200-191] - Section of human colon carcinoma (left) and mouse renal cell carcinoma (right). Antibody: Affinity purified rabbit anti-DDX6 used at a dilution of 1:200 (1ug/ml). Detection: DAB.



Immunoprecipitation: DDX6 Antibody [NB200-191] - Detection of human DDX6 by Western Blot and Immunoprecipitation. Samples: Whole cell lysate (15 and 50 ug) from HeLa cells. Antibodies: Affinity purified rabbit anti-DDX6 antibody NB200-191 used at 0.04 ug/ml for WB (A and B) and at 3 ug/mg lysate for IP. DDX6 was also immunoprecipitated using rabbit anti-DDX6 antibody NB200-192, but not with 'DDX6 Ab', which is to another epitope from DDX6. Normal rabbit IgG was used as a negative control. Detection: Chemiluminescence with an exposure time of 1 second (A & B).



Western Blot: DDX6 Antibody [NB200-191] - Staufen1 protein but not mRNA steady-state levels are increased in neurodegenerative disease cells & tissues. Western blot analysis of SCA2- FBs (a) & LBCs (b) show increased STAU1 levels compared with normal controls. DDX6 levels are unchanged. HD & SCA3 patient (polyQ expanded) FBs were used as additional controls. Four normal & five SCA2 FBs, & two normal & three SCA2 LBCs were used. c, d Western blot analyses of ATXN2Q127 (c) & BAC-Q72 (d) mouse cerebellar extracts (24 weeks of age) showing increased Stau1 levels compared with wild-type or BAC-Q22 controls (n = 2–3 animals per group). e Western blot of FB extracts from an ALS patient with the TDP-43G298S mutation show increased STAU1 levels. β -Actin was used as loading control & representative blots of three independent experiments are shown. f–h STAU1 RNA levels are unaltered in SCA2 & ALS cells & SCA2 mice. qRT-PCR analyses of STAU1 mRNA in SCA2 FBs & ALS FB with TDP-43G298S mutation (f) or SCA2 LBCs (g). h qRT-PCR analyses of cerebellar RNAs from ATXN2Q127 & BAC-Q72 mice compared to wild-type littermates (24 weeks of age; n = animals per group). Gene expression levels were normalized to Actb Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/30194296>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Publications

Heim A, Cheng S, Orth J et al. Translational repression by 4E-T is crucial to maintain the prophase-I arrest in vertebrate oocytes. *Nature Communications* 2025-08-28 [PMID: 40877279]

Venkataramanan S, Gadek M, Calviello L et al. DDX3X and DDX3Y are redundant in protein synthesis RNA 2021-12-01 [PMID: 34535544]

Vladimir Majerciak, Tongqing Zhou, Michael J Kruhlak, Zhi-Ming Zheng RNA helicase DDX6 and scaffold protein GW182 in P-bodies promote biogenesis of stress granules *Nucleic Acids Research* 2023-09-22 [PMID: 37427791]

Heim A, Niedermeier ML, Stengel F, Mayer TU The translation regulator Zar1l controls timing of meiosis in *Xenopus* oocytes *Development (Cambridge, England)* 2022-11-01 [PMID: 36278895]

Mehravar M, Kumar Y, Olshansky M et al. MOV10 facilitates messenger RNA decay in an N6-methyladenosine (m6A) dependent manner to maintain the mouse embryonic stem cells state *bioRxiv* 2021-08-12

Sung CC, Poll B, Lin SH et al. Early Molecular Events Mediating Loss of Aquaporin-2 during Ureteral Obstruction in Rat *Journal of the American Society of Nephrology : JASN* 2022-08-02 [PMID: 35918145] (IHC-P, Rat)

Details:

Dilutions: 1:500

Hallacli E, Kayatekin C, Nazeen S et al. The Parkinson's disease protein alpha-synuclein is a modulator of processing bodies and mRNA stability *Cell* 2022-06-09 [PMID: 35688132] (ICC/IF, WB, Human)

Shah A, Bhandari R IP6K1 upregulates the formation of processing bodies by influencing protein-protein interactions on the mRNA cap *Journal of cell science* 2021-11-29 [PMID: 34841428]

Somasekharan SP, Gleave M SARS-CoV-2 nucleocapsid protein interacts with immunoregulators and stress granules and phase separates to form liquid droplets *FEBS letters* 2021-11-15 [PMID: 34780058] (ICC/IF, Human)

Rink MR, Baptista MAP, Flomm FJ Et al. Concatemeric Broccoli reduces mRNA stability and induces aggregates *PloS one* 2021-08-04 [PMID: 34347781] (ICC/IF)

Shah A, Bhandari R IP6K1 upregulates the formation of processing bodies by promoting proteome remodeling on the mRNA cap *bioRxiv* 2020-07-14 [PMID: 34841428] (WB, ICC/IF, Human)

Cougnoux A, Drummond RA, Fellmeth M, et al. GC content shapes mRNA storage and decay in human cells *Elife* 2019-12-19 [PMID: 31855182] (WB, ICC/IF, Human)

More publications at <http://www.novusbio.com/NB200-191>





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NB200-191

NBL1-09810	DDX6 Overexpression Lysate
NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB200-191

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

