

# Product Datasheet

## O-GlcNAc Antibody (RL2) - BSA Free NB300-524

Unit Size: 0.1 ml

Store at -20C. Avoid freeze-thaw cycles.

[www.novusbio.com](http://www.novusbio.com)



[technical@novusbio.com](mailto:technical@novusbio.com)

**Reviews: 2 Publications: 35**

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:  
[www.novusbio.com/NB300-524](http://www.novusbio.com/NB300-524)

Updated 9/9/2025 v.20.1

**Earn rewards for product  
reviews and publications.**

Submit a publication at [www.novusbio.com/publications](http://www.novusbio.com/publications)

Submit a review at [www.novusbio.com/reviews/destination/NB300-524](http://www.novusbio.com/reviews/destination/NB300-524)



**NB300-524**

O-GlcNAc Antibody (RL2) - BSA Free

Product Information	
Unit Size	0.1 ml
Concentration	1.0 mg/ml
Storage	Store at -20C. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	RL2
Preservative	0.02% Sodium Azide
Isotype	IgG1
Purity	Protein A purified
Buffer	PBS


Product Description	
Description	Novus Biologicals Mouse O-GlcNAc Antibody (RL2) - BSA Free (NB300-524) is a monoclonal antibody validated for use in IHC, WB, ELISA, Flow, ICC/IF, IP and ChIP. Anti-O-GlcNAc Antibody: Cited in 30 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Mouse
Species	Human, Mouse, Rat, Porcine, Bovine, Drosophila, Fish, Hamster, Primate, Virus, Xenopus
Reactivity Notes	Porcine reactivity reported in scientific literature (PMID: 26004176). Xenopus reactivity reported in scientific literature (PMID: 17329255). Please note that this antibody is reactive to Mouse and derived from the same host, Mouse. Additional Mouse on Mouse blocking steps may be required for IHC and ICC experiments. Please contact Technical Support for more information.
Specificity/Sensitivity	Detects nuclear pore complex (NPC), cytoplasmic and intranuclear O-linked glycoproteins from human, mouse, and rat tissues.
Immunogen	Pore complex-lamina fraction purified from rat liver nuclear envelopes.

Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, Chromatin Immunoprecipitation, Dot Blot, ELISA, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunoprecipitation, Chromatin Immunoprecipitation (ChIP)
Recommended Dilutions	Western Blot 1:1000, Chromatin Immunoprecipitation 1:10 - 1:500. Use reported in scientific literature (PMID 20404350), Flow Cytometry 1:10 - 1:1000, ELISA 1:100 - 1:2000. Use reported in scientific literature (PMID 12029848), Immunohistochemistry 1:10 - 1:500, Immunocytochemistry/ Immunofluorescence, Immunoprecipitation 1:10 - 1:500, Immunohistochemistry-Paraffin 1:200, Dot Blot 1:800, Chromatin Immunoprecipitation (ChIP) 1:10-1:500




## Images


**Immunohistochemistry-Paraffin: O-GlcNAc Antibody (RL2) [NB300-524] -** Analysis of a FFPE tissue section of the mouse colon using 1:200 dilution of O-GlcNAc [RL2] antibody (NB300-524). The signal was developed using HRP-DAB method which followed counterstaining of the cells with hematoxylin.




**Western Blot: O-GlcNAc Antibody (RL2) [NB300-524] - Impact of OGA inhibition by thiamet G in primary cultures of NCM.** Representative western blots (left panel) and quantification (right panel) of O-GlcNAcylated proteins levels in control (Ctrl) and NCM treated with 100 nM of thiamet G (TG) during 24 h (n = 12). Image collected and cropped by Citeab from the following publication (Interplay Between Phosphorylation and O-GlcNAcylation of Sarcomeric Proteins in Ischemic Heart Failure. Front Endocrinol (Lausanne) (2018) licensed under a CC-BY license.




**Immunocytochemistry/Immunofluorescence: O-GlcNAc Antibody (RL2) [NB300-524] -** Neuro2a cells were fixed for 10 minutes using 10% formalin and then permeabilized for 5 minutes using 1X PBS + 0.5% Triton X-100. The cells were incubated with anti-O-GlcNAc (RL2) at 5 ug/mL overnight at 4C and detected with an anti-mouse Dylight 488 (Green) at a 1:500 dilution. Nuclei were counterstained with DAPI (Blue). Cells were imaged using a 40X objective.




**Flow Cytometry: O-GlcNAc Antibody (RL2) [NB300-524] -** An intracellular stain was performed on Neuro2a cells with O-GlcNAc Antibody [RL2] NB300-524 (blue) and a matched isotype control (orange). Cells were fixed with 4% PFA and then permeabilized with 0.1% saponin. Cells were incubated in an antibody dilution of 1.0 ug/mL for 30 minutes at room temperature, followed by Mouse IgG (H+L) Cross-Adsorbed Secondary Antibody, Dylight 550 (35503, Thermo Fisher).




**Western Blot: O-GlcNAc Antibody (RL2) [NB300-524]** - Analysis of mouse cortical brain lysates using O-Linked N-Acetylglucosamine Monoclonal Antibody. Blots containing cortical extracts from 4 individual C57BL/6 mice (Lanes 1-4) were blocked with 5% milk in TBST, and probed with MA1-072 at 1:1000, followed by a fluorophore-conjugated goat anti-mouse IgG secondary antibody. Data courtesy of the Innovators Program.




**Immunocytochemistry/Immunofluorescence: O-GlcNAc Antibody (RL2) [NB300-524]** - HeLa cells were fixed for 10 minutes using 10% formalin and then permeabilized for 5 minutes using 1X PBS + 0.5% Triton X-100. The cells were incubated with anti-O-GlcNAc (RL2) at 5 ug/mL overnight at 4C and detected with an anti-mouse Dylight 488 (Green) at a 1:500 dilution. Nuclei were counterstained with DAPI (Blue). Cells were imaged using a 40X objective.




**Flow Cytometry: O-GlcNAc Antibody (RL2) [NB300-524]** - Analysis using Alexa Fluor (R) 647 conjugate of NB300-524. An intracellular stain was performed on Jurkat cells with O-GlcNAc antibody (RL2) NB300-524 (blue) and a matched isotype control NBP2-27287 (orange). Cells were fixed with 4% PFA and then permeablized with 0.1% saponin. 1 ug of antibody was added to 100 uL of staining buffer and cells were incubated for 30 minutes at room temperature. Both antibodies were conjugated to Alexa Fluor 647.



**Flow Cytometry: O-GlcNAc Antibody (RL2) [NB300-524]** - An intracellular stain was performed on U-937 cells with O-GlcNAc antibody (RL2) NB300-524AF647 (blue) and a matched isotype control. Cells were fixed with 4% PFA and then permeablized with 0.1% saponin. Cells were incubated in an antibody dilution of 2.5 ug/mL for 30 minutes at room temperature. Both antibodies were conjugated to Alexa Fluor 647.




Flow Cytometry: O-GlcNAc Antibody (RL2) [NB300-524] - An intracellular stain was performed on Jurkat cells with O-GlcNAc antibody (RL2) NB300-524PE (blue) and a matched isotype control. Cells were fixed with 4% PFA and then permeabilized with 0.1% saponin. Cells were incubated in an antibody dilution of 2.5 ug/mL for 30 minutes at room temperature. Both antibodies were conjugated to Phycoerythrin.




Copyright © 2018 Novus Biologicals

Flow Cytometry: O-GlcNAc Antibody (RL2) [NB300-524] - An intracellular stain was performed on SK-MEL-28 cells with O-GlcNAc antibody (RL2) NB300-524AF647 (blue) and a matched isotype control. Cells were fixed with 4% PFA and then permeabilized with 0.1% saponin. Cells were incubated in an antibody dilution of 2.5 ug/mL for 30 minutes at room temperature. Both antibodies were conjugated to Alexa Fluor 647.




Copyright © 2018 Novus Biologicals

Flow Cytometry: O-GlcNAc Antibody (RL2) [NB300-524] - An intracellular stain was performed on HeLa cells with O-GlcNAc Antibody [RL2] Antibody NB300-524AF647 (blue) and a matched isotype control (orange). Cells were fixed with 4% PFA and then permeabilized with 0.1% saponin. Cells were incubated in an antibody dilution of 2.5 ug/mL for 30 minutes at room temperature. Both antibodies were conjugated to Alexa Fluor 647.




Copyright © 2018 Novus Biologicals

Flow Cytometry: O-GlcNAc Antibody (RL2) [NB300-524] - An intracellular stain was performed on Neuro2a cells with O-GlcNAc Antibody [RL2] NB300-524AF647 (blue) and a matched isotype control (orange). Cells were fixed with 4% PFA and then permeabilized with 0.1% saponin. Cells were incubated in an antibody dilution of 2.5 ug/mL for 30 minutes at room temperature. Both antibodies were conjugated to Alexa Fluor 647.




Copyright © 2019 Novus Biologicals


**Flow Cytometry: O-GlcNAc Antibody (RL2) [NB300-524]** - An intracellular stain was performed on RH30 cells with O-GlcNAc [RL2] Antibody NB300-524AF647 (blue) and a matched isotype control (orange). Cells were fixed with 4% PFA and then permeabilized with 0.1% saponin. Cells were incubated in an antibody dilution of 2.5 ug/mL for 30 minutes at room temperature. Both antibodies were conjugated to Alexa Fluor 647.




**Flow Cytometry: O-GlcNAc Antibody (RL2) [NB300-524]** - An intracellular stain was performed on Jurkat cells with O-GlcNAc Antibody [RL2] NB300-524 (blue) and a matched isotype control (orange). Cells were fixed with 4% PFA and then permeabilized with 0.1% saponin. Cells were incubated in an antibody dilution of 1.0 ug/mL for 30 minutes at room temperature, followed by Mouse IgG (H+L) Cross-Adsorbed Secondary Antibody, Dylight 550 (35503, Thermo Fisher).



**Western Blot: O-GlcNAc Antibody (RL2) - BSA Free [NB300-524]** - Analysis of O-GlcNAcylated LV proteins by Western blot & WGA-SDS-PAGE gel electrophoresis. (A) Red ponceau staining (left panel) & western blot (right panel) of O-GlcNAcylated proteins (50 µg) extracted from sham- & HF-rats treated or not with thiamet G. The positions of molecular weight are indicated as kilodalton (kDa) on the left. (B) Red ponceau staining (left panel) & WGA-SDS-PAGE of O-GlcNAcylated proteins levels (middle panel) of O-GlcNAcylated desmin levels (right panel) from the same samples. The arrow in desmin WGA gels indicates the non-O-GlcNAcylated form. Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/30344511>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



**Western Blot: O-GlcNAc Antibody (RL2) - BSA Free [NB300-524]** - Effect of OGA inhibition by thiamet G in isolated perfused heart. (A) Description of the protocol designed for thiamet G (TG) perfusion in sham- (n = 6) & HF- (n = 7) rats 6 weeks post-MI. (B) Western blot (left panel) & quantification (right panel) of O-GlcNAcylated proteins levels measured in proteins extracted from LVs of isolated perfused sham- & HF-rat hearts treated or not with 100 µM thiamet G for 2 h (n = 7 in each group). (C) Western blots (upper panel) & quantification (lower panel) of total desmin levels in the same samples. (D) Phosphorylation profiles of desmin were analyzed in the same samples by Phos-tag™ gel. Graphs show mean ± SEM values expressed in arbitrary units (A.U.). The positions of molecular weight are indicated as kilodalton (kDa) on the left. \*P < 0.05; \*\* < 0.01. Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/30344511>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



## Publications

Akihito Ishikita, Shouji Matsushima, Soichiro Ikeda, Kosuke Okabe, Ryohei Nishimura, Tomonori Tadokoro, Nobuyuki Enzan, Taishi Yamamoto, Masashi Sada, Yoshitomo Tsutsui, Ryo Miyake, Masataka Ikeda, Tomomi Ide, Shintaro Kinugawa, Hiroyuki Tsutsui GFAT2 mediates cardiac hypertrophy through HBP-O-GlcNAcylation-Akt pathway iScience 2021-12-17 [PMID: 34934932]

Hugo SE, Schlegel A., , et Al. A Genetic Model to Study Increased Hexosamine Biosynthetic Flux Endocrinology 2017-06-06 [PMID: 28582574]

Taub DG, Awal MR, Gabel CV., et Al. O-GlcNAc Signaling Orchestrates the Regenerative Response to Neuronal Injury in *Caenorhabditis elegans* Cell Rep 2018-08-23 [PMID: 30134155]

Shi Q, Shen Q, Liu Y et Al. Increased glucose metabolism in TAMs fuels O-GlcNAcylation of lysosomal Cathepsin B to promote cancer metastasis and chemoresistance Cancer Cell 2022-10-13 [PMID: 36084651]

Czajewski I, Swain B, Xu J et Al. Rescuable sleep and synaptogenesis phenotypes in a *Drosophila* model of O-GlcNAc transferase intellectual disability Elife 2024-11-13 [PMID: 39535175]

Florence Authier, Nina Ondruskova, Andrew T Ferenbach, Alison McNeilly, Daan M F van Aalten Neurodevelopmental defects in a mouse model of O-GlcNAc transferase intellectual disability. Disease models & mechanisms 2024-04-03 [PMID: 38566589]

Authier F, Attianese B, Bartual S et al. Intellectual disability and neurogenesis defects associated with increased turnover of an O-GlcNAcase variant medRxiv 2023-11-24 (WB)

Kadosaka T, Watanabe M, Natsui H et al. Empagliflozin attenuates arrhythmogenesis in diabetic cardiomyopathy by normalizing intracellular Ca<sup>2+</sup> handling in ventricular cardiomyocytes American journal of physiology. Heart and circulatory physiology 2023-03-01 [PMID: 36607794]

Murray M, Davidson L, Ferenbach A et al. Neuroectoderm phenotypes in a human stem cell model of O-GlcNAc transferase intellectual disability bioRxiv 2023-09-21 (WB, Human)

Details:  
1:1000 dilution

Authier F, Ondruskova N, Ferenbach A et al. Neurodevelopmental defects in a mouse model of O-GlcNAc transferase intellectual disability bioRxiv 2023-08-24 (WB, Mouse)

Feng Z, Wang T, Sun Y et al. Sulforaphane suppresses paraquat-induced oxidative damage in bovine  $\square$ in vitro-matured oocytes through Nrf2 transduction pathway Ecotoxicology and environmental safety 2023-04-01 [PMID: 36907095] (Western Blot, Bovine)

Czajewski I, McDowall L, Ferenbach A Et al Rescuable sleep and synaptogenesis phenotypes in a *Drosophila* model of O-GlcNAc transferase intellectual disability bioRxiv 2023-06-30 (WB)

More publications at <http://www.novusbio.com/NB300-524>

## Procedures

### Immunohistochemistry-Paraffin protocol for O-GlcNAc Antibody (NB300-524)

#### Immunohistochemistry-Paraffin Embedded Sections

##### Antigen Unmasking:

Bring slides to a boil in 10 mM sodium citrate buffer (pH 6.0) then maintain at a sub-boiling temperature for 10 minutes. Cool slides on bench-top for 30 minutes.

##### Staining:

1. Wash sections in deionized water three times for 5 minutes each.
2. Wash sections in wash buffer for 5 minutes.
3. Block each section with 100-400 ul blocking solution for 1 hour at room temperature.
4. Remove blocking solution and add 100-400 ul diluted primary antibody. Incubate overnight at 4 C.
5. Remove antibody solution and wash sections in wash buffer three times for 5 minutes each.
6. Add 100-400 ul biotinylated diluted secondary antibody. Incubate 30 minutes at room temperature.
7. Remove secondary antibody solution and wash sections three times with wash buffer for 5 minutes each.
8. Add 100-400 ul Streptavidin-HRP reagent to each section and incubate for 30 minutes at room temperature.
9. Wash sections three times in wash buffer for 5 minutes each.
10. Add 100-400 ul DAB substrate to each section and monitor staining closely.
11. As soon as the sections develop, immerse slides in deionized water.
12. Counterstain sections in hematoxylin.
13. Wash sections in deionized water two times for 5 minutes each.
14. Dehydrate sections.
15. Mount coverslips.





### **Novus Biologicals USA**

10730 E. Briarwood Avenue  
Centennial, CO 80112  
USA  
Phone: 303.730.1950  
Toll Free: 1.888.506.6887  
Fax: 303.730.1966  
nb-customerservice@bio-techne.com

### **Bio-Techne Canada**

21 Canmotor Ave  
Toronto, ON M8Z 4E6  
Canada  
Phone: 905.827.6400  
Toll Free: 855.668.8722  
Fax: 905.827.6402  
canada.inquires@bio-techne.com

### **Bio-Techne Ltd**

19 Barton Lane  
Abingdon Science Park  
Abingdon, OX14 3NB, United Kingdom  
Phone: (44) (0) 1235 529449  
Free Phone: 0800 37 34 15  
Fax: (44) (0) 1235 533420  
info.EMEA@bio-techne.com

### **General Contact Information**

www.novusbio.com  
Technical Support: nb-technical@bio-techne.com  
Orders: nb-customerservice@bio-techne.com  
General: novus@novusbio.com

### **Products Related to NB300-524**

---

NBL1-13919	O-GlcNAc Transferase p110 subunit Overexpression Lysate
NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB7539	Goat anti-Mouse IgG (H+L) Secondary Antibody [HRP]
NBP1-97005-0.5mg	Mouse IgG1 Isotype Control (MG1)

---

### **Limitations**

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit [www.novusbio.com/guarantee](http://www.novusbio.com/guarantee)

Earn gift cards/discounts by submitting a review: [www.novusbio.com/reviews/submit/NB300-524](http://www.novusbio.com/reviews/submit/NB300-524)

Earn gift cards/discounts by submitting a publication using this product:

[www.novusbio.com/publications](http://www.novusbio.com/publications)

