

Product Datasheet

Ki67/MKI67 Antibody - BSA Free NB500-170

Unit Size: 0.1 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Reviews: 8 Publications: 196

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB500-170

Updated 9/9/2025 v.20.1

**Earn rewards for product
reviews and publications.**

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB500-170



NB500-170


Ki67/MKI67 Antibody - BSA Free

Product Information	
Unit Size	0.1 ml
Concentration	1.0 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.05% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	PBS
Target Molecular Weight	359 kDa
Product Description	
Description	Novus Biologicals Knockout (KO) Validated Rabbit Ki67/MKI67 Antibody - BSA Free (NB500-170) is a polyclonal antibody validated for use in IHC, WB, Flow, ICC/IF and IP. Anti-Ki67/MKI67 Antibody: Cited in 185 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	4288
Gene Symbol	MKI67
Species	Human, Mouse, Rat, Porcine, Avian
Reactivity Notes	Use in Avian reported in scientific literature (PMID:34986451) Ki67/MKI67 Antibody reacted with Mouse (PMID: 22033079), Rat (PMID: 22521325), and Porcine (PMID: 29146772). .
Marker	Proliferation Marker
Immunogen	The immunogen for this KI67/MKI67 Antibody was made using a synthetic peptide from the internal region of Human KI67/MKI67, between amino acids: 1550-1600 [Uniprot: P46013].
Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, Flow Cytometry, Immunoblotting, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Frozen, Immunoprecipitation, Knockout Validated
Recommended Dilutions	Western Blot 1-2 ug/ml, Flow Cytometry reported in scientific literature (PMID 34946963), Immunohistochemistry 1:50-1:300, Immunocytochemistry/ Immunofluorescence 1:20-1:100, Immunoprecipitation reported by customer review, Immunohistochemistry-Paraffin 1:50-1:300, Immunohistochemistry-Frozen 1:50-1:300. Use reported in scientific literature (PMID 22493503), Immunoblotting reported in scientific literature (PMID 24248265), Knockout Validated
Application Notes	Formalin fixed paraffin embedded tissue sections require high temperature antigen unmasking with 10 mM citrate buffer (pH 6.0) prior to immunostaining. This antibody will not work without optimal antigen retrieval, and is a critical step. NOTE: We suggest an incubation period of 30 minutes at room temperature and to use DAB to stain the protein.




Images


Immunocytochemistry/Immunofluorescence: Ki67/MKI67 Antibody [NB500-170] - HeLa cells were fixed in 4% paraformaldehyde for 10 minutes and permeabilized in 0.5% Triton X-100 in PBS for 5 minutes. The cells were incubated with anti- NB500-170 at 2 ug/ml overnight at 4C and detected with an anti-rabbit Dylight 488 (Green) at a 1:1000 dilution for 60 minutes. Alpha tubulin (DM1A) NB100-690 was used as a co-stain at a 1:1000 dilution and detected with an anti-mouse Dylight 550 (Red) at a 1:1000 dilution. Nuclei were counterstained with DAPI (Blue). Cells were imaged using a 100X objective and digitally deconvolved.




Simple Western: Ki67/MKI67 Antibody [NB500-170] - Detection of Ki67/MKI67 by Simple Western™. Simple Western lane view shows lysates of HeLa parental cell line and Ki67 knockout (KO) HeLa cell line. A specific band was detected for Ki67/MKI67 at approximately 320 kDa (as indicated) in the parental cell line, but is not detectable in the knockout HeLa cell line using 20 ug/mL of Rabbit Anti-Ki67/MKI67 Polyclonal Antibody (Catalog # NB500-170). GAPDH is shown as a loading control. This experiment was conducted under reducing conditions and using the 12-230 kDa separation system.




Immunohistochemistry: Ki67/MKI67 Antibody [NB500-170] - Senescent MSCs are characterized by loss of nestin expression. Double-immunofluorescence images of femoral metaphysis sections from 4-, 6-, 8-, and 12-week-old male Nestin-GFP mice using antibodies against GFP (green) and Ki67 (red). DAPI stains nuclei blue. GP growth plate. PS primary spongiosa. Quantification of the percentage of GFP+ cells that express Ki67. Image collected and cropped by CiteAb from the following publication (<https://www.nature.com/articles/s41467-017-01509-0>) licensed under a CC-BY license.




Immunohistochemistry-Paraffin: Ki67/MKI67 Antibody [NB500-170] - Ki67/MKI67 Antibody [NB500-170] - Tissue section of human tonsil using 1:50 dilution of rabbit anti-Ki67 antibody. The staining was developed with HRP labeled anti-rabbit IgG secondary antibody and DAB reagent, and nuclei of cells were counter-stained with hematoxylin. This Ki67 antibody generated a specific nuclear staining in the cells in germinal centers of the tested tonsil tissue.




Immunohistochemistry-Paraffin: Ki67/MKI67 Antibody [NB500-170] - Ki67/MKI67 Antibody [NB500-170] - Human tonsil using 1:200 dilution of rabbit anti-Ki67 antibody. The staining was developed with HRP labeled anti-rabbit IgG secondary antibody and DAB reagent, and nuclei of cells were counter-stained with hematoxylin. This Ki67 antibody generated a specific nuclear staining in the cells in germinal centers of the tested tonsil tissue.




Western Blot: Ki67/MKI67 Antibody - BSA Free [NB500-170] - Whole cell lysates from SUM159, MDA-MB-231 and MCF-7 cells were loaded with 30 ug/lane. 10% SDS-PAGE. Ki67/MKI67 antibody (NB500-170) was used for primary antibody: 1:2000, 4C, overnight. Image from verified customer review.




Immunocytochemistry/Immunofluorescence: Ki67/MKI67 Antibody [NB500-170] - NIH3T3 cells were fixed in 4% paraformaldehyde for 10 minutes and permeabilized in 0.5% Triton X-100 in PBS for 5 minutes. The cells were incubated with anti-Ki67/MKI67 Antibody NB500-170 at 2 ug/ml overnight at 4C and detected with an anti-rabbit Dylight 488 (Green) at a 1:1000 dilution for 60 minutes. Nuclei were counterstained with DAPI (Blue). Cells were imaged using a 100X objective and digitally deconvolved.




Flow Cytometry: Ki67/MKI67 Antibody - BSA Free [NB500-170] - An intracellular stain was performed on U-251 MG cells with Ki67/MKI67 Antibody NB500-170AF594 (blue) and a matched isotype control NBP2-24891 (orange). Cells were fixed with 4% PFA and then permeabilized with 0.1% saponin. Cells were incubated in an antibody dilution of 2.5 ug/mL for 30 minutes at room temperature. Both antibodies were conjugated to Alexa Fluor 594.




Western Blot: Ki67/MKI67 Antibody [NB500-170] - Ki-67/MKI67 Antibody [NB500-170] - Analysis of A431 (A) and Hek293 (B) cell lysate using Ki67 antibody (NB500-170) at 2 ug/ml.




Immunocytochemistry/Immunofluorescence: Ki67/MKI67 Antibody [NB500-170] - Ki67 antibody was tested at 1:25 in HeLa cells with FITC (green). Nuclei and actin were counterstained with Dapi (blue) and Phalloidin (red).




Immunocytochemistry/Immunofluorescence: Ki67/MKI67 Antibody [NB500-170] - Confocal immunofluorescent analysis of MCF7 cells using Ki67 antibody (NB500-170, 1:5). An Alexa Fluor 488-conjugated Goat to rabbit IgG was used as secondary antibody (green). Actin filaments were labeled with Alexa Fluor 568 phalloidin (red). DAPI was used to stain the cell nuclei (blue).




Immunocytochemistry/Immunofluorescence: Ki67/MKI67 Antibody [NB500-170] - A431 cells were fixed in 4% paraformaldehyde for 10 minutes and permeabilized in 0.5% Triton X-100 in PBS for 5 minutes. The cells were incubated with anti- NB500-170 at 2 ug/ml overnight at 4C and detected with an anti-rabbit Dylight 488 (Green) at a 1:1000 dilution for 60 minutes. Alpha tubulin (DM1A) NB100-690 was used as a co-stain at a 1:1000 dilution and detected with an anti-mouse Dylight 550 (Red) at a 1:1000 dilution. Nuclei were counterstained with DAPI (Blue). Cells were imaged using a 100X objective and digitally deconvolved.




Immunohistochemistry-Paraffin: Ki67/MKI67 Antibody [NB500-170] - Ki67/MKI67 Antibody [NB500-170] - Human tonsil.




Ki67/MKI67 was detected in immersion fixed MCF7 human breast cancer cell line using Rabbit anti-Ki67/MKI67 Affinity Purified Polyclonal Antibody conjugated to Biotin (Catalog # NB500-170B) at 5 µg/mL overnight at 4C. Cells were stained using Streptavidin conjugated to DyLight 550 (red) and counterstained with DAPI (blue). Cells were imaged using a 100X objective and digitally deconvolved.



Ki67/MKI67 was detected in immersion fixed A431 human skin carcinoma cell line using Rabbit anti- Ki67/MKI67 Affinity Purified Polyclonal Antibody conjugated to Alexa Fluor® 488 (Catalog # NB500-170AF488) (green) at 2 µg/mL overnight at 4C. Cells were counterstained with DAPI (blue). Cells were imaged using a 100X objective and digitally deconvolved.




An intracellular stain was performed on A431 human skin carcinoma cell line using Rabbit anti-Ki67/MKI67 Affinity Purified Polyclonal Antibody conjugated to DyLight 650 (Catalog # NB500-170C, blue histogram) or matched control antibody (orange histogram) at 2.5 µg/mL for 30 minutes at RT.




Ki67/MKI67 was detected in immersion fixed A431 human skin carcinoma cell line using Rabbit anti- Ki67/MKI67 Affinity Purified Polyclonal Antibody conjugated to DyLight 488 (Catalog # NB500-170G) (green) at 5 $\mu\text{g}/\text{mL}$ overnight at 4C. Cells were counterstained with DAPI (blue). Cells were imaged using a 100X objective and digitally deconvolved.




Ki67/MKI67 was detected in immersion fixed U-251 MG human glioblastoma cell line using Rabbit anti- Ki67/MKI67 Affinity Purified Polyclonal Antibody conjugated to DyLight 650 (Catalog # NB500-170C) (light blue) at 5 $\mu\text{g}/\text{mL}$ overnight at 4C. Cells were counterstained with DAPI (blue). Cells were imaged using a 100X objective and digitally deconvolved.




Immunohistochemistry: Ki67/MKI67 Antibody - BSA Free [NB500-170] - PKD2 enzymatic deficiency had no effect on the proliferation or apoptosis of intestinal epithelial cells in mice exposed to DSS. (a) Representative 200 \times immunohistochemical images (left) & quantification of Ki-67+ proliferating intestinal epithelial cells (right) in wild-type & PKD2SSAA/SSAA mice. Bar represents 5 μm . (b) Representative 200 \times immunohistochemical images (left) & quantification of cleaved Caspase-3+ apoptotic intestinal epithelial cells (right) in wild-type & PKD2SSAA/SSAA mice. Bar represents 5 μm . Results are representative of three separate experiments with three mice per group. All $p > 0.05$ as measured by one way-ANOVA among four groups. Image collected & cropped by CiteAb from the following publication (<https://www.nature.com/articles/srep34079>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Immunohistochemistry: Ki67/MKI67 Antibody - BSA Free [NB500-170] - Deletion of Vangl2, but not Vangl1, in the mammary gland results in narrow ducts & low cell turnover. (A) RT-qPCR analysis of Vangl1 & Vangl2 mRNA levels in the mammary glands of MMTV-Cre, MMTV-Cre;Vangl2flox/+ & MMTV-Cre;Vangl2flox/flox at 10 weeks of age. (B) Representative images of 10 week old, carmine stained mammary glands & quantification of total branch number from MMTV-Cre, MMTV-Cre;Vangl2flox/+ & MMTV-Cre;Vangl2flox/flox mice (n = 3 mice per genotype). (C) Histological analysis by H&E staining show normal duct formation in 10 week old MMTV-Cre & MMTV-Cre;Vangl2flox/+ glands, whereas ducts from MMTV-Cre;Vangl2flox/flox glands are narrow & have significantly diminished lumens (n = 3 mice/genotype). (D) Immunostaining in mammary ducts show normal distribution of luminal (Cytokeratin 8 (K8), green), & basal (K5), red; SMA magenta) cell populations. (E) Immunofluorescence of adhesion junctional protein E-Cadherin (CDH1, green) & apical membrane marker pERM (red). (F,G) Immunostaining in 10 week old MMTV-Cre;Vangl2+/+ & MMTV-Cre;Vangl2flox/flox glands & quantitation of (F) Ki67 & (G) Cleaved Caspase 3 (n = 3–5 mice per genotype). Lu indicates lumen. * indicates non-magnified duct. Data are represented as mean \pm SEM. Scale bar represents 100 μ m (C,F,G) 50 μ m (D) & 10 μ m (E). Student's t-test *p < 0.05 & ****p < 0.0001. Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/31068622>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Immunocytochemistry/ Immunofluorescence: Ki67/MKI67 Antibody - BSA Free [NB500-170] - Senescent MSPCs are characterized by loss of nestin expression. a, b Representative images of GFP immunofluorescence staining (green) & quantitative analysis of GFP+ cells in femoral primary spongiosa from 4, 6, 8, & 12-week-old male Nestin-GFP mice. Images in upper panels in a are lower power w/ boxes outlining area of higher power in lower panels. Numbers of GFP+ cells per mm² tissue area in primary spongiosa (N. GFP+ cells/PS.Ar) b. 4W, 6W, 8W, & 12W represent 4-, 6-, 8-, & 12-week-old mice, respectively. DAPI stains nuclei blue. c, d Representative images of flow cytometry analysis (c) & % of CD45-GFP+ cells (d) in femoral metaphysis from 4-, 6-, 8-, & 12-week-old male Nestin-GFP mice. e, f Double-immunofluorescence images of femoral metaphysis sections from 4-, 6-, 8-, & 12-week-old male Nestin-GFP mice using antibodies against GFP (green) & Ki67 (red) e. DAPI stains nuclei blue. GP growth plate. PS primary spongiosa. Quantification of % of GFP+ cells that express Ki67 f. Five mice per group. Data are represented as mean \pm s.e.m. *P < 0.05 as determined by ANOVA. g Diagram showing isolation of Nestin-GFP+ (red) & Nestin-GFP- (purple) mesenchymal stem/progenitor cells (MSPCs) by fluorescence-activated cell sorting. Detailed information on isolation of MSPCs from femoral metaphysis from Nestin-GFP mice are described in Supplementary Fig. 3 & Methods section. h–j The sorted cells cultured, & SA- β Gal staining (h), BrdU incorporation (i), & p16INK4a immunostaining (j) performed, & representative images shown. k–m Quantification of % of cells that express SA- β Gal (k), BrdU (l), & p16INK4a (m). n = 5. Data are represented as mean \pm s.e.m. *P < 0.01 as determined by Student's t-tests Image collected & cropped by CiteAb from following publication (<https://pubmed.ncbi.nlm.nih.gov/29101351>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.





Immunocytochemistry/ Immunofluorescence: Ki67/MKI67 Antibody - BSA Free [NB500-170] - Cellular senescence occurs in primary spongiosa of long bone during late puberty. a–e Representative senescence-associated β -galactosidase (SA- β Gal) staining (blue) & quantitative analysis of SA- β Gal+ cells in femoral metaphysis (a–c) & diaphysis (d, e) sections from increasing ages of male mice. 4, 6, 8, & 12W represent 4-, 6-, 8-, & 12-week-old mice, respectively. Images in a are lower power with boxes outlining the area of higher power in b. Numbers of SA- β Gal+ cells per mm² tissue area in primary spongiosa (N. SA- β Gal+ cells/PS.Ar) (c) & diaphysis (N. SA- β Gal+ cells/DP.Ar) (e).

Counterstained with eosin (pink). f, g Representative images of immunofluorescence staining (f) & quantitative analysis of ki67+ (g) cells (red) in femoral primary spongiosa from 4, 6, 8, & 12-week-old male mice. DAPI stains nuclei blue. Images in upper panels in f are lower power with boxes outlining the area of higher power in lower panels. Five mice per group. Data are represented as mean \pm s.e.m. Ar tissue area, DP diaphysis, GP growth plate, N number, PS primary spongiosa. *P < 0.01 as determined by ANOVA Image collected & cropped by CiteAb from the following publication

(<https://pubmed.ncbi.nlm.nih.gov/29101351>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.

Ki67/MKI67 was detected in immersion fixed A431 human skin carcinoma cell line using Rabbit anti-Ki67/MKI67 Antigen Affinity Purified Polyclonal Antibody conjugated to Biotin (Catalog # NB500-170B) at 2 μ g/mL overnight at 4C. Cells were stained using Streptavidin conjugated to DyLight 550 (red) and counterstained with DAPI (blue). Cells were imaged using a 100X objective and digitally deconvolved.



Publications

Abdulla N, Aronson R, Plessis T et al. Protocol for the establishment and characterization of South African patient-derived intestinal organoids STAR protocols 2025-09-19 [PMID: 40705597]

Xie H, Lu L, Wang S et al. Deciphering the crucial role of IGF2BP3 in modulating stemness traits of salivary adenoid cystic carcinoma Medical oncology (Northwood, London, England) 2025-10-13 [PMID: 41082045]

Liu X, Yao Y, Yan J et al. RPS27L Enhances Myogenesis and Muscle Mass by Targeting IGF1 Through Liquid-Liquid Phase Separation. Advanced science (Weinheim, Baden-Wuerttemberg, Germany) 2025-08-31 [PMID: 40886325]

Li K, Yi Y, Ling R et al. PCIF1 drives oesophageal squamous cell carcinoma progression via m6Am-mediated suppression of MTF2 translation. Clinical and translational medicine 2025-03-29 [PMID: 40156159]

Gunasekara, N;Clauss, D;Voss, A;Schurz, K;Fleck, K;Neu-Gil, P;Bloch, W; The Influence of an Acute Endurance Intervention on Breast Cancer Cell Growth-A Pilot Study International journal of molecular sciences 2025-04-23 [PMID: 40362215]

Scalzone A, Cerqueni G, Wang XN et al. A cytokine-induced spheroid-based in vitro model for studying osteoarthritis pathogenesis Frontiers in Bioengineering and Biotechnology 2023-05-09 [PMID: 37229489]

Jia Y, Li HY, Wang J et al. Tripartite motif containing 69 elicits ERK2-dependent EYA4 turnover to impart pancreatic tumorigenesis Journal of Cancer 2023-01-01 [PMID: 36741265]

Xie XP, Laks DR, Sun D et al. Quiescent human glioblastoma cancer stem cells drive tumor initiation, expansion, and recurrence following chemotherapy Developmental Cell 2022-01-01 [PMID: 35016005]

Ye M, Liu Z, Gao C et al. Iturin A Rescued STb-R-Induced Pork Skeletal Muscle Growth Restriction through the Hypothalamic-Pituitary-mTORC1 Growth Axis Animals (Basel) 2022-06-17 [PMID: 35739903]

Galhenage P, Zhou Y, Perry E, Loc B et Al. Replication stress and defective checkpoints make fallopian tube epithelial cells putative drivers of high-grade serous ovarian cancer Cell Rep 2023-09-20 [PMID: 37729060]

Zhang Y, Liu G, Zeng Q, Wu W et Al. CCL19-producing fibroblasts promote tertiary lymphoid structure formation enhancing anti-tumor IgG response in colorectal cancer liver metastasis Cancer Cell 2024-08-13 [PMID: 39137726]

Xi G, Feng P, Zhang X et Al. iPSC-derived cells stimulate ABCG2(+)/NES(+) endogenous trabecular meshwork cell proliferation and tissue regeneration Cell Prolif 2024-02-14 [PMID: 38356373]

More publications at <http://www.novusbio.com/NB500-170>

Procedures

Immunocytochemistry/Immunofluorescence Protocol for Ki67/MKI67 Antibody (NB500-170)

Immunocytochemistry Protocol

Culture cells to appropriate density in 35 mm culture dishes or 6-well plates.

1. Remove culture medium and wash the cells briefly in PBS. Add 10% formalin to the dish and fix at room temperature for 10 minutes.
2. Remove the formalin and wash the cells in PBS.
3. Permeabilize the cells with 0.1% Triton X100 or other suitable detergent for 10 min.
4. Remove the permeabilization buffer and wash three times for 10 minutes each in PBS. Be sure to not let the specimen dry out.
5. To block nonspecific antibody binding, incubate in 10% normal goat serum from 1 hour to overnight at room temperature.
6. Add primary antibody at appropriate dilution and incubate overnight at 4C.
7. Remove primary antibody and replace with PBS. Wash three times for 10 minutes each.
8. Add secondary antibody at appropriate dilution. Incubate for 1 hour at room temperature.
9. Remove secondary antibody and replace with PBS. Wash three times for 10 minutes each.
10. Counter stain DNA with DAPI if required.

Immunohistochemistry-Paraffin Protocol for Ki67/MKI67 Antibody (NB500-170)

Immunohistochemistry-Paraffin Embedded Sections

Antigen Unmasking:

Bring slides to a boil in 10 mM sodium citrate buffer (pH 6.0) then maintain at a sub-boiling temperature for 10 minutes. Cool slides on bench-top for 30 minutes (keep slides in the sodium citrate buffer at all times).

Staining:

1. Wash sections in deionized water three times for 5 minutes each.
2. Wash sections in PBS for 5 minutes.
3. Block each section with 100-400 ul blocking solution (1% BSA in PBS) for 1 hour at room temperature.
4. Remove blocking solution and add 100-400 ul diluted primary antibody. Incubate overnight at 4 C.
5. Remove antibody solution and wash sections in wash buffer three times for 5 minutes each.
6. Add 100-400 ul HRP polymer conjugated secondary antibody. Incubate 30 minutes at room temperature.
7. Wash sections three times in wash buffer for 5 minutes each.
8. Add 100-400 ul DAB substrate to each section and monitor staining closely.
9. As soon as the sections develop, immerse slides in deionized water.
10. Counterstain sections in hematoxylin.
11. Wash sections in deionized water two times for 5 minutes each.
12. Dehydrate sections.
13. Mount coverslips.



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NB500-170

NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB500-170

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

