

Product Datasheet

Numb Antibody - BSA Free NB500-178

Unit Size: 0.1 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 7

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB500-178

Updated 9/9/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB500-178



NB500-178

Numb Antibody - BSA Free

Product Information	
Unit Size	0.1 ml
Concentration	2.0 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.05% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	Tris-Glycine and 0.15M NaCl


Product Description	
Description	Novus Biologicals Rabbit Numb Antibody - BSA Free (NB500-178) is a polyclonal antibody validated for use in WB, ICC/IF and Simple Western. Anti-Numb Antibody: Cited in 7 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	8650
Gene Symbol	NUMB
Species	Human, Mouse, Chicken
Reactivity Notes	Immunogen sequence has 90% identity to rat.
Specificity/Sensitivity	This is specific for all four isoforms of the NUMB protein.
Immunogen	A synthetic peptide made to a C-terminal region of mouse NUMB (between residues 600-653). [UniProt# Q9QZS3]

Product Application Details	
Applications	Western Blot, Simple Western, Immunocytochemistry/ Immunofluorescence
Recommended Dilutions	Western Blot 0.2-0.5 ug/ml, Simple Western 1:200, Immunocytochemistry/ Immunofluorescence 1:500
Application Notes	<p>In WB one may see any or all of the four isoforms. The molecular weight of human and mouse NUMB is 72 kDa (isoform 1), 66 kDa (isoform 2), 71 kDa (isoform 3), and 65 kDa (isoform 4). Multiple non-specific bands may be seen with lower dilutions and/or longer exposure times than suggested in the protocol we used to obtain our data.</p> <p>In Simple Western only 10 - 15 uL of the recommended dilution is used per data point.</p> <p>See Simple Western Antibody Database for Simple Western validation: Tested in A431 lysate 0.5 mg/mL, separated by Size, antibody dilution of 1:200, apparent MW was 83 kDa. Separated by Size-Wes, Sally Sue/Peggy Sue.</p>




Images

Western Blot: NUMB Antibody [NB500-178] - Detection of NUMB isoforms 1 and 2 in A431 whole cell lysate (20 ug) using 0.5 ug/ml of NB500-178. ECL detection: 30 seconds.




Immunocytochemistry/Immunofluorescence: Numb Antibody [NB500-178] - Rat FR cells were fixed in 4% paraformaldehyde for 10 minutes and permeabilized in 0.05% Triton X-100 in PBS for 5 minutes. The cells were incubated with (NB500-178) at 1ug/ml overnight at 4C and detected with an anti-rabbit DyLight 488 (Green) at a 1:1000 dilution for 60 minutes. Nuclei were counterstained with DAPI (Blue). Cells were imaged using a 40X objective.



Immunocytochemistry/Immunofluorescence: NUMB Antibody [NB500-178] - The NUMB antibody was tested in HepG2 cells at a 1:500 dilution against Dylight 488 (Green). Alpha-tubulin and nuclei were counterstained against Dylight 550 (Red) and DAPI (Blue), respectively.




Immunocytochemistry/Immunofluorescence: Numb Antibody [NB500-178] - HepG2 cells were fixed in 4% paraformaldehyde for 10 minutes and permeabilized in 0.05% Triton X-100 in PBS for 5 minutes. The cells were incubated with Numb Antibody (NB500-178) at 1ug/ml overnight at 4C and detected with an anti-rabbit DyLight 488 (Green) at a 1:1000 dilution for 60 minutes. Nuclei were counterstained with DAPI (Blue). Cells were imaged using a 40X objective.



Simple Western: Numb Antibody [NB500-178] - Simple Western lane view shows a specific band for NUMB in 0.5 mg/ml of A431 lysate. This experiment was performed under reducing conditions using the 12-230 kDa separation system.



Characterisation of LNX1 interacting proteins. (A) Schematic diagram of the domain structure of LNX1p80 and LNX2 showing the RING and four PDZ domains. N represents the NUMB-binding NPAY/NPAF motif. (B) The ability of the indicated proteins to interact with transfected GFP-tagged LNX constructs was assessed in HEK 293 cells. For each interacting protein, top panels show western blots of cell lysates (Lys), while the bottom panels show the output of a GFP "pull down" assay (PD). In the panels on the left, the specificity of interactions for LNX1 versus LNX2 was assessed, while on the right the interaction site on LNX1 was mapped to individual protein domains. Binding of endogenous proteins to LNX was assessed for liprin α -1, KIF7 and NUMB. For the other proteins, interactions of transfected HA or GST epitope-tagged proteins were assessed. For AKAP13, the mapping to LNX domains was performed in two separate experiments. Successful expression and pull down of GFP-tagged LNX proteins was verified in all assays and representative "pull down" blots probed for GFP are shown. n = 2–3. Image collected and cropped by CiteAb from the following open publication (<https://pubmed.ncbi.nlm.nih.gov/29121065>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Publications

Das A, Adhikary S, Chowdhury AR, Barui A Chirality-induced Lineage Enforcement of Mechanosensitive Mesenchymal Stem Cells Across Germ Layer Boundaries Stem cell reviews and reports 2023-11-16 [PMID: 37971671]

Das A, Adhikary S, Chowdhury AR Et al. Leveraging Substrate Stiffness to Promote Stem Cell Asymmetric Division via Mechanotransduction-Polarity Protein Axis and Its Bayesian Regression Analysis Rejuvenation Res 2022-03-22 [PMID: 35316074] (ICC/IF)

Details:

Citation using the Texas Red version of this antibody.

Huang SC, Vu LV, Yu FH Et al. Multifunctional protein 4.1R regulates the asymmetric segregation of Numb during terminal erythroid maturation The Journal of biological chemistry 2021-08-06 [PMID: 34364872]

Sikorski K, Mehta A et al. A high-throughput pipeline for validation of antibodies. Nat Methods 2018-01-11 [PMID: 30377371] (Human)

Details:

Antibody validation based on denaturing gel electrophoresis of biotinylated cell lysates (PAGE) followed by mass spectrometry (MS) and antibody array analysis (MAP).

Lenihan JA, Saha O, Young PW et al. Proteomic analysis reveals novel ligands and substrates for LNX1 E3 ubiquitin ligase PLoS One. 2017-11-08 [PMID: 29121065] (WB, Human)

Wang H, Xiang D, Liu B et al. Inadequate DNA Damage Repair Promotes Mammary Transdifferentiation, Leading to BRCA1 Breast Cancer Cell 2019-06-27 [PMID: 31251913] (Mouse)

Lenihan JA, Saha O, Heimer-McGinn V et al. Decreased Anxiety-Related Behaviour but Apparently Unperturbed NUMB Function in Ligand of NUMB Protein-X (LNX) 1/2 Double Knockout Mice. Mol. Neurobiol. 2016-11-26 [PMID: 27889896] (WB, Mouse)



Procedures

Western Blot protocol for Numb Antibody (NB500-178)

Numb Antibody:

Western Blot Protocol

1. Perform SDS-PAGE (4-12%) on samples to be analyzed, loading 20ug of total protein per lane.
2. Transfer proteins to Nitrocellulose according to the instructions provided by the manufacturer of the transfer apparatus.
3. Stain the blot using ponceau S for 1-2 minutes to access the transfer of proteins onto the nitrocellulose membrane. Rinse the blot in water to remove excess stain and mark the lane locations and locations of molecular weight markers using a pencil.
4. Rinse the blot in TBS for approximately 5 minutes.
5. Block the membrane using 5% non-fat dry milk in TBS for 1 hour.
6. Dilute the rabbit anti-Numb primary antibody (NB 500-178) in blocking buffer and incubate 2 hours at room temperature.
7. Wash the membrane in water for 5 minutes and apply the diluted rabbit-IgG HRP-conjugated secondary antibody in blocking buffer (as per manufacturer's instructions) and incubate 1 hour at room temperature.
8. Wash the blot in TBS containing 0.05-0.1% Tween-20 for 10-20 minutes.
9. Wash the blot in type I water for an additional 10-20 minutes (this step can be repeated as required to reduce background).
10. Apply the detection reagent of choice in accordance with the manufacturer's instructions (Amersham's ECL is the standard reagent used at Novus Biologicals).

Note: Tween-20 can be added to the blocking buffer at a final concentration of 0.05-0.2%, provided it does not interfere with antibody-antigen binding.





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NB500-178

NB820-59461	A-431 Whole Cell Lysate
NB500-178PEP	Numb Antibody Blocking Peptide
NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB500-178

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications



