

Product Datasheet

Caspase-7 Antibody (Mch3 1-1-11) - BSA Free NB500-206

Unit Size: 0.1 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 2

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB500-206

Updated 9/9/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB500-206



NB500-206

Caspase-7 Antibody (Mch3 1-1-11) - BSA Free

Product Information	
Unit Size	0.1 ml
Concentration	This product is unpurified. The exact concentration of antibody is not quantifiable.
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	Mch3 1-1-11
Preservative	0.1% Sodium Azide
Isotype	IgG1 Kappa
Purity	Unpurified
Buffer	Ascites
Target Molecular Weight	35 kDa


Product Description	
Description	Novus Biologicals Mouse Caspase-7 Antibody (Mch3 1-1-11) - BSA Free (NB500-206) is a monoclonal antibody validated for use in WB and Simple Western. Anti-Caspase-7 Antibody: Cited in 2 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Mouse
Gene ID	840
Gene Symbol	CASP7
Species	Human, Mouse, Rat
Immunogen	Full-length recombinant human Caspase 7 [UniProt# P55210]

Product Application Details	
Applications	Western Blot, Simple Western, Knockdown Validated
Recommended Dilutions	Western Blot 1:500-1:1000, Simple Western 1:500, Knockdown Validated
Application Notes	<p>This Caspase 7 (Mch3 1-1-11) antibody is useful for Western blot, where a band can be seen at approx. 35 kDa.</p> <p>In Simple Western only 10 - 15 uL of the recommended dilution is used per data point.</p> <p>See Simple Western Antibody Database for Simple Western validation: Tested in Jurkat +Etoposide, HeLa lysate 0.5 mg/mL, separated by Size, antibody dilution of 1:50, 1:500, apparent MW was 41, 44 kDa. Separated by Size-Wes, Sally Sue/Peggy Sue.</p> <p>The observed molecular weight of the protein may vary from the listed predicted molecular weight due to post translational modifications, post translation cleavages, relative charges, and other experimental factors.</p>




Images


Western Blot: Caspase-7 Antibody (Mch3 1-1-11) [NB500-206] - Western blot shows lysates of A549 human lung epithelial parental cell line and Caspase-7 knockout (KO) A549 cell line. PVDF membrane was probed with 1:500 of Mouse Anti-Human Caspase-7 Monoclonal Antibody (Catalog # NB500-106) followed by HRP-conjugated Anti-Mouse IgG Secondary Antibody (Catalog #HAF018). Specific band was detected for Caspase-7 at approximately 40 kDa (as indicated) in the parental A549 cell line, but is not detectable in the knockout A549 cell line. This experiment was conducted under reducing conditions.



Western Blot: Caspase-7 Antibody (Mch3 1-1-11) [NB500-206] - HeLa whole cell lysate.



Simple Western: Caspase-7 Antibody (Mch3 1-1-11) [NB500-206] - Simple Western lane view shows a specific band for Caspase 7 in 0.5 mg/ml of HeLa lysate. This experiment was performed under reducing A246 conditions using the 12-230 kDa separation system.



Publications

Badr DM, Hafez HF, Agha AM, Shouman SA. The Combination of alpha-Tocopheryl Succinate and Sodium Selenite on Breast Cancer: A Merit or a Demerit. *Oxid Med Cell Longev*. 2016-04-29 [PMID: 27127548] (WB, Human)

Blanc-Brude OP et al. Therapeutic targeting of the survivin pathway in cancer: initiation of mitochondrial apoptosis and suppression of tumor-associated angiogenesis. *Clin Cancer Res*;9(7):2683-92. 2003-07-01 [PMID: 12855648]

Procedures

Protocol specific for Caspase 7 Antibody (NB500-206)

Caspase-7 Antibody (Mch3 1-1-11):

Procedure Guide for NB 500-206 Monoclonal anti-Caspase 7

Western Protocol

- 1) Prepare samples for a NuPAGE gel and heat them at 70 degree Celsius for 10 minutes.
- 2) Run a 4-12% NuPAGE Bis-Tris gel, loading 50-100 mg of whole cell extract per lane, for 45 minutes at 200 Volts.
- 3) Soak transfer sponges and gel 15 minutes and nitrocellulose membrane 5 minutes prior to transfer.
- 4) Transfer the proteins to the nitrocellulose membrane for 1 hour at 30 Volts, in 4 degrees C transfer buffer.
- 5) Block the membrane in TBST + 5% non-fat dry milk, for 1 hour at room temperature (RT), shaking gently.
- 6) Rinse the membrane twice with TBST.
- 7) Incubate the membrane in anti-Caspase 7 [NB 500-206], diluted 1:500-1:1,000 in TBS + 1% BSA, for 90 minutes at RT.
- 8) Wash the membrane once for 15 minutes, then 3 times 5 minutes, in TBST at RT.
- 9) Incubate the membrane in anti-mouse conjugated to HRP (secondary antibody), diluted in TBS + 1% BSA for 35 minutes at RT.
- 10) Wash the membrane once for 15 minutes, then 3 times 5 minutes, in TBST at RT.
- 11) Develop membrane in chemiluminescent reagents, as instructed by kit-vendor.

*Positive control(s): HeLa whole cell extracts (cat# NB 800-PC1)

Jurkat whole cell extracts (cat# NB 800-PC2)





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NB500-206

NB800-PC1	HeLa Whole Cell Lysate
NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB7539	Goat anti-Mouse IgG (H+L) Secondary Antibody [HRP]
NBP1-43319-0.5mg	Mouse IgG1 Kappa Isotype Control (P3.6.2.8.1)

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB500-206

Earn gift cards/discounts by submitting a publication using this product:

www.novusbio.com/publications

