

Product Datasheet

beta-Actin Antibody - BSA Free NB600-503

Unit Size: 0.05 ml

Store at 4C. Do not freeze.

www.novusbio.com



technical@novusbio.com

Reviews: 4 Publications: 134

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB600-503

Updated 9/9/2025 v.20.1

**Earn rewards for product
reviews and publications.**

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB600-503



NB600-503

beta-Actin Antibody - BSA Free

Product Information	
Unit Size	0.05 ml
Concentration	1.0 mg/ml
Storage	Store at 4C. Do not freeze.
Clonality	Polyclonal
Preservative	0.09% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	Tris-Citrate/Phosphate (pH 7.0 - 8.0)
Target Molecular Weight	42 kDa


Product Description	
Description	Novus Biologicals Rabbit beta-Actin Antibody - BSA Free (NB600-503) is a polyclonal antibody validated for use in IHC, WB, ELISA, Flow, ICC/IF, Simple Western and IP. Anti-beta-Actin Antibody: Cited in 132 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	60
Gene Symbol	ACTB
Species	Human, Mouse, Rat, Porcine, Avian, Bovine, Chinese Hamster, Fish, Primate, Rabbit
Reactivity Notes	Rabbit reactivity and Fish reactivity reported in scientific literature (PMID: 23813946 and 25842206 respectively). Expected to cross-react with a wide range of species due to sequence identity. Bovine reactivity reported in scientific literature (PMID:33066332).
Immunogen	This beta-Actin Antibody was made from a synthetic peptide made to an N-terminal region of human Beta Actin. [UniProt P60709]

Product Application Details	
Applications	Western Blot, Simple Western, Immunohistochemistry-Paraffin, ELISA, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunoprecipitation, Proximity Ligation Assay, Block/Neutralize
Recommended Dilutions	Western Blot 1:100-1:2000, Simple Western 1:12.5, Flow Cytometry 1:10-1:1000, ELISA, Immunohistochemistry 1:100, Immunocytochemistry/ Immunofluorescence, Immunoprecipitation 1:10-1:500, Immunohistochemistry-Paraffin 1:100, Proximity Ligation Assay, Block/Neutralize
Application Notes	See Simple Western Antibody Database for Simple Western validation: tested in HeLa lysate (0.1 mg/ml); antibody dilution of 1:12.5; separated by size; detects a band at 50 kDa




Images


Immunocytochemistry/Immunofluorescence: beta-Actin Antibody [NB600-503] - Cultured pig trabecular meshwork cells stained with beta-Actin antibody at a dilution of 1:500 followed by an anti-rabbit Alexa Fluor488 at 1:1000 dilution (green). Image from verified customer review.




Western Blot: beta-Actin Antibody [NB600-503] - Rabbit polyclonal at 1/5000.



Simple Western: beta-Actin Antibody [NB600-503] - Simple Western lane view shows a specific band for Beta Actin in 0.1 mg/ml of HeLa lysate. This experiment was performed under reducing conditions using the 12-230 kDa separation system.




Flow Cytometry: beta-Actin Antibody [NB600-503] - An intracellular stain was performed on HeLa cells with beta-actin Antibody NB600-503AF488 (blue) and a matched isotype control (orange). Cells were fixed with 4% PFA and then permeabilized with 0.1% saponin. Cells were incubated in an antibody dilution of 5 ug/mL for 30 minutes at room temperature. Both antibodies were conjugated to Alexa Fluor 488.




Copyright © 2018 Novus Biologicals


Western Blot: beta-Actin Antibody [NB600-503] - Western blot analysis of Jurkat, COS, and CHO cell lysate using beta actin antibody at 1:100.




Immunohistochemistry-Paraffin: beta-Actin Antibody [NB600-503] - Analysis of Beta Actin in mouse epidermis using DAB with hematoxylin counterstain.




Immunocytochemistry/Immunofluorescence: beta-Actin Antibody [NB600-503] - IF Confocal analysis of HeLa cells using Beta Actin antibody (NB600-503, 1:5). An Alexa Fluor 488-conjugated Goat to rabbit IgG was used as secondary antibody (green). DAPI was used to stain the cell nuclei (blue).




Flow Cytometry: beta-Actin Antibody [NB600-503] - Analysis of HeLa cells using mouse Monoclonal beta-Actin antibody (Orange) and Isotype control Antibody (Blue).




Simple Western: beta-Actin Antibody [NB600-503] - Early start GM1 administration did not affect alpha-synuclein expression or transport to the striatum. When assessed 1 week following AAV-A53T alpha-synuclein injection, levels of striatal alpha-synuclein were no different in saline (N=6) vs. GM1-treated animals (N=6), suggesting no influence of GM1 on A53T alpha-synuclein transduction or transport to the striatum. Representative Wes Simple Western blots are shown after cropping (full length images of blots are presented as Supplementary Fig. 2. Image collected and cropped by CiteAb from the following publication (<https://www.nature.com/articles/s41598-019-42847-x>), licensed under a CC-BY license.




Western Blot: beta-Actin Antibody [NB600-503] - Regulation of inflammatory signaling in hSMC.ND (open bars) & T2D (solid bars) cells extracted after 48 hr treatment with LPS, Pioglitazone (Pio), palmitate or oleate. Results expressed relative to untreated control for each individual set of cells, Ave + SEM. (A) IκBα protein, n = 7–12 & 6–15 for ND & T2D, respectively. (B) Total & phospho-p38, n = 6–14 & 5–8. (C) Total & phospho-p44/42, n = 7–13 & 7–11. (D) Total & phospho-JNK, n = 9–12 & 3–8. * p<0.05 T2D vs ND. †p<0.05 T2D response vs ND response Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/27453994>), licensed under a CC0-1.0 license. Not internally tested by Novus Biologicals.




Western Blot: beta-Actin Antibody [NB600-503] - Impact of T2D on inflammatory signaling in hSMC. Proteins extracted from ND (open bars) & T2D (solid bars) hSMC under control conditions. (A) Representative western blots. (B) Quantification of protein expression normalized to loading control (β-actin). Ave + SEM, n = 13–21 for ND, 10–16 for T2D. (C) Protein phosphorylation, expressed as ratio of phosphorylated to total protein. Ave + SEM, n = 13–17 for ND, 9–15 for T2D. *p<0.05 vs ND Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/27453994>), licensed under a CC0-1.0 license. Not internally tested by Novus Biologicals.



Western Blot: beta-Actin Antibody [NB600-503] - Early start GM1 administration did not affect α-synuclein expression or transport to the striatum. (A,B) When assessed 1 week following AAV-A53T α-synuclein injection, levels of striatal α-synuclein were no different in saline (N = 6) vs. GM1-treated animals (N = 6), suggesting no influence of GM1 on A53T α-synuclein transduction or transport to the striatum. Representative Wes Western blots are shown after cropping (full length images of blots are presented as Supplementary Fig. 2. (C) Double label immunofluorescence 1 week after AAV-A53T α-synuclein injection showed no differences between saline & GM1-treated animals in α-synuclein accumulation (green) in TH+ neurons (red) in the SNc. Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/31182727>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Western Blot: beta-Actin Antibody [NB600-503] - Anti-hnRNP A1 antibodies alter protein levels as measured by Western blot. SK-N-SH cells were cultured & treated with anti-hnRNP A1 antibodies or control IgG. Following a 48 hour incubation, cells were lysed & protein lysate was run on 10% Tris-glycine gels for Western blot analysis & probed for Beta-actin (A) or GAPDH (B)) (control), hnRNP A1, SPG4, SPG7, & SPG20. Results revealed that there was a marked reduction of SPG 4 & SPG 7 protein levels in anti-hnRNP A1 antibody compared to control isotype IgG treated cells. There was a variable response to hnRNP A1 protein & a modest reduction of SPG 20. Parentheses show relative percent reduction of signal comparing anti-hnRNP A1 antibody to control isotype IgG treatment of cells. (L=lysate, IgG=control isotype IgG, A1=anti-hnRNP A1-M9 antibody treatment of cells). Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/27375925>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Publications

E Aladowicz, L Granieri, F Marocchi, S Punzi, G Giardina, PF Ferrucci, G Mazzarol, M Capra, G Viale, S Confalonie, S Gandini, F Lotti, L Lanfrancon ShcD Binds DOCK4, Promotes Ameboid Motility and Metastasis Dissemination, Predicting Poor Prognosis in Melanoma Cancers (Basel), 2020-11-13;12(11):. 2020-11-13 [PMID: 33202906] (Western Blot, Human, Mouse, Hamster - Cricetulus (Chinese Hamster))

Shastri S, Shinde T, Perera AP et al. Idebenone Protects against Spontaneous Chronic Murine Colitis by Alleviating Endoplasmic Reticulum Stress and Inflammatory Response Biomedicines 2020-09-28 [PMID: 32998266] (Western Blot, Human, Mouse, Hamster - Cricetulus (Chinese Hamster))

Krysa SJ, Allen LH. Metabolic Reprogramming Mediates Delayed Apoptosis of Human Neutrophils Infected With Francisella tularensis Frontiers in Immunology 2022-05-25 [PMID: 35693822] (Western Blot, Human, Mouse, Hamster - Cricetulus (Chinese Hamster))

Shastri S, Shinde T, Woolley KL et al. Short-Chain Naphthoquinone Protects Against Both Acute and Spontaneous Chronic Murine Colitis by Alleviating Inflammatory Responses Frontiers in Pharmacology 2021-08-23 [PMID: 34497514] (Western Blot, Human, Mouse, Hamster - Cricetulus (Chinese Hamster))

Miroshnikova VV, Panteleeva AA, Bazhenova EA et al. [Regulation of ABCA1 and ABCG1 gene expression in the intraabdominal adipose tissue] Biomeditsinskaya Khimiya 2016-07-16 [PMID: 27420620] (Western Blot, Human, Mouse, Hamster - Cricetulus (Chinese Hamster))

Anderson T, Sharma S, Kelberman MA, Ware C et Al. Obesity during preclinical Alzheimer's disease development exacerbates brain metabolic decline J Neurochem 2023-06-30 [PMID: 37391269]

Davis EA, Wald HS, Suarez AN, Zubcevic J et Al. Ghrelin Signaling Affects Feeding Behavior, Metabolism, and Memory through the Vagus Nerve Curr Biol 2020-09-18 [PMID: 32946754]

Zhou Y, Li ZL, Ding L, Zhang XJ et Al. Long noncoding RNA SNHG5 promotes podocyte injury via the microRNA-26a-5p/TRPC6 pathway in diabetic nephropathy J Biol Chem 2022-10-18 [PMID: 36257404]

Li J, Jeong SY, Xiong B et Al. Repurposing pyridoxamine for therapeutic intervention of intravascular cell-cell interactions in mouse models of sickle cell disease Haematologica 2019-10-31 [PMID: 33054081]

Bostanci A, Doganlar O., et Al. MELATONIN ENHANCES TEMOZOLOMIDE-INDUCED APOPTOSIS IN GLIOBLASTOMA AND NEUROBLASTOMA CELLS Exp Oncol 2024-10-13 [PMID: 39396175]

Md. Aejazur Rahman, Bridgette M. Cumming, Kelvin W. Addicott, Hayden T. Pacl, Shannon L. Russell, Kievershen Nargan, Threnesan Naidoo, Pratistadevi K. Ramdial, John H. Adamson, Rui Wang, Adrie J. C. Steyn Hydrogen sulfide dysregulates the immune response by suppressing central carbon metabolism to promote tuberculosis Proceedings of the National Academy of Sciences of the United States of America 2020-03-24 [PMID: 32139610]

Mueller, J;van Muilekom, DR;Ehlers, J;Suhr, M;Hornburg, SC;Bang, C;Wilkes, M;Schultheiß, T;Maser, E;Rebl, A;Goldammer, T;Seibel, H;Schulz, C; Dietary Chlorella vulgaris supplementation modulates health, microbiota and the response to oxidative stress of Atlantic salmon Scientific reports 2024-10-10 [PMID: 39389986]

More publications at <http://www.novusbio.com/NB600-503>



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NB600-503

NB800-PC1	HeLa Whole Cell Lysate
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB600-503

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

