

Product Datasheet

CRHR1/CRF1 Antibody NBP1-00175

Unit Size: 0.1 mg

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 6

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP1-00175

Updated 9/9/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP1-00175



NBP1-00175**CRHR1/CRF1 Antibody**

Product Information	
Unit Size	0.1 mg
Concentration	0.5 mg/ml
Storage	Store at -20C. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.02% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	Tris saline (20 mM Tris pH 7.3, 150 mM NaCl), 0.5% BSA


Product Description	
Description	Novus Biologicals Goat CRHR1/CRF1 Antibody (NBP1-00175) is a polyclonal antibody validated for use in IHC, WB, ELISA, ICC/IF and Simple Western. Anti-CRHR1/CRF1 Antibody: Cited in 6 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Goat
Gene ID	1394
Gene Symbol	CRHR1
Species	Human, Mouse, Rat
Immunogen	Peptide with sequence C-NEEKKSKVHYH corresponding to internal region according to NP_004373.2.

Product Application Details	
Applications	Western Blot, Simple Western, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Peptide ELISA
Recommended Dilutions	Western Blot 0.1 - 0.3 ug/mL, Simple Western 1:100, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 10 ug/mL, Peptide ELISA Detection limit 1:128000
Application Notes	Western blot: Approx 60kDa band observed in Human Brain (Cerebellum), Colon, Rat Brain and Mouse spinal cord lysates (calculated MW of 47.7kDa according to NP_004373.2). Primary incubation was 1 hour. IHC reported in scientific literature (PMID: 22570383). Use in ICC/IF reported in scientific literature (PMID: 30439403). See Simple Western Antibody Database for Simple Western validation: Tested in Brain, separated by Size, antibody dilution of 1:100




Images


Western Blot: CRHR1/CRF1 Antibody [NBP1-00175] - Staining of Human Colon (A), Cerebellum (B), Rat Brain (C) and Mouse spinal Cord (D) lysates (35 ug protein in RIPA buffer). Antibody at 1 ug/mL. Detected by chemiluminescence.



Immunocytochemistry/Immunofluorescence: CRHR1/CRF1 Antibody [NBP1-00175] - Immunofluorescence analysis of paraformaldehyde fixed MCF7 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10 ug/mL) followed by Alexa Fluor 488 secondary antibody (2 ug/mL), showing cytoplasmic and vesicle staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10 ug/mL) followed by Alexa Fluor 488 secondary antibody (2 ug/mL).



Immunocytochemistry/Immunofluorescence: CRHR1/CRF1 Antibody [NBP1-00175] - Immunofluorescence analysis of paraformaldehyde fixed Neuro2a cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10 ug/mL) followed by Alexa Fluor 488 secondary antibody (2 ug/mL), showing plasma membrane, and vesicle staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10 ug/mL) followed by Alexa Fluor 488 secondary antibody (2 ug/mL).



Publications

Ritchie JL, Walters JL, Galliou JMC Et al. Basolateral amygdala corticotropin-releasing factor receptor type 1 regulates context-cocaine memory strength during reconsolidation in a sex-dependent manner *Neuropharmacology* 2021-12-01 [PMID: 34610289] (Simple Western, Rat)

Wang S, Wang C, Yu Z et al. Agarwood Essential Oil Ameliorates Restrain Stress-Induced Anxiety and Depression by Inhibiting HPA Axis Hyperactivity. *Int J Mol Sci.* 2018-11-05 [PMID: 30400578] (WB, Mouse)

D'Costa S, Ayyadurai S, Gibson AJ et al. Mast Cell CRF2 Suppresses Mast Cell Degranulation and Limits the Severity of Anaphylaxis and Stress-Induced Intestinal Permeability. *J. Allergy Clin. Immunol.* 2018-11-12 [PMID: 30439403] (ICC/IF, Mouse)

Rissman RA, Lee KF, Vale W, Sawchenko PE. Corticotropin-releasing factor receptors differentially regulate stress-induced tau phosphorylation. *J Neurosci* 2007-06-13 [PMID: 17567816]

Xu G, Rabadan-Diehl C, Nikodemova M et al. Inhibition of corticotropin releasing hormone type-1 receptor translation by an upstream AUG triplet in the 5' untranslated region. *Mol Pharmacol* 2001-03-01 [PMID: 11179443]

Jungling K, Liu X, Lesting J et al. Activation of neuropeptide S expressing neurons in the locus coeruleus by corticotropin-releasing factor. *J Physiol* 2012-05-08 [PMID: 22570383] (IHC-P, Mouse)





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP1-00175

NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF017	Rabbit anti-Goat IgG Secondary Antibody [HRP (Horseradish Peroxidase)]
HAF109	Donkey anti-Goat IgG Secondary Antibody [HRP (Horseradish Peroxidase)]
NB410-28088-1mg	Goat IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP1-00175

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications



