

Product Datasheet

FANCG Antibody NBP1-06035

Unit Size: 0.1 mg

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 2

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP1-06035

Updated 9/9/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP1-06035



NBP1-06035

FANCG Antibody


Product Information	
Unit Size	0.1 mg
Concentration	0.5 mg/ml
Storage	Store at -20C. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.02% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	Tris saline (20 mM Tris pH 7.3, 150 mM NaCl), 0.5% BSA

Product Description	
Description	Novus Biologicals Goat FANCG Antibody (NBP1-06035) is a polyclonal antibody validated for use in IHC, WB and ELISA. Anti-FANCG Antibody: Cited in 2 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Goat
Gene ID	2189
Gene Symbol	FANCG
Species	Human
Immunogen	Peptide with sequence LEEFRTSLPKSCDL corresponding to C-Terminus according to NP_004620.1.


Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, Immunohistochemistry, Peptide ELISA
Recommended Dilutions	Western Blot 0.5 - 1 ug/ml, Immunohistochemistry, Immunohistochemistry-Paraffin 3 ug/ml, Peptide ELISA Detection limit 1:32000
Application Notes	WB: Approx. 70 kDa band observed in HeLa lysates (calculated MW of 68.6 kDa band according to NP_004620.1). In transfected HEK293 transiently expressing FANCG a band of approx. 65 kDa is observed. This band is not observed in the non-transfected HEK293. IHC: Paraffin embedded Human Uterus and Spleen.

Images

Western Blot: FANCG Antibody [NBP1-06035] - Staining of Jurkat nuclear lysate (35ug protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.




Immunohistochemistry-Paraffin: FANCG Antibody [NBP1-06035] - staining of paraffin embedded Human Uterus. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.




Western Blot: FANCG Antibody [NBP1-06035] - Analysis of FANCG using this antibody at 0.5 ug/ml in HeLa cell lysate (35ug protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

250kDa
150kDa
100kDa
75kDa
50kDa
37kDa
25kDa
20kDa
15kDa

Western Blot: FANCG Antibody [NBP1-06035] - HEK293 overexpressing FANCG and probed with (mock transfection in first lane).



Immunohistochemistry-Paraffin: FANCG Antibody [NBP1-06035] - staining of paraffin embedded Human Spleen. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



Publications

Liu N, Lamerdin JE, Tucker JD et al. The human XRCC9 gene corrects chromosomal instability and mutagen sensitivities in CHO UV40 cells. Proc Natl Acad Sci U S A 1997-08-19 [PMID: 9256465]

Tomida J, Itaya A, Shigechi T et al. A novel interplay between the Fanconi anemia core complex and ATR-ATRIP kinase during DNA cross-link repair. Nucleic Acids Res 2013-05-30 [PMID: 23723247] (WB, Human)





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP1-06035

NBL1-10589	FANCG Overexpression Lysate
NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF017	Rabbit anti-Goat IgG Secondary Antibody [HRP (Horseradish Peroxidase)]
HAF109	Donkey anti-Goat IgG Secondary Antibody [HRP (Horseradish Peroxidase)]
NB410-28088-1mg	Goat IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP1-06035

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications



