

# Product Datasheet

## EIF3A Antibody - BSA Free

### NBP1-18891

Unit Size: 100 ul

Store at 4C. Do not freeze.

[www.novusbio.com](http://www.novusbio.com)



[technical@novusbio.com](mailto:technical@novusbio.com)

**Reviews: 1** **Publications: 11**

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:  
[www.novusbio.com/NBP1-18891](http://www.novusbio.com/NBP1-18891)

Updated 9/9/2025 v.20.1

Earn rewards for product  
reviews and publications.


Submit a publication at [www.novusbio.com/publications](http://www.novusbio.com/publications)

Submit a review at [www.novusbio.com/reviews/destination/NBP1-18891](http://www.novusbio.com/reviews/destination/NBP1-18891)






TRS functions similarly to eIF4G & acts as an eIF4F analog. a Pull-down assay of co-expressed TRS-Strep with eIF4A- or eIF4G-FLAG in 293 T cells. TRS-Strep was pulled down with Strep-Tactin beads, & co-precipitation of eIF4A or 4 G was determined by immunoblotting with anti-FLAG antibody. EV, empty vector. \* indicates a nonspecific band. b Immunoassay of the co-expression of different combinations of plasmids in 293T cells. Myc-TRS was immunoprecipitated with anti-Myc antibody, & co-precipitation of other proteins was determined using tag-specific antibodies. c Immunoassay of co-expressed eIF4A-FLAG with GST-fused full-length TRS or its various domains in 293T cells. eIF4A-FLAG was immunoprecipitated with anti-FLAG antibody, & co-precipitated TRS proteins were determined by immunoblotting with anti-GST antibody. d Pull-down assay of endogenous translation initiation factors with m7GTP-Sepharose beads in 293 T cells transfected with siRNAs against TRS, 4EHP, or a non-targeting control (siCont). Cap-bound proteins were eluted from beads & immunoblotted with the indicated antibodies. Sepharose beads were used as a negative control. e Pull-down assay of endogenous translation initiation factors with m7GTP-Sepharose beads in 293T cells transfected with siRNAs against eIF4G, eIF4E, or siCont, & their suppression effects on cap-binding of other components, were determined as in (d). The data are representative of at least three experiments, each with similar results Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/30902983>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



TRS functions similarly to eIF4G & acts as an eIF4F analog. a Pull-down assay of co-expressed TRS-Strep with eIF4A- or eIF4G-FLAG in 293 T cells. TRS-Strep was pulled down with Strep-Tactin beads, & co-precipitation of eIF4A or 4 G was determined by immunoblotting with anti-FLAG antibody. EV, empty vector. \* indicates a nonspecific band. b Immunoassay of the co-expression of different combinations of plasmids in 293T cells. Myc-TRS was immunoprecipitated with anti-Myc antibody, & co-precipitation of other proteins was determined using tag-specific antibodies. c Immunoassay of co-expressed eIF4A-FLAG with GST-fused full-length TRS or its various domains in 293T cells. eIF4A-FLAG was immunoprecipitated with anti-FLAG antibody, & co-precipitated TRS proteins were determined by immunoblotting with anti-GST antibody. d Pull-down assay of endogenous translation initiation factors with m7GTP-Sepharose beads in 293 T cells transfected with siRNAs against TRS, 4EHP, or a non-targeting control (siCont). Cap-bound proteins were eluted from beads & immunoblotted with the indicated antibodies. Sepharose beads were used as a negative control. e Pull-down assay of endogenous translation initiation factors with m7GTP-Sepharose beads in 293T cells transfected with siRNAs against eIF4G, eIF4E, or siCont, & their suppression effects on cap-binding of other components, were determined as in (d). The data are representative of at least three experiments, each with similar results Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/30902983>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



## Publications

Duan H, Zhang S, Zarai Y et al. eIF3 mRNA selectivity profiling reveals eIF3k as a cancer-relevant regulator of ribosome content *The EMBO Journal* 2023-06-15 [PMID: 37155573] (Western Blot, Human)

Hassan Hayek, Lauriane Gross, Aurélie Janvier, Laure Schaeffer, Franck Martin, Gilbert Eriani, Christine Allmang eIF3 interacts with histone H4 messenger RNA to regulate its translation *The Journal of Biological Chemistry* 2021-03-23 [PMID: 33766559]

Kejiou NS, Ilan L, Aigner S et al. Pyruvate Kinase M (PKM) binds ribosomes in a poly-ADP ribosylation dependent manner to induce translational stalling *Nucleic acids research* 2023-05-24 [PMID: 37224531] (Western Blot, Human)

Zhao Y, Zhao H, Zhang D et al. YTHDF3 Facilitates eIF2AK2 and eIF3A Recruitment on mRNAs to Regulate Translational Processes in Oxaliplatin-Resistant Colorectal Cancer *ACS chemical biology* 2022-06-15 [PMID: 35708211]

Gong C, Krupka JA, Gao J Et al. Sequential inverse dysregulation of the RNA helicases DDX3X and DDX3Y facilitates MYC-driven lymphomagenesis *Molecular cell* 2021-08-18 [PMID: 34437837] (WB, Human)

Lin Y, Li F, Huang L et al. eIF3 Associates with 80S Ribosomes to Promote Translation Elongation, Mitochondrial Homeostasis, and Muscle Health *Mol. Cell* 2020-08-20 [PMID: 32589965] (WB)

Lin Y, Li F, Huang L et al. eIF3 promotes early translation elongation to ensure mitochondrial homeostasis and skeletal muscle health *bioRxiv* 1905-07-11 (WB, Human)

Jeong SJ, Park S, Nguyen LT et al. A threonyl-tRNA synthetase-mediated translation initiation machinery *Nat Commun* 2019-03-22 [PMID: 30902983] (WB, Human)

Toompuu M, Karblane K, Pata P et al. ABCE1 is essential for S phase progression in human cells. *Cell Cycle*. 2016-03-17 [PMID: 26985706] (WB, Human)

Lee AS, Kranzusch PJ, Cate JH. eIF3 targets cell-proliferation messenger RNAs for translational activation or repression *Nature* 2015-04-06 [PMID: 25849773] (WB, Human)

Chazal PE, Dagueneat E, Wendling C et al. EJC core component MLN51 interacts with eIF3 and activates translation *Proc Natl Acad Sci U S A* 2013-03-25 [PMID: 23530232] (WB, Human)





### **Novus Biologicals USA**

10730 E. Briarwood Avenue  
Centennial, CO 80112  
USA  
Phone: 303.730.1950  
Toll Free: 1.888.506.6887  
Fax: 303.730.1966  
nb-customerservice@bio-techne.com

### **Bio-Techne Canada**

21 Canmotor Ave  
Toronto, ON M8Z 4E6  
Canada  
Phone: 905.827.6400  
Toll Free: 855.668.8722  
Fax: 905.827.6402  
canada.inquires@bio-techne.com

### **Bio-Techne Ltd**

19 Barton Lane  
Abingdon Science Park  
Abingdon, OX14 3NB, United Kingdom  
Phone: (44) (0) 1235 529449  
Free Phone: 0800 37 34 15  
Fax: (44) (0) 1235 533420  
info.EMEA@bio-techne.com

### **General Contact Information**

www.novusbio.com  
Technical Support: nb-technical@bio-techne.com  
Orders: nb-customerservice@bio-techne.com  
General: novus@novusbio.com

### **Products Related to NBP1-18891**

---

NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

---

### **Limitations**

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit [www.novusbio.com/guarantee](http://www.novusbio.com/guarantee)

Earn gift cards/discounts by submitting a review: [www.novusbio.com/reviews/submit/NBP1-18891](http://www.novusbio.com/reviews/submit/NBP1-18891)

Earn gift cards/discounts by submitting a publication using this product:  
[www.novusbio.com/publications](http://www.novusbio.com/publications)

