

Product Datasheet

Collagen I alpha 1 Antibody - Azide and BSA Free NBP1-30054

Unit Size: 0.1 ml

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Reviews: 2 Publications: 82

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP1-30054

Updated 4/13/2025 v.20.1

**Earn rewards for product
reviews and publications.**

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP1-30054



NBP1-30054

Collagen I alpha 1 Antibody - Azide and BSA Free

Product Information	
Unit Size	0.1 ml
Concentration	Please see the vial label for concentration. If unlisted please contact technical services.
Storage	Store at -20C. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	No Preservative
Isotype	IgG
Purity	Antigen Affinity-purified
Buffer	PBS
Target Molecular Weight	140 kDa
Product Description	
Description	Recommended that the undiluted antibody be aliquoted into smaller working volumes (10-30 uL/vial depending on usage).
Host	Rabbit
Gene ID	1277
Gene Symbol	COL1A1
Species	Human, Mouse, Rat, Amphibian, Avian, Mammal, Sheep
Reactivity Notes	Based on the homology of the immunogen, this antibody is expected to recognize the collagen I alpha1 polypeptide in all mammals, birds, and amphibians.
Specificity/Sensitivity	This Collagen I alpha 1 antibody is specific for the ~ 140 kDa telopeptide portion of the collagen I alpha 1 polypeptide. The antibody works well for immunohistochemistry on paraformaldehyde-fixed sections with a simple antigen-retrieval protocol (incubate slides for 20 minutes at 90 degrees C in 10 mM sodium citrate (pH 6.0)/ 0.1 % Tween-20). Note that in paraffin sections of formaldehydefixed fibrotic mouse lung tissue, the antibody recognizes mature collagen I that has formed fibrils in the extracellular matrix.
Immunogen	This Collagen I alpha 1 antibody was raised against synthetic peptide corresponding to amino acid residues within the C-terminal telo peptide portion (aa1193-1218) of the human COL1A1, conjugated to KLH. Accession # P02452
Product Application Details	
Applications	Western Blot, Simple Western, Immunohistochemistry-Paraffin, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Frozen
Recommended Dilutions	Western Blot 1:1000, Simple Western 1:100, Immunohistochemistry 1:10-1:500, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin 1:100, Immunohistochemistry-Frozen 1:10-1:500



Application Notes


Use in Immunohistochemistry-Frozen sections was reported in the scientific literature (PMID: 21939397), IHC reactivity reported in scientific literature (PMID: 26045736). Use in Immunocytochemistry/immunofluorescence reported in scientific literature (PMID: 26651081).

In Simple Western only 10 - 15 μ L of the recommended dilution is used per data point.


See [Simple Western Antibody Database](#) for Simple Western validation: Tested in Human Lung and Human Kidney lysate 0.5 mg/mL, separated by Size, antibody dilution of 1:100. Separated by Size-Wes, Sally Sue/Peggy Sue.

Images


Western Blot: Collagen I alpha 1 Antibody - Azide and BSA Free [NBP1-30054] - Collagen I Antibody [NBP1-30054] - Rat lung lysate showing specific immunolabeling of the collagen protein with an observed molecular weight \sim 140.




Immunohistochemistry: Collagen I alpha 1 Antibody - Azide and BSA Free [NBP1-30054] - Immunohistochemical staining targeting Collagen I alpha 1 revealed severe disorganization of the annulus fibrosus (AF), and a decrease in staining intensity for group D. On the contrary, the arcshaped and layered AF structure with immunoreactivity for type I collagen staining were mostly maintained in group S when compared to group C. Presented images show the AF at site of surgery, defect, and transplant. Abbreviations: D: AF defect only group, S: AF defect treated with sheet transplantation group, C: sham group, NC: negative control of sample stained without primary antibody (scale bar=300 μ m). Image collected and cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/31463464/>) licensed under a CC-BY license.




Immunohistochemistry: Collagen I alpha 1 Antibody - Azide and BSA Free [NBP1-30054] - Collagen I Antibody [NBP1-30054] - Formaldehyde-fixed fibrotic mouse lung tissue. The antibody recognizes mature collagen I (red) that has formed fibrils in the extracellular matrix.



Simple Western: Collagen I alpha 1 Antibody - Azide and BSA Free [NBP1-30054] - Simple Western lane view shows a specific band for Collagen I Alpha 1 in 0.5 mg/ml of Human Lung (left) and Human Kidney (right) lysate. This experiment was performed under reducing conditions using the 66-440 kDa separation system. * Non-specific interaction with the 230 kDa Simple Western standard may be seen with this antibody



Western Blot: Collagen I alpha 1 Antibody - Azide and BSA Free [NBP1-30054] - TNC deficiency reduced kidney fibrosis in animal models. Deletion of TNC (TNC^{-/-}) significantly attenuated the induction of collagen I assessed by IF following UUO by approximately 30% at day 7 & 10 compared with their wild-type littermates (A, n = 4 for each time point, two-way ANOVA p < 0.05). Consistently, the expression of fibrosis markers, such as collagen I α , fibronectin & plasminogen activator inhibitor-1 (PAI-1), were significantly lower in TNC^{-/-} mice at UUO day 7 & 10 (B, n = 4 for each time point, two-way ANOVA, p < 0.05). Western blot showed that the proteins of collagen I α & α -SMA were also significantly reduced in TNC^{-/-} mice at UUO day 7 (C, n = 7, p < 0.05). Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/36522320>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Publications

Cho S, Rhee S, Madl C et al. Selective inhibition of stromal mechanosensing suppresses cardiac fibrosis *Nature* 2025-06-01 [PMID: 40307543]

C Lee, J Kim, J Han, D Oh, M Kim, H Jeong, TJ Kim, SW Kim, JN Kim, YS Seo, A Suzuki, JH Kim, Y Jung Formyl peptide receptor 2 determines sex-specific differences in the progression of nonalcoholic fatty liver disease and steatohepatitis *Nature Communications*, 2022-01-31;13(1):578. 2022-01-31 [PMID: 35102146] (Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Mouse)

Chiang YP, Li Z, He M et al. Sphingomyelin synthase-related protein SMSr is a phosphatidylethanolamine phospholipase C that promotes nonalcoholic fatty liver disease *Journal of Biological Chemistry* 2023-08-14 [PMID: 37586586] (Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Mouse)

Olmsted ZT, Paluh JL. A combined human gastruloid model of cardiogenesis and neurogenesis *iScience* 2022-06-17 [PMID: 35721464] (Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Mouse)

Yamanaka Y, Tajima T, Tsujimura Y et al. Adiponectin inhibits fibrosis of the palmar aponeurosis in Dupuytren's contracture in male patients *Bone & Joint Research* 2023-08-04 [PMID: 37536684] (Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Mouse)

S Dey, LM Udari, P RiveraHern, JJ Kwon, B Willis, JJ Easler, EL Fogel, S Pandol, J Kota Loss of miR-29a/b1 promotes inflammation and fibrosis in acute pancreatitis *JCI Insight*, 2021-10-08;0(0):. 2021-10-08 [PMID: 34464354] (Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Mouse)

Wang Q, Chen Z, Peng X et al. Neuraminidase 1 Exacerbating Aortic Dissection by Governing a Pro-Inflammatory Program in Macrophages *Frontiers in Cardiovascular Medicine* 2021-11-18 [PMID: 34869700] (Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Mouse)

Yang P, Hong MS, Fu C et al. Preexisting smooth muscle cells contribute to neointimal cell repopulation at an incidence varying widely among individual lesions. *Surgery* 2016-02-01 [PMID: 26387788] (Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Mouse)

Xie Q, Zhang M, Mao X et al. Matrix protein Tenascin-C promotes kidney fibrosis via STAT3 activation in response to tubular injury *Cell Death & Disease* 2022-12-15 [PMID: 36522320] (Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Mouse)

Zhuang R, Chen J, Cheng HS et al. Perivascular Fibrosis Is Mediated by a KLF10-IL-9 Signaling Axis in CD4+ T Cells *Circulation Research* 2022-05-27 [PMID: 35440172] (Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Mouse)

Ji S, Gumina D, McPeak K et al. Human placental villous stromal extracellular matrix regulates fetoplacental angiogenesis in severe fetal growth restriction *Clinical science (London, England : 1979)* 2021-04-27 [PMID: 33904582] (Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Mouse)

Wang Q, Sundar I, Li D et al. E-cigarette-Induced Pulmonary Inflammation and Dysregulated Repair are Mediated by nAChR $\alpha 7$ Receptor: Role of nAChR $\alpha 7$ in ACE2 Covid-19 receptor regulation *Research Square* 2022-11-07 [PMID: 32702718] (Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Mouse)

More publications at <http://www.novusbio.com/NBP1-30054>



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP1-30054

HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control
NBP1-77457PEP	Collagen I alpha 1 Antibody Blocking Peptide

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP1-30054

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

