

Product Datasheet

PGD Antibody - BSA Free NBP1-31589

Unit Size: 100 ul

Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 4

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP1-31589

Updated 9/9/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP1-31589



NBP1-31589

PGD Antibody - BSA Free


Product Information	
Unit Size	100 ul
Concentration	Concentrations vary lot to lot. See vial label for concentration. If unlisted please contact technical services.
Storage	Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.01% Thimerosal
Isotype	IgG
Purity	Antigen Affinity-purified
Buffer	0.1M Tris, 0.1M Glycine, 10% Glycerol
Target Molecular Weight	53 kDa

Product Description	
Description	Novus Biologicals Rabbit PGD Antibody - BSA Free (NBP1-31589) is a polyclonal antibody validated for use in IHC, WB and ICC/IF. Anti-PGD Antibody: Cited in 4 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	5226
Gene Symbol	PGD
Species	Human, Mouse
Reactivity Notes	Xenopus laevis (85%) and Zebrafish (86%).
Immunogen	Recombinant protein encompassing a sequence within the center region of human PGD. The exact sequence is proprietary.

Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry
Recommended Dilutions	Western Blot 1:500-1:3000, Immunohistochemistry 1:100-1:1000, Immunocytochemistry/ Immunofluorescence 1:100-1:1000, Immunohistochemistry-Paraffin 1:100-1:1000

Images

Western Blot: PGD Antibody [NBP1-31589] - Sample(30 ug whole cell lysate)A:293T B:MOLT4 C:Raji 10% SDS PAGE, antibody diluted at 1:1000.




PGD antibody [N1N3] detects PGD protein at cytoplasm by immunofluorescent analysis.

Sample: HeLa cells were fixed in 4% paraformaldehyde at RT for 15 min.

Green: PGD protein stained by PGD antibody [N1N3] (NBP1-31589) diluted at 1:1000.

Red: alpha Tubulin, a cytoskeleton marker, stained by alpha Tubulin antibody [GT114] diluted at 1:1000.

Blue: Hoechst 33342 staining.




PGD antibody [N1N3] detects PGD protein at cytoplasm by immunofluorescent analysis.

Sample: HeLa cells were fixed in ice-cold MeOH for 5 min.


Green: PGD protein stained by PGD antibody [N1N3] (NBP1-31589) diluted at 1:500.

Blue: Hoechst 33342 staining.



PGD antibody [N1N3] detects PGD protein by western blot analysis.

Mouse tissue extracts (50 ug) was separated by 10% SDS-PAGE, and the membrane was blotted with PGD antibody [N1N3] (NBP1-31589) diluted at 1:500. The HRP-conjugated anti-rabbit IgG antibody was used to detect the primary antibody.




PGD antibody [N1N3] detects PGD protein at cytoplasm on A549 xenograft by immunohistochemical analysis.


Sample: Paraffin-embedded A549 xenograft.

PGD antibody [N1N3] (NBP1-31589) dilution: 1:500.

Antigen Retrieval: (EDTA based, pH 8.0) buffer, 15min



PGD antibody [N1N3] detects PGD protein by western blot analysis. Mouse tissue extracts (50 ug) was separated by 10% SDS-PAGE, and the membrane was blotted with PGD antibody [N1N3] (NBP1-31589) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody was used to detect the primary antibody.



Publications

Emmanuel Benichou, Bolaji Seffou, Selin Topçu, Ophélie Renoult, Véronique Lenoir, Julien Planchais, Caroline Bonner, Catherine Postic, Carina Prip-Buus, Claire Pecqueur, Sandra Guilmeau, Marie-Clotilde Alves-Guerra, Renaud Dentin The transcription factor ChREBP Orchestrates liver carcinogenesis by coordinating the PI3K/AKT signaling and cancer metabolism *Nature Communications* 2024-02-29 [PMID: 38424041]

Pushpak Bhattacharjee, Die Wang, Dovile Anderson, Joshua N Buckler, Eveline de Geus, Feng Yan, Galina Polekhina, Ralf Schittenhelm, Darren J Creek, Lawrence D Harris, Anthony J Sadler The immune response to RNA suppresses nucleic acid synthesis by limiting ribose 5-phosphate *The EMBO Journal* 2024-05-22 [PMID: 38778156]

Elf S, Lin R, Xia S et al. Targeting 6-phosphogluconate dehydrogenase in the oxidative PPP sensitizes leukemia cells to antimalarial agent dihydroartemisinin. *Oncogene* 2016-06-06 [PMID: 27270429] (Human)

Lemons JM, Feng XJ, Bennett BD et al. Quiescent fibroblasts exhibit high metabolic activity. *PLoS Biol* 2010-10-01 [PMID: 21049082] (WB)



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP1-31589

HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control
NBP1-85925PEP	PGD Recombinant Protein Antigen

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP1-31589

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

