

Product Datasheet

SNX27 Antibody - BSA Free

NBP1-45283

Unit Size: 0.1 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 2

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP1-45283

Updated 9/9/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP1-45283



NBP1-45283


SNX27 Antibody - BSA Free

Product Information	
Unit Size	0.1 ml
Concentration	1 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.1% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	PBS and 30% Glycerol
Target Molecular Weight	65 kDa
Product Description	
Description	Novus Biologicals Rabbit SNX27 Antibody - BSA Free (NBP1-45283) is a polyclonal antibody validated for use in IHC, WB, ICC/IF, Simple Western and IP. Anti-SNX27 Antibody: Cited in 2 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	81609
Gene Symbol	SNX27
Species	Human, Mouse, Rat
Reactivity Notes	Predicted to react with rat, bovine and Zebrafish based on 100% sequence homology. Use in Rat reported in scientific literature (PMID:32413996).
Immunogen	Synthetic peptide made to an internal portion of human SNX27 (within residues 450-500). [Swiss-Prot# Q96L92]
Product Application Details	
Applications	Western Blot, Simple Western, Immunohistochemistry-Paraffin, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunoprecipitation
Recommended Dilutions	Western Blot 1 ug/ml, Simple Western 10 ug/ml, Immunohistochemistry 1:100, Immunocytochemistry/ Immunofluorescence 1:50-1:200, Immunoprecipitation reported in scientific literature (PMID 32413996), Immunohistochemistry-Paraffin 1:100
Application Notes	<p>In Western blot, a band is seen at approx. 65 kDa. Prior to immunostaining paraffin tissues, antigen retrieval with sodium citrate buffer (pH 6.0) is recommended.</p> <p>In Simple Western only 10 - 15 uL of the recommended dilution is used per data point.</p> <p>See Simple Western Antibody Database for Simple Western validation: Tested in Human Cerebellum and Mouse Cerebellum lysate 0.5 mg/mL, separated by Size, antibody dilution of 10 ug/mL. Separated by Size-Wes, Sally Sue/Peggy Sue.</p> <p>The observed molecular weight of the protein may vary from the listed predicted molecular weight due to post translational modifications, post translation cleavages, relative charges, and other experimental factors.</p>




Images


Western Blot: SNX27 Antibody [NBP1-45283] - Analysis of SNX27 on mouse brain extracts.




Immunocytochemistry/Immunofluorescence: SNX27 Antibody [NBP1-45283] - Detection of SNX27 (Green) in HeLa cells using NBP1-45283. Nuclei (Blue) are counterstained using Hoechst 33258.




Immunohistochemistry: SNX27 Antibody [NBP1-45283] - IHC analysis of SNX27 in mouse bladder using DAB with hematoxylin counterstain.



Western Blot: SNX27 Antibody [NBP1-45283] - Analysis of SNX27 antibody in A. 293T + tagged SNX27 protein and B. 293T empty vector.



Simple Western: SNX27 Antibody [NBP1-45283] - Simple Western lane view shows a specific band for SNX27 in 0.5 mg/ml of Human Cerebellum (left) and Mouse Cerebellum (right) lysate. This experiment was performed under reducing conditions using the 12-230 kDa separation system.



Publications

Choi HJ, Jang HJ, Park E et al. Sorting Nexin 27 Regulates the Lysosomal Degradation of Aquaporin-2 Protein in the Kidney Collecting Duct Cells 2020-05-13 [PMID: 32413996] (IP, Rat)

Ish-Shalom E, Meirow Y, Sade-Feldman M et al. Impaired SNX9 Expression in Immune Cells during Chronic Inflammation: Prognostic and Diagnostic Implications. *J. Immunol.* 2016-01-01 [PMID: 26608909] (WB, Mouse)

Procedures

Protocol specific for SNX27 Antibody (NBP1-45282)

SNX27 Antibody:

Western Blot Protocol

1. Perform SDS-PAGE (4-12% MOPS) on samples to be analyzed, loading 35 ug of total protein per lane.
2. Transfer proteins to Nitrocellulose according to the instructions provided by the manufacturer of the transfer apparatus.
3. Rinse membrane with dH₂O and then stain the blot using Ponceau S for 1-2 minutes to access the transfer of proteins onto the nitrocellulose membrane. Rinse the blot in water to remove excess stain and mark the lane locations and locations of molecular weight markers using a pencil.
4. Rinse the blot in TBS for approximately 5 minutes.
5. Block the membrane using 5% NFD_M + 1% BSA in TBS + Tween, 1 hour at RT.
6. Rinse the membrane in dH₂O and then wash the membrane in wash buffer [TBS + 0.1% Tween] 3 times for 10 minutes each.
7. Dilute the rabbit anti-SNX27 primary antibody (NBP1-45282) in blocking buffer and incubate 1 hour at room temperature.
8. Rinse the membrane in dH₂O and then wash the membrane in wash buffer [TBS + 0.1% Tween] 3 times for 10 minutes each.
9. Apply the diluted rabbit-IgG HRP-conjugated secondary antibody in blocking buffer (as per manufacturers instructions) and incubate 1 hour at room temperature.
10. Wash the blot in wash buffer [TBS + 0.1% Tween] 3 times for 10 minutes each (this step can be repeated as required to reduce background).
11. Apply the detection reagent of choice in accordance with the manufacturers instructions (Pierce ECL).

Note: Tween-20 can be added to the blocking or antibody dilution buffer at a final concentration of 0.05-0.2%, provided it does not interfere with antibody-antigen binding.



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP1-45283

H00006604-M01-100ug	SMARCD3 Antibody (1G6) - Azide and BSA Free
NBP1-45283PEP	SNX27 Antibody Blocking Peptide
NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP1-45283

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications



