

Product Datasheet

N-Cadherin Antibody (13A9) - BSA Free NBP1-48309

Unit Size: 0.1 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Reviews: 2 **Publications: 42**

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP1-48309

Updated 9/9/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP1-48309




NBP1-48309**N-Cadherin Antibody (13A9) - BSA Free**


Product Information	
Unit Size	0.1 ml
Concentration	1 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	13A9
Preservative	0.05% Sodium Azide
Isotype	IgG1
Purity	Protein G purified
Buffer	PBS
Target Molecular Weight	140 kDa
Product Description	
Description	Novus Biologicals Mouse N-Cadherin Antibody (13A9) - BSA Free (NBP1-48309) is a monoclonal antibody validated for use in IHC, WB, Flow, ICC/IF, Simple Western and IP. Anti-N-Cadherin Antibody: Cited in 42 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Mouse
Gene ID	1000
Gene Symbol	CDH2
Species	Human, Mouse, Rat
Marker	Mesenchymal Cells Marker
Immunogen	This N-Cadherin Antibody (13A9) was developed against the cytoplasmic domain of human N Cadherin [Swiss-Prot# P19022].
Product Application Details	
Applications	Western Blot, Simple Western, Immunohistochemistry-Paraffin, Flow Cytometry, Flow (Intracellular), Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunoprecipitation
Recommended Dilutions	Western Blot 0.5 ug/ml, Simple Western 1:50, Flow Cytometry, Immunohistochemistry 1:50-1:200, Immunocytochemistry/ Immunofluorescence 1:100, Immunoprecipitation 1:10-1:500, Immunohistochemistry-Paraffin 1:50-1:100, Flow (Intracellular)
Application Notes	In Western Blot a band is observed at approx. 140 kDa. In Simple Western only 10 - 15 uL of the recommended dilution is used per data point. See Simple Western Antibody Database for Simple Western validation: Tested in HeLa lysate 1.0 mg/mL, separated by Size, antibody dilution of 1:50, apparent MW was 155 kDa. Separated by Size-Wes, Sally Sue/Peggy Sue. The observed molecular weight of the protein may vary from the listed predicted molecular weight due to post translational modifications, post translation cleavages, relative charges, and other experimental factors.

Images


Immunocytochemistry: N-Cadherin Antibody (13A9) [NBP1-48309] - Immunostaining of original tumor, low passage, and high passage KCI-MENG1 cells. The original patient-derived tumor (top row) showed moderate and patchy immunoreactivity for epithelial membrane antigen (EMA); strong and diffuse immunostaining for progesterone receptor (PR); and a Ki-67 proliferative index of 2-3%. There was also strong immunostaining for N-cadherin and vimentin. KCI-MENG1-LP cells (middle row) and KCI-MENG1-HP cells (bottom row) maintained expression of EMA, N-cadherin, and vimentin but had significantly reduced PR expression compared to the original tumor. Whereas Ki-67 labeling was found in only a small number of cells in the original tumor and low passage cells, it was positive in virtually all P84 cells. Scale bar 50 μ m. Image collected and cropped by CiteAb from the following publication (<https://www.translational-medicine.com/content/13/1/227>), licensed under a CC-BY license.




Western Blot: N-Cadherin Antibody (13A9) [NBP1-48309] - Western blots showing a reduction in epithelial mesenchymal transition (EMT) markers upon PDK1 knockdown in Colo205 and HT29 cells using E cadherin (NBP2-19051), N cadherin (NBP1-48309) and B-actin antibody (NB600-501). The corresponding secondary antibodies used were either goat anti-rabbit IgG-HRP (NB7160) or goat anti-mouse IgG-HRP (NB7539). Image collected and cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/33738242>) licensed under a CC-BY license.




Western Blot: N-Cadherin Antibody (13A9) [NBP1-48309] - Analysis of N-Cadherin expression in HeLa whole cell lysate.



Immunocytochemistry: N-Cadherin Antibody (13A9) [NBP1-48309] - Human meningioma mouse xenograft model KCI-MENG1-LPSX generated with the spontaneously immortal cell line KCI-MENG1-LP. Tumors from immunocompromised SCID mice were dissected and the derivative cell line KCI-MENG1-LPSX CL was generated. The EMA, PR, and N-cadherin IHC of the mouse tumor highly resembled the original patient-derived tumor. The vimentin- and Ki-67-stained cells in the mouse tumor tissue were markedly more abundant and more intensely stained than in the original tumor. KCI-MENG1-LPSX CL cells displayed the same patterns of immunostaining as the high passage parent cell line KCI-MENG1-HP, including the loss of PR staining. Scale bar 50 μ m. Image collected and cropped by CiteAb from the following publication (<https://www.translational-medicine.com/content/13/1/227>), licensed under a CC-BY license.




Flow (Intracellular): N-Cadherin Antibody (13A9) [NBP1-48309] - An intracellular stain was performed on HeLa with NBP1-48309 and a matched isotype control. Cells were fixed with 4% PFA and then permeabilized with 0.1% saponin. Cells were incubated in an antibody dilution of 1 ug/mL for 30 minutes at room temperature, followed by Mouse F(ab)2 IgG (H+L) PE-conjugated Antibody.




Simple Western: N-Cadherin Antibody (13A9) [NBP1-48309] - Simple Western lane view shows a specific band for N Cadherin in 1.0 mg/mL of HeLa lysate. This experiment was performed under reducing conditions using the 12-230 kDa separation system.




Western Blot: N-Cadherin Antibody (13A9) [NBP1-48309] - Mitochondrial fission regulates migration and invasion in BRAF^{V600E} CRC cells through glucose metabolic reprogramming. Western blot showing a reduction in EMT markers upon vemurafenib treatment in BRAF^{V600E} cells. Image collected and cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/33738242/>) licensed under a CC-BY license.




Immunocytochemistry/Immunofluorescence: N-Cadherin Antibody (13A9) [NBP1-48309] - Adult mouse neural stem cells stained for N-Cadherin. Antibody at 1:100. ICC/IF image submitted by a verified customer review.




Immunohistochemistry: N-Cadherin Antibody (13A9) [NBP1-48309] - Immunohistochemical characterization of tissue cultures. Nuclei labeled with DAPI (blue). Whole-mount immunostaining of a cardiomyocyte-only spheroid 1 month in culture, N-cadherin (red) and EH-myomesin (green). Image collected and cropped by CiteAb from the following publication ([//pubmed.ncbi.nlm.nih.gov/32118040/](https://pubmed.ncbi.nlm.nih.gov/32118040/)) licensed under a CC-BY license.




Flow Cytometry: N-Cadherin Antibody (13A9) [NBP1-48309] - An intracellular stain was performed on U-251 MG cells with N-Cadherin Antibody (13A9) NBP1-48309AF594 (blue) and a matched isotype control (orange). Cells were fixed with 4% PFA and then permeabilized with 0.1% saponin. Cells were incubated in an antibody dilution of 2.5 ug/mL for 30 minutes at room temperature. Both antibodies were conjugated to Alexa Fluor 594.




Immunocytochemistry/Immunofluorescence: N-Cadherin Antibody (13A9) [NBP1-48309] - HEK 293 cells were fixed for 10 minutes using 10% formalin and then permeabilized for 5 minutes using 1X PBS + 0.5% Triton X-100. The cells were incubated with anti- at 5 ug/mL overnight at 4C and detected with an anti-mouse DyLight 488 (Green) at a 1:500 dilution. Actin was detected with Phalloidin 568 (Red) at a 1:200 dilution. Nuclei were counterstained with DAPI (Blue). Cells were imaged using a 40X objective.




Immunohistochemistry: N-Cadherin Antibody (13A9) [NBP1-48309] - IHC analysis of N Cadherin in mouse liver using DAB with hematoxylin counterstain.



Immunohistochemistry-Paraffin: N-Cadherin Antibody (13A9) [NBP1-48309] - Analysis of a FFPE tissue section of human brain using 1:200 dilution of N-Cadherin antibody. The staining was developed using HRP labeled anti-mouse secondary antibody and DAB reagent, and nuclei of cells were counter-stained with hematoxylin.




Flow Cytometry: N-Cadherin Antibody (13A9) [NBP1-48309] - An intracellular stain was performed on HeLa cells with NBP1-48309AF488 and a matched isotype control (orange). Cells were fixed with 4% PFA and then permeabilized with 0.1% saponin. Cells were incubated in an antibody dilution of 10 ug/mL for 30 minutes at room temperature. Both antibodies were conjugated to Alexa Fluor 488.




Copyright © 2018 Novus Biologicals

N-Cadherin (13A9) was detected in immersion fixed U-2 OS human osteosarcoma cell line using Mouse anti- N-Cadherin (13A9) Protein G Purified Monoclonal Antibody conjugated to Alexa Fluor® 647 (Catalog # NBP1-48309AF647) (light blue) at 10 µg/mL overnight at 4C. Cells were counterstained with DAPI (blue). Cells were imaged using a 100X objective and digitally deconvolved.




Copyright © 2025 Novus Biologicals, a Bio-Techne Brand

N-Cadherin (13A9) was detected in immersion fixed U-251 MG human glioblastoma cell line using Mouse anti-N-Cadherin (13A9) Protein G Purified Monoclonal Antibody conjugated to DyLight 550 (Catalog # NBP1-48309R) (red) at 10 µg/mL overnight at 4C. Cells were counterstained with DAPI (blue). Cells were imaged using a 100X objective and digitally deconvolved.



Copyright © 2025 Novus Biologicals, a Bio-Techne Brand

Immunohistochemical characterization of tissue, 2D-, and 3D-cultures. Nuclei are labeled with DAPI (blue) in all images. (A) Cryosection of an adult mouse heart immunostained for myomesin (red) and vimentin (green). (B) Confocal optical section on the substrate level of a spheroid made of hiPSC-CM and CF cultured for 1 month. Immunostaining for myosin heavy chain (red), and vimentin (green). (C) Confocal optical section above the substrate level of the same spheroid as shown in (B). (D) 2D-cultured cardiac fibroblasts immunostained for alpha-SMA (red) and all actin (green) shown as a maximum intensity projection of optical sections. (E) Single confocal optical sections perpendicular to the substrate through a CF-only, small spheroid (above) cultured for 5 days and immunostained for all actin (red) and alpha-SMA (green), and an optical section at midlevel (below). (F) Cryosection of fibroblast-only spheroid cultured for 3 weeks immunostained for vimentin (red) and alpha-SMA (green). (G) Western blot experiments and quantitative assessment of the ratio of alpha-SMA/vimentin ($n = 4$, $***p < 0.001$) demonstrating the difference of alpha-SMA content in 2D vs. 3D cultures of pure CF. (H) Whole-mount immunostaining of a cardiomyocyte-only spheroid 1 month in culture, N-cadherin (red) and EH-myomesin (green). (I) Cardiomyocyte-only spheroid 1 month in culture, whole-mount immunostained for all-actin (red) and laminin (green). Image collected and cropped by CiteAb from the following open publication (<https://pubmed.ncbi.nlm.nih.gov/32118040>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Publications

Padder RA, Bhat ZI, Ahmad Z et al. DRP1 Promotes BRAF(V600E)-Driven Tumor Progression and Metabolic Reprogramming in Colorectal Cancer *Frontiers in Oncology* 2021-03-02 [PMID: 33738242] (Western Blot, Human)

Donaldson RI, Buchanan OJ, Graham TL, Ross JD. Development of a Novel Epidural Hemorrhage Model in Swine. *Military medicine* 2023-01-10 [PMID: 34676417]

Belghasem M, Yin W, Lotfollahzadeh S et al. Tryptophan Metabolites Target Transmembrane and Immunoglobulin Domain-Containing 1 Signaling to Augment Renal Tubular Injury *The American journal of pathology* 2023-10-01 [PMID: 37676196] (WB, Mouse)

Lin YK, Zhang F, Lei WJ et al. Amnion-derived serum amyloid A1 participates in sterile inflammation of fetal membranes at parturition *Inflammation research : official journal of the European Histamine Research Society ... [et al.]* 2023-03-06 [PMID: 36879064]

?wierczewska M, Sterzy?ska K, Ruci?ski M et al. The response and resistance to drugs in ovarian cancer cell lines in 2D monolayers and 3D spheroids *Biomedicine & pharmacotherapy = Biomedecine & pharmacotherapie* 2023-07-11 [PMID: 37442067] (IHC, Human)

Huang M, Wang C, Yao Y et al. Mebendazole-Induced Blood-Testis Barrier Injury in Mice Testes by Disrupting Microtubules in Addition to Triggering Programmed Cell Death *International journal of molecular sciences* 2022-04-11 [PMID: 35457043]

Karim S, Burzangi AS, Ahmad A Et al. PI3K-AKT Pathway Modulation by Thymoquinone Limits Tumor Growth and Glycolytic Metabolism in Colorectal Cancer *Int J Mol Sci* 2022-02-26 [PMID: 35216429] (WB, Human)

Details:

Citation using the HRP version of this antibody.

Prasad A, Mahmood A, Gupta R Et Al. In cardiac muscle cells, both adrenergic agonists and antagonists induce reactive oxygen species from NOX2 but mutually attenuate each other's effects *European journal of pharmacology* 2021-07-13 [PMID: 34265295] (WB)

Matta R, Yousafzai MS, Murrell M, Gonzalez AL Endothelial cell secreted metalloproteinase-2 enhances neural stem cell N-cadherin expression, clustering, and migration *FASEB journal : official publication of the Federation of American Societies for Experimental Biology* 2021-02-01 [PMID: 33417253]

SAenz-de-Santa-Marla I, Celada L, San JosE MartInez A et al. Blockage of Squamous Cancer Cell Collective Invasion by FAK Inhibition Is Released by CAFs and MMP-2 *Cancers* 2020-12-10 [PMID: 33321813] (WB, Human)

Beauchamp P, Jackson C B et al. 3D Co-culture of hiPSC-Derived Cardiomyocytes With Cardiac Fibroblasts Improves Tissue-Like Features of Cardiac Spheroids. *Front Mol Biosci* 2020-03-03 [PMID: 32118040] (IF/IHC, Mouse)

Becker A B, Qian J et al. Heart and Nervous System Pathology in Compound Heterozygous Friedreich Ataxia. *J Neuropathol Exp Neurol* 2017-01-08 [PMID: 28789479] (IF/IHC, Human)

More publications at <http://www.novusbio.com/NBP1-48309>



Procedures

Protocol specific for N Cadherin Antibody (13A9) [NBP1-48309]

Western Blot Protocol

1. Perform SDS-PAGE (4-12% MOPS) on samples to be analyzed, loading 25 ug of total protein per lane.
 2. Transfer proteins to Nitrocellulose according to the instructions provided by the manufacturer of the transfer apparatus.
 3. Rinse membrane with dH₂O and then stain the blot using Ponceau S for 1-2 minutes to access the transfer of proteins onto the nitrocellulose membrane. Rinse the blot in water to remove excess stain and mark the lane locations and locations of molecular weight markers using a pencil.
 4. Rinse the blot in TBS for approximately 5 minutes.
 5. Block the membrane using 5% NFDM + 1% BSA in TBS + Tween, 1 hour at RT.
 6. Rinse the membrane in dH₂O and then wash the membrane in wash buffer [TBS + 0.1% Tween] 3 times for 10 minutes each.
 7. Dilute the anti-N Cadherin primary antibody (NBP1-48309) in blocking buffer and incubate 1 hour at room temperature.
 8. Rinse the membrane in dH₂O and then wash the membrane in wash buffer [TBS + 0.1% Tween] 3 times for 10 minutes each.
 9. Apply the diluted mouse-IgG HRP-conjugated secondary antibody in blocking buffer (as per manufacturers instructions) and incubate 1 hour at room temperature.
 10. Wash the blot in wash buffer [TBS + 0.1% Tween] 3 times for 10 minutes each (this step can be repeated as required to reduce background).
 11. Apply the detection reagent of choice in accordance with the manufacturers instructions (Pierce ECL).
- Note: Tween-20 can be added to the blocking or antibody dilution buffer at a final concentration of 0.05-0.2%, provided it does not interfere with antibody-antigen binding.

Immunohistochemistry-Paraffin Embedded Sections

Antigen Unmasking:

Bring slides to a boil in 10 mM sodium citrate buffer (pH 6.0) then maintain at a sub-boiling temperature for 10 minutes. Cool slides on bench-top for 30 minutes.

Staining:

1. Wash sections in deionized water three times for 5 minutes each.
2. Wash sections in wash buffer for 5 minutes.
3. Block each section with 100-400 ul blocking solution for 1 hour at room temperature.
4. Remove blocking solution and add 100-400 ul diluted primary antibody. Incubate overnight at 4 C.
5. Remove antibody solution and wash sections in wash buffer three times for 5 minutes each.
6. Add 100-400 ul biotinylated diluted secondary antibody. Incubate 30 minutes at room temperature.
7. Remove secondary antibody solution and wash sections three times with wash buffer for 5 minutes each.
8. Add 100-400 ul Streptavidin-HRP reagent to each section and incubate for 30 minutes at room temperature.
9. Wash sections three times in wash buffer for 5 minutes each.
10. Add 100-400 ul DAB substrate to each section and monitor staining closely.
11. As soon as the sections develop, immerse slides in deionized water.
12. Counterstain sections in hematoxylin.
13. Wash sections in deionized water two times for 5 minutes each.
14. Dehydrate sections.
15. Mount coverslips.

Immunocytochemistry Protocol

Culture cells to appropriate density in 35 mm culture dishes or 6-well plates.

1. Remove culture medium and add 10% formalin to the dish. Fix at room temperature for 30 minutes.
2. Remove the formalin and add ice cold methanol. Incubate for 5-10 minutes.
3. Remove methanol and add washing solution (i.e. PBS). Be sure to not let the specimen dry out. Wash three times for 10 minutes.

4. To block nonspecific antibody binding incubate in 10% normal goat serum from 1 hour to overnight at room temperature.
5. Add primary antibody at appropriate dilution and incubate at room temperature from 2 hours to overnight at room temperature.
6. Remove primary antibody and replace with washing solution. Wash three times for 10 minutes.
7. Add secondary antibody at appropriate dilution. Incubate for 1 hour at room temperature.
8. Remove antibody and replace with wash solution, then wash for 10 minutes. Add Hoechst 33258 to wash solution at 1:25,000 and incubate for 10 minutes. Wash a third time for 10 minutes.
9. Cells can be viewed directly after washing. The plates can also be stored in PBS containing Azide covered in Parafilm (TM). Cells can also be cover-slipped using Fluoromount, with appropriate sealing.

*The above information is only intended as a guide. The researcher should determine what protocol best meets their needs. Please follow safe laboratory procedures.





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP1-48309

NB800-PC1	HeLa Whole Cell Lysate
NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB7539	Goat anti-Mouse IgG (H+L) Secondary Antibody [HRP]
NBP1-97005-0.5mg	Mouse IgG1 Isotype Control (MG1)

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP1-48309

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

