

Product Datasheet

Cytokeratin, pan Antibody (C-11) - BSA Free NBP1-48348

Unit Size: 0.1 mg

Store at 4C. Do not freeze.

www.novusbio.com



technical@novusbio.com

Publications: 9

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP1-48348

Updated 9/9/2025 v.20.1

**Earn rewards for product
reviews and publications.**

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP1-48348



NBP1-48348


Cytokeratin, pan Antibody (C-11) - BSA Free

Product Information	
Unit Size	0.1 mg
Concentration	1 mg/ml
Storage	Store at 4C. Do not freeze.
Clonality	Monoclonal
Clone	C-11
Preservative	15mM Sodium Azide
Isotype	IgG1
Purity	Protein A purified
Buffer	Phosphate buffered saline (PBS), pH 7.4
Product Description	
Description	Novus Biologicals Mouse Cytokeratin, pan Antibody (C-11) - BSA Free (NBP1-48348) is a monoclonal antibody validated for use in IHC, WB, Flow, ICC/IF and IP. Anti-Cytokeratin, pan Antibody: Cited in 9 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Mouse
Gene ID	3848
Gene Symbol	KRT1
Species	Mouse, Mammal
Reactivity Notes	Mouse reactivity reported in (PMID:26574781). Please note that this antibody is reactive to Mouse and derived from the same host, Mouse. Mouse-On-Mouse blocking reagent may be needed for IHC and ICC experiments to reduce high background signal. You can find these reagents under catalog numbers PK-2200-NB and MP-2400-NB. Please contact Technical Support if you have any questions.
Specificity/Sensitivity	This Mouse monoclonal Cytokeratin (Pan-reactive) antibody reacts with Cytokeratin peptides 4, 5, 6, 8, 10, 13, 18. Cytokeratins are a member of intermediate filaments subfamily represented in epithelial tissues.
Immunogen	This Cytokeratin, pan Antibody (C-11) was developed against a keratin-enriched preparation from human epidermoid carcinoma cell line A431.
Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, Flow (Intracellular), Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunoprecipitation, CyTOF-ready
Recommended Dilutions	Western Blot 1-2 ug/ml, Immunohistochemistry 2-8 ug/ml, Immunocytochemistry/ Immunofluorescence 1:10 - 1:500, Immunoprecipitation 1:50, Immunohistochemistry-Paraffin 1:200, Flow (Intracellular) 1 ug/mL, CyTOF-ready
Application Notes	This antibody is CyTOF ready.




Images

Flow Cytometry: Cytokeratin, pan Antibody (C-11) [NBP1-48348] - An intracellular stain was performed on HeLa cells with pan Cytokeratin antibody (C-11) NBP1-48348AF488 (blue) and a matched isotype control (orange). Cells were fixed with 4% PFA and then permeabilized with 0.1% saponin. Cells were incubated in an antibody dilution of 5 ug/mL for 30 minutes at room temperature. Both antibodies were conjugated to Alexa Fluor 488.




Copyright © 2018 Novus Biologicals


Immunohistochemistry-Paraffin: Cytokeratin, pan Antibody (C-11) [NBP1-48348] - Staining of cytokeratin on guinea pig breast carcinoma.




Western Blot: Cytokeratin, pan Antibody (C-11) - BSA Free [NBP1-48348] - Analysis of human cytokeratins using mouse monoclonal antibody C-11 on lysates of HeLa cell line and MOLT-4 cell line (cytokeratin non-expressing cell line; negative control) under non-reducing and reducing conditions. Nitrocellulose membrane was probed with 2 ug/ml of mouse monoclonal antibody anti-cytokeratins followed by IRDye800-conjugated anti-mouse secondary antibody. Specific bands were detected for cytokeratins at approximately 45-55 kDa.




Immunocytochemistry/Immunofluorescence: Cytokeratin, pan Antibody (C-11) - BSA Free [NBP1-48348] - Staining of cytokeratins in Hep-2 cells using pan-cytokeratin antibody C-11 (diluted 1:400), detected with GAM IgG-Alexa Fluor®488 (diluted 1:200; green).




Flow (Intracellular): Cytokeratin, pan Antibody (C-11) - BSA Free [NBP1-48348] - Intracellular staining pattern of human peripheral whole blood spiked with MCF-7 cells stained using anti-Cytokeratins (C-11) purified antibody (concentration in sample 3 ug/ml, GAM APC).




Immunocytochemistry/Immunofluorescence: Cytokeratin, pan Antibody (C-11) [NBP1-48348] - HeLa cells were fixed for 10 minutes using 10% formalin and then permeabilized for 5 minutes using 1X PBS + 0.05% Triton X-100. The cells were incubated with anti-pan Cytokeratin (C-11) conjugated to Alexa Fluor 488 [NBP2-33134AF488] at 10 ug/mL for 1 hour at room temperature. Nuclei were counterstained with DAPI (Blue). Cells were imaged using a 40X objective. Image from the Alexa Fluor 488 version of this antibody.




Flow (Intracellular): Cytokeratin, pan Antibody (C-11) [NBP1-48348] - An intracellular stain was performed on HeLa cells with pan Cytokeratin (C-11) antibody NBP1-48348 (blue) and a matched isotype control NBP2-27287 (orange). Cells were fixed with 4% PFA and then permeabilized with 0.1% saponin. Cells were incubated in an antibody dilution of 1 ug/mL for 30 minutes, followed by mouse F(ab)2 IgG (H+L) PE-conjugated secondary antibody (F0102B, R&D Systems).




Flow Cytometry: Cytokeratin, pan Antibody (C-11) [NBP1-48348] - Intracellular flow cytometry analysis of cytokeratin expression in HT-29 human Caucasian colon adenocarcinoma cell line using anti-cytokeratin antibody (C-11) PE. Overlay with Isotype mouse IgG1 control.




Flow (Intracellular): Cytokeratin, pan Antibody (C-11) [NBP1-48348] - An intracellular stain was performed on HeLa cells with pan Cytokeratin antibody (C-11) NBP1-48348AF647 (blue) and a matched isotype control NBP2-27287AF647 (orange). Cells were fixed with 4% PFA and then permeabilized with 0.1% saponin. Cells were incubated in an antibody dilution of 2 $\mu\text{g}/\text{mL}$ for 30 minutes at room temperature. Both antibodies were conjugated to Alexa Fluor 647.



Flow Cytometry: Cytokeratin, pan Antibody (C-11) [NBP1-48348] - An intracellular stain was performed on A549 cells with pan Cytokeratin Antibody (C-11) NBP1-48348F488 (blue) and a matched isotype control (orange). Cells were fixed with 4% PFA and then permeabilized with 0.1% saponin. Cells were incubated in an antibody dilution of 5 $\mu\text{g}/\text{mL}$ for 30 minutes at room temperature. Both antibodies were conjugated to Alexa Fluor 488.



Analysis of human cytokeratin using mouse monoclonal antibody C-11 on lysates of HeLa cell line and MOLT-4 cell line (cytokeratin non-expressing cell line; negative control) under non-reducing and reducing conditions. Nitrocellulose membrane was probed with 2 $\mu\text{g}/\text{ml}$ of mouse monoclonal antibody anti-cytokeratin followed by IRDye800-conjugated anti-mouse secondary antibody. Specific bands were detected for cytokeratin at approximately 45-55 kDa.



Publications

Kleeman S, Thakir T, Demestichas B et al. Cystatin C is glucocorticoid responsive, directs recruitment of Trem2+ macrophages, and predicts failure of cancer immunotherapy Cell Genomics 2023-06-23 [PMID: 37601967] (IHC-Fr, Mouse)

McGowan SE, McCoy DM. Regulation of fibroblast lipid storage and myofibroblast phenotypes during alveolar septation in mice Am J Physiol Lung Cell Mol Physiol 2014-08-24 [PMID: 25150063]

Details:

Citation using the PE version of this antibody.

Papadaki MA, Mala A, Merodoulaki AC Et al. Investigating the Role of CTCs with Stem/EMT-like Features in Metastatic Breast Cancer Patients Treated with Eribulin Mesylate Cancers (Basel) 2022-08-26 [PMID: 36010896] (ICC/IF, Human)

Details:

Citation using the Alexa Fluor 488 version of this antibody.

Polioudaki H, Mala A, Gkimprxi E et al. Epithelial/Mesenchymal Characteristics and PD-L1 Co-Expression in CTCs of Metastatic Breast Cancer Patients Treated with Eribulin: Correlation with Clinical Outcome Cancers 2020-12-11 [PMID: 33322610]

Details:

We developed a method combining immunocytochemistry, indirect and direct immunofluorescence.

Papadaki MA, Koutsopoulos AV, Tsoulfas PG et al. Clinical Relevance of Immune Checkpoints on Circulating Tumor Cells in Breast Cancer Cancers (Basel) 2020-02-06 [PMID: 32041353] (ICC/IF, Human)

Ijsselsteijn ME, Brouwer TP, Abdulrahman Z et al. Cancer immunophenotyping by 7 colour multispectral imaging without tyramide signal amplification J Pathol Clin Res. 2018-01-01 [PMID: 30191683] (ICC/IF, Human)

Details:

Citation using the Alexa Fluor 647 form of this antibody.

Consentius C, Mirenska A, Jurisch A et al. In situ detection of CD73+ CD90+ CD105+ lineage: Mesenchymal stromal cells in human placenta and bone marrow specimens by chipcytometry. Cytometry A 2018-07-01 [PMID: 30211969] (FLOW, Human)

Ogata A, Kawashima M, Wakamiya T et al. Carotid artery stenosis with a high-intensity signal plaque on time-of-flight magnetic resonance angiography and association with evidence of intraplaque hypoxia J. Neurosurg. 2016-07-01 [PMID: 27367236] (ICC/IF, Human)

Sinha M, Lowell CA. Immune Defense Protein Expression in Highly Purified Mouse Lung Epithelial Cells. Am. J. Respir. Cell Mol. Biol. 2015 Nov 17 [PMID: 26574781]





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP1-48348

NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB7539	Goat anti-Mouse IgG (H+L) Secondary Antibody [HRP]
NBP1-97005-0.5mg	Mouse IgG1 Isotype Control (MG1)

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP1-48348

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

