

Product Datasheet

Goat anti-Bovine IgG (H+L) Secondary Antibody [DyLight 488] NBP1-74759

Unit Size: 1 mg

Store lyophilized antibody at 4C. Mix reconstituted liquid with an equal volume of glycerol, aliquot contents and freeze at -20C or below for long term storage.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP1-74759

Updated 10/30/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP1-74759



NBP1-74759

Goat anti-Bovine IgG (H+L) Secondary Antibody [DyLight 488]

Product Information	
Unit Size	1 mg
Concentration	LYOPH mg/ml
Storage	Store lyophilized antibody at 4C. Mix reconstituted liquid with an equal volume of glycerol, aliquot contents and freeze at -20C or below for long term storage.
Clonality	Polyclonal
Preservative	0.05% Sodium Azide
Reconstitution Instructions	Rehydrate with 1.1 ml of deionized water and let stand 30 minutes at room temperature to dissolve. (Product has been overfilled to ensure complete recovery.) Centrifuge to remove any particulates. Prepare fresh working dilution daily.
Isotype	IgG
Conjugate	DyLight 488
Purity	Affinity purified
Buffer	10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG free

Product Description	
Description	<p>Purity > 95% based on SDS-PAGE. Goat serum was obtained from healthy animals of US origin and under the care of a registered veterinarian. After reconstituted, prepare fresh working dilution daily.</p> <p>Product is stable for up to 4 weeks at 2-8C after rehydration. For extended storage after rehydration, add an equal volume of glycerol and store at -20C.</p>
Host	Goat
Species	Bovine
Reactivity Notes	Based on IEP, no reactivity is observed to non-immunoglobulin bovine serum immunoglobulins
Specificity/Sensitivity	Based on IEP, this Goat anti-Bovine IgG (H+L) Secondary Antibody [DyLight 488] reacts with heavy gamma chains on bovine IgG and light chains on all bovine immunoglobulins.
Immunogen	This Goat anti-Bovine IgG (H+L) Secondary Antibody [DyLight 488] was developed against purified bovine IgG (H&L).
Notes	DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.

Product Application Details	
Applications	Flow Cytometry, Immunomicroscopy
Recommended Dilutions	Flow Cytometry 1:20-1:2000, Immunomicroscopy 1:20-1:2000
Application Notes	This conjugate is suitable for immunomicroscopy and flow cytometry.



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Secondary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP1-74759

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

