

# Product Datasheet

## XBP1 Antibody - BSA Free NBP1-77253

Unit Size: 0.1 mg

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

[www.novusbio.com](http://www.novusbio.com)



[technical@novusbio.com](mailto:technical@novusbio.com)

**Reviews: 1 Publications: 4**

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:  
[www.novusbio.com/NBP1-77253](http://www.novusbio.com/NBP1-77253)

Updated 3/4/2026 v.20.1

**Earn rewards for product  
reviews and publications.**

Submit a publication at [www.novusbio.com/publications](http://www.novusbio.com/publications)

Submit a review at [www.novusbio.com/reviews/destination/NBP1-77253](http://www.novusbio.com/reviews/destination/NBP1-77253)



**NBP1-77253**

XBP1 Antibody - BSA Free


Product Information	
Unit Size	0.1 mg
Concentration	1 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.02% Sodium Azide
Isotype	IgG
Purity	Peptide affinity purified
Buffer	PBS
Target Molecular Weight	28, 40 kDa

Product Description	
Host	Rabbit
Gene ID	7494
Gene Symbol	XBP1
Species	Human, Mouse, Rat
Specificity/Sensitivity	Human XBP1 has two isoforms, including isoform 1 (261aa, 29kD) and isoform 2 (376aa, 40kD). Mouse XBP1 also has two isoforms, including isoform 1 (267aa, 29.6kD) and isoform 2 (371aa, 40kD). And Rat XBP1 has two isoforms as well, including isoform 1 (267aa, 30kD) and isoform 2 (371aa, 40kD). NBP1-77253 can detect human, mouse and rat.
Immunogen	Antibody was raised against a peptide corresponding to 17 amino acids near the amino terminus of human XBP1. The immunogen is located within amino acids 40 - 90 of XBP1. Amino Acid Sequence: GLPQARKRQRLTHLSPE


Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, ELISA, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry
Recommended Dilutions	Western Blot 0.25-2 ug/mL, ELISA 1:100-1:2000, Immunohistochemistry 2-5 ug/ml, Immunocytochemistry/ Immunofluorescence 10-20 ug/ml, Immunohistochemistry-Paraffin 2-5 ug/mL

**Images**


Immunocytochemistry/ Immunofluorescence: XBP1 Antibody - BSA Free [NBP1-77253] - Validation of XBP-1 in HeLa Cells Immunofluorescent analysis of methanol-fixed HeLa cells labeling XBP-1 with at 10 u/mL, followed by goat anti-rabbit IgG secondary antibody at 1/1000 dilution (red) and DAPI staining (blue). Alpha tubulin was stained with anti-alpha tubulin antibody following by goat anti-mouse IgG secondary antibody (green). Images were captured with confocal microscopy.




Western Blot: XBP1 Antibody - BSA Free [NBP1-77253] - KO Validation in 293T Cells. Loading: 10 u of lysate Antibodies: XBP-1, 4 u/mL and beta-actin, 1u/mL, 1 h incubation at RT in 5% NFDm/TBST. Secondary: Goat Anti-Rabbit IgG HRP conjugate at 1:10000 dilution.




Western Blot: XBP1 Antibody [NBP1-77253] - Loading: 15 ug of lysates per lane. Antibodies: XBP-1 (1 ug/mL), 1h incubation at RT in 5% NFDm/TBST. Secondary: Goat anti-rabbit IgG HRP conjugate at 1:10000 dilution. (1) the absence and (2) the p




Western Blot: XBP1 Antibody [NBP1-77253] - Loading: 15 ug of HepG2 cell lysate Antibodies: XBP-1, 1h incubation at RT in 5% NFDm/TBST. Secondary: Goat anti-rabbit IgG HRP conjugate at 1:10000 dilution. Lane1: 1 ug/mL Lane2: 2 ug/mL




Western Blot: XBP1 Antibody [NBP1-77253] - Loading: 15 ug of A549 cell lysate Antibodies: XBP-1, 1h incubation at RT in 5% NFDm/TBST. Secondary: Goat anti-rabbit IgG HRP conjugate at 1:10000 dilution. Lane1: 0.25 ug/mL Lane2: 0.5 ug/mL Lane3: 1 ug/mL




Western Blot: XBP1 Antibody [NBP1-77253] - Loading: 100 ng of human XBP-1 recombinant protein per lane. Antibodies: XBP-1 (A: 0.5 ug/mL, B: 1 ug/mL and C: 2 ug/mL), 1h incubation at RT in 5% NFDN/TBST. Secondary: Goat anti-rabbit IgG HRP conjugate at




Western Blot: XBP1 Antibody - BSA Free [NBP1-77253] - Validation in Mouse Pancreas Tissue. Loading: 15 u of lysates per lane. Antibodies: XBP-1, (4 u /mL), 1h incubation at RT in 5% NFDN/TBST. Secondary: Goat anti-rabbit IgG HRP conjugate at 1:10000 dilution.




Immunocytochemistry/ Immunofluorescence: XBP1 Antibody - BSA Free [NBP1-77253] - Validation of XBP-1 in Mouse Pancreas Tissue. Immunofluorescent analysis of 4% paraformaldehyde-fixed mouse pancreas tissue labeling XBP-1 with at 20 u/mL, followed by goat anti-rabbit IgG secondary antibody at 1/500 dilution (green) and DAPI staining (blue).




Immunocytochemistry/ Immunofluorescence: XBP1 Antibody - BSA Free [NBP1-77253] - Validation of XBP-1 in Human Liver Tissue. Immunofluorescent analysis of 4% paraformaldehyde-fixed Human Liver Tissue labeling XBP-1 with at 20 u/mL, followed by goat anti-rabbit IgG secondary antibody at 1/500 dilution (green) and DAPI staining (blue).




Immunocytochemistry/ Immunofluorescence: XBP1 Antibody - BSA Free [NBP1-77253] - Validation of XBP-1 in Human Pancreas Tissue. Immunofluorescent analysis of 4% paraformaldehyde-fixed human pancreas tissue labeling XBP-1 with at 20 u/mL, followed by goat anti-rabbit IgG secondary antibody at 1/500 dilution (green) and DAPI staining (blue).




Immunohistochemistry: XBP1 Antibody - BSA Free [NBP1-77253] - Validation of XBP1 in Mouse Spleen Tissue. Immunohistochemical analysis of paraffin-embedded mouse spleen tissue using anti-XBP1 antibody at 1 u/ml. Tissue was fixed with formaldehyde and blocked with 10% serum for 1 h at RT; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody overnight at 4 C. A goat anti-rabbit IgG H&L (HRP) at 1/250 was used as secondary. Counter stained with Hematoxylin.



Immunohistochemistry: XBP1 Antibody - BSA Free [NBP1-77253] - Validation of XBP1 in Human Liver Tissue. Immunohistochemical analysis of paraffin-embedded human liver tissue using anti-XBP1 antibody at 1 u/ml. Tissue was fixed with formaldehyde and blocked with 10% serum for 1 h at RT; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody overnight at 4C. A goat anti-rabbit IgG H&L (HRP) at 1/250 was used as secondary. Counter stained with Hematoxylin.



Immunohistochemistry: XBP1 Antibody - BSA Free [NBP1-77253] - Validation of XBP1 in Rat Spleen Tissue. Immunohistochemical analysis of paraffin-embedded rat spleen tissue using anti-XBP1 antibody at 1 u/ml. Tissue was fixed with formaldehyde and blocked with 10% serum for 1 h at RT; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody overnight at 4 C. A goat anti-rabbit IgG H&L (HRP) at 1/250 was used as secondary. Counter stained with Hematoxylin.



## Publications

Hu JM, Chang YL, Hsieh CC, Huang SM The Synergistic Cytotoxic Effects of GW5074 and Sorafenib by Impacting Mitochondrial Functions in Human Colorectal Cancer Cell Lines *Frontiers in oncology* 2022-06-07 [PMID: 35747833] (WB)

Liu Y, Xu D, Wang L et al. MBTPS2 exacerbates albuminuria in streptozotocin-induced type I diabetic nephropathy by promoting endoplasmic reticulum stress-mediated renal damage *Arch. Physiol. Biochem.* 2020-04-07 [PMID: 32255378]

Lee HJ, Kim JH, Hong S et al. Proteomics-based functional studies reveal that galectin-3 plays a protective role in the pathogenesis of intestinal Behcet's disease *Sci Rep* 2019-08-12 [PMID: 31406212] (WB, Mouse, Human)

Zhang W, Zhong W, Sun Q et al. Hepatic overproduction of 13- HODE due to ALOX15 upregulation contributes to alcohol-induced liver injury in mice. *Sci Rep* 2017-08-21 [PMID: 28827690] (WB, Mouse)





### **Novus Biologicals USA**

10730 E. Briarwood Avenue  
Centennial, CO 80112  
USA  
Phone: 303.730.1950  
Toll Free: 1.888.506.6887  
Fax: 303.730.1966  
nb-customerservice@bio-techne.com

### **Bio-Techne Canada**

21 Canmotor Ave  
Toronto, ON M8Z 4E6  
Canada  
Phone: 905.827.6400  
Toll Free: 855.668.8722  
Fax: 905.827.6402  
canada.inquires@bio-techne.com

### **Bio-Techne Ltd**

19 Barton Lane  
Abingdon Science Park  
Abingdon, OX14 3NB, United Kingdom  
Phone: (44) (0) 1235 529449  
Free Phone: 0800 37 34 15  
Fax: (44) (0) 1235 533420  
info.EMEA@bio-techne.com

### **General Contact Information**

www.novusbio.com  
Technical Support: nb-technical@bio-techne.com  
Orders: nb-customerservice@bio-techne.com  
General: novus@novusbio.com

### **Products Related to NBP1-77253**

---

NBP1-77253PEP	XBP1 Antibody Blocking Peptide
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

---

### **Limitations**

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit [www.novusbio.com/guarantee](http://www.novusbio.com/guarantee)

Earn gift cards/discounts by submitting a review: [www.novusbio.com/reviews/submit/NBP1-77253](http://www.novusbio.com/reviews/submit/NBP1-77253)

Earn gift cards/discounts by submitting a publication using this product:  
[www.novusbio.com/publications](http://www.novusbio.com/publications)

