

Product Datasheet

FAM98B Antibody - BSA Free NBP1-83475

Unit Size: 0.1 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 3

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP1-83475

Updated 12/2/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP1-83475



NBP1-83475

FAM98B Antibody - BSA Free


Product Information	
Unit Size	0.1 ml
Concentration	Concentrations vary lot to lot. See vial label for concentration. If unlisted please contact technical services.
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.02% Sodium Azide
Isotype	IgG
Purity	Affinity purified
Buffer	PBS (pH 7.2) and 40% Glycerol

Product Description	
Description	Novus Biologicals Rabbit FAM98B Antibody - BSA Free (NBP1-83475) is a polyclonal antibody validated for use in IHC and WB. Anti-FAM98B Antibody: Cited in 3 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	283742
Gene Symbol	FAM98B
Species	Human
Reactivity Notes	Reactivity reported in scientific literature (PMID: 24608264)
Immunogen	This antibody was developed against Recombinant Protein corresponding to amino acids: STTSDIPHMLNQVESKVKDILSKVQKNHVGKPLLKMDLNSEQAEQLERINDALS CEYECRRRMLMKRLDVTVQSGFGWSDRAKVKTTDDIARIYQPKRYALSPKTTITM AHLLAAREDLSKIIRTSSGTSREKTACAI


Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, Immunohistochemistry
Recommended Dilutions	Western Blot 0.04-0.4 ug/ml, Immunohistochemistry 1:20 - 1:50, Immunohistochemistry-Paraffin 1:20 - 1:50
Application Notes	WB reported in scientific literature (PMID: 25378478). For IHC-Paraffin, HIER pH 6 retrieval is recommended.

Images


Immunohistochemistry-Paraffin: FAM98B Antibody [NBP1-83475] - Staining of human testis shows strong nuclear positivity in cells in seminiferous ducts.




Western Blot: FAM98B Antibody [NBP1-83475] - Analysis using Anti-FAM98B antibody NBP1-83475 (A) shows similar pattern to independent antibody NBP1-83474 (B).




Immunohistochemistry-Paraffin: FAM98B Antibody [NBP1-83475] - Staining of human colon.




Immunohistochemistry-Paraffin: FAM98B Antibody [NBP1-83475] - Staining of human testis.




Immunohistochemistry-Paraffin: FAM98B Antibody [NBP1-83475] - Staining of human cerebral cortex shows strong nuclear positivity in neuronal cells.




Immunohistochemistry-Paraffin: FAM98B Antibody [NBP1-83475] - Staining of human cerebral cortex, gastrointestinal, skeletal muscle and testis using Anti-FAM98B antibody NBP1-83475 (A) shows similar protein distribution across tissues to independent antibody NBP1-83474 (B).




Immunohistochemistry-Paraffin: FAM98B Antibody [NBP1-83475] - Staining of human colon shows strong nuclear positivity in glandular cells.




Immunohistochemistry-Paraffin: FAM98B Antibody [NBP1-83475] - Staining of human parathyroid gland shows strong nuclear positivity in glandular cells.



Immunohistochemistry-Paraffin: FAM98B Antibody [NBP1-83475] - Staining of human skeletal muscle shows strong nuclear positivity in myocytes.



Western Blot: FAM98B Antibody [NBP1-83475] - RTCB is required for induction of XBP1s during plasma cell differentiation. B220+ splenocytes of control (Rtcbfl/fl or Rtcbfl/+), Rtcbfl/+Cd23-Cre or Rtcbfl/flCd23-Cre mice were stimulated with 20 µg/ml LPS for 4 days. A, B Protein levels of RTCB, tRNA ligase complex members (DDX1, FAM98B, CGI-99) & XBP1 were monitored by Western blot analysis (n > 3). C FACS-sorted CD138+ CD22- plasmablasts or CD138- CD22low pre-plasmablasts were probed for expression of the indicated proteins by Western blot analysis (n = 2). D Relative mRNA levels of *Rtcb*, *Xbp1s*, total *Xbp1*, *Edem1*, µM & µS were analyzed by RT-qPCR in fractionated LPS-stimulated cells at day 4 (n = 4, mean expression levels & SEM are displayed). Expression levels were normalized to *Actb* mRNA levels & to B cells from control mice. An unpaired Student's t-test was used to analyze the statistical significance of differences in mRNA levels (*P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001). Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/25378478>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Publications

Jurkin J, Henkel T, Nielsen AF et al. The mammalian tRNA ligase complex mediates splicing of XBP1 mRNA and controls antibody secretion in plasma cells. *EMBO J* 2014-12-17 [PMID: 25378478] (WB, Human)

Popow J, Jurkin J, Schleiffer A et al. Analysis of orthologous groups reveals Archease and DDX1 as tRNA splicing factors. *Nature* 2014-07-03 [PMID: 24870230]

Perez-Gonzalez A, Pazo A, Navajas R et al. hCLE/C14orf166 Associates with DDX1-HSPC117-FAM98B in a Novel Transcription-Dependent Shuttling RNA-Transporting Complex. *PLoS One* 2014-01-01 [PMID: 24608264]



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP1-83475

NBP1-83475PEP	FAM98B Recombinant Protein Antigen
NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP1-83475

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

