

# Product Datasheet

## FoxP2 Antibody - BSA Free

### NBP1-86671

Unit Size: 0.1 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

[www.novusbio.com](http://www.novusbio.com)



[technical@novusbio.com](mailto:technical@novusbio.com)

**Publications: 5**

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:  
[www.novusbio.com/NBP1-86671](http://www.novusbio.com/NBP1-86671)

Updated 1/25/2026 v.20.1

Earn rewards for product  
reviews and publications.

Submit a publication at [www.novusbio.com/publications](http://www.novusbio.com/publications)

Submit a review at [www.novusbio.com/reviews/destination/NBP1-86671](http://www.novusbio.com/reviews/destination/NBP1-86671)



**NBP1-86671**

FoxP2 Antibody - BSA Free

Product Information	
<b>Unit Size</b>	0.1 ml
<b>Concentration</b>	Concentrations vary lot to lot. See vial label for concentration. If unlisted please contact technical services.
<b>Storage</b>	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
<b>Clonality</b>	Polyclonal
<b>Preservative</b>	0.02% Sodium Azide
<b>Isotype</b>	IgG
<b>Purity</b>	Affinity purified
<b>Buffer</b>	PBS (pH 7.2) and 40% Glycerol


Product Description	
<b>Description</b>	Novus Biologicals Rabbit FoxP2 Antibody - BSA Free (NBP1-86671) is a polyclonal antibody validated for use in IHC, WB and ICC/IF. Anti-FoxP2 Antibody: Cited in 5 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
<b>Host</b>	Rabbit
<b>Gene ID</b>	93986
<b>Gene Symbol</b>	FOXP2
<b>Species</b>	Human, Mouse, Rat
<b>Reactivity Notes</b>	Rat reactivity reported in scientific literature (PMID: 25926446). Mouse reactivity reported in scientific literature (PMID: 26407299).
<b>Immunogen</b>	This antibody was developed against Recombinant Protein corresponding to amino acids: AQQLVFQQQLLQMQQLQQQHLHLSLQRQGLISIPPGQAALPVQSLPQAGLSPA EIQQLWKEVTGVHSMEDNGIKHGGLDLTTNNSSTTSSNTSKASPPITHHS

Product Application Details	
<b>Applications</b>	Western Blot, Immunohistochemistry-Paraffin, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry
<b>Recommended Dilutions</b>	Western Blot 0.04-0.4 ug/ml, Immunohistochemistry 1:200 - 1:500, Immunocytochemistry/ Immunofluorescence Reported in scientific literature (PMID: 25926446 and 25926446)., Immunohistochemistry-Paraffin 1:200-1:500
<b>Application Notes</b>	IHC-Paraffin, HIER pH 6 retrieval is recommended.




## Images


FoxP2-Antibody-Immunohistochemistry-NBP1-86671-img0009.jpg




Immunohistochemistry-Paraffin: FoxP2 Antibody [NBP1-86671] - Staining of human cerebral cortex shows moderate to strong nuclear positivity in neurons.




Immunohistochemistry-Paraffin: FoxP2 Antibody [NBP1-86671] - Staining of human rectum shows moderate nuclear positivity in glandular cells.




Immunohistochemistry-Paraffin: FoxP2 Antibody [NBP1-86671] - Staining of human skeletal muscle shows no positivity in striated muscle fibers as expected.




Immunohistochemistry-Paraffin: FoxP2 Antibody [NBP1-86671] - Staining of human skin shows moderate nuclear positivity in deep epidermal cells.




Immunocytochemistry/ Immunofluorescence: FoxP2 Antibody [NBP1-86671] - Patterns of neocortical layer markers in the PPC. A) TBR-1 heavily labeled cells in Layer 2 as well as scattered cells in Layer 3. As in the APC many cells in layers 2 & 3 exhibited the deep marker CTIP2 (B). Only widely scattered cells exhibited FOXP2 & DAARP 32 & NOR1 (B,C). The other three makers exhibited very different patterns: CUX 1 staining (E) was strong throughout layers 2 & 3, BRN2 staining much more modest in the same regions, CART was restricted to the middle of layer 2 (F). Scale bar = 200µm. Dorsal to top, lateral to right. Image collected & cropped by CiteAb from the following publication (<https://dx.plos.org/10.1371/journal.pone.0138541>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.




Immunocytochemistry/ Immunofluorescence: FoxP2 Antibody [NBP1-86671] - Patterns of neocortical layer markers in the OT. A) TBR-1 (A), BRN2 (D) & CART (E) were only found scattered in the very deepest regions of the OT. All four of the deep laminar markers heavily labeled the region. Both CTIP2 & FOXP2 cells were broadly present in Layer 2 & scattered in Layer 3 (Fig 6b). On the medial side most FOXP2 cells coexpressed CTIP2 but the percentage of cells with both markers was reduced laterally. DARRP-32 cells were dense on the lateral side near the APC & in deep regions of the OT, while NOR1 cells were found in Layer 2 in the medial OT (Fig 6c). Scale bar = 200µm. Dorsal to top, lateral to right. Image collected & cropped by CiteAb from the following publication (<https://dx.plos.org/10.1371/journal.pone.0138541>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.




Immunocytochemistry/ Immunofluorescence: FoxP2 Antibody [NBP1-86671] - Patterns of neocortical layer markers in the APC. A) TBR-1 heavily labeled cells in Layer 2 as well as scattered cells in Layer 3. Of the 4 deep layer markers (B,C), only CTIP2 exhibited dense staining. The other three (FOXP2, NOR1 & DAARP32) labeled sparse number in Layers 1–3. The dense staining for FOXP2 & DAARP32 seen at the bottom of the figures sharply demarcates the APC from the more ventral OT. The other three makers exhibited very different patterns: BRN2 staining was found more in the ventral APC (D), CUX 1 in the deeper portions of both Layer 2 & 3 (E), & CART in the middle of Layer 2 (F). Scale bar = 200µm. Dorsal to top, lateral to right. Image collected & cropped by CiteAb from the following publication (<https://dx.plos.org/10.1371/journal.pone.0138541>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Immunocytochemistry/ Immunofluorescence: FoxP2 Antibody [NBP1-86671] - Patterns of neocortical layer markers in the AONpP.A). TBR1-labelled cells were found throughout Layer 2 of the AONpP as well as in the tenia tecta & mitral cell layer of the OB. B, C) Deep markers were differentially distributed in the region. Layer 2 exhibited dense & evenly-spread CTIP2-positive cells (Fig 3b), while NOR1 was found primarily in the dorsal portion of the structure (Fig 3c, top) Cells expressing the other two marker were rare & found primarily in layer 1: DARRP-32 (note dense staining in the glomerular layer of the OB at left, an area containing large numbers of dopaminergic interneurons, Fig 3c; Liu et al, 2013) & FOXP2 (most often found near the OB, Fig 3b). CTIP2 stained cells were also found in layer 1 but never in cells that expressed one of the other markers. The superficial markers were also differentially distributed. Both BRN2 (Fig 3d) & CUX1 (Fig 3e) were observed primarily in deep cells (except in pars medialis, where CUX1-labeled cells spanned the entire region) with highest densities in the region under the LOT (pars lateralis). All CUX1 cells also expressed BRN2, & over 90% of CUX1 & BRN2 cells also expressed CTIP2. The anti-BRN2 antibody also labeled the LOT (right) & RMS (core of the olfactory peduncle). Scale bar = 200µm. Dorsal to top, lateral to right. Image collected & cropped by CiteAb from the following publication (<https://dx.plos.org/10.1371/journal.pone.0138541>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Analysis in human cell line RH-30.



## Publications

Aquiles A, Fiordelisio T, Luna-Munguia H, Concha L. Altered functional connectivity and network excitability in a model of cortical dysplasia *Scientific Reports* 2023-07-30 [PMID: 37518675]

Brunjes PC, Osterberg SK, et al. Developmental Markers Expressed in Neocortical Layers Are Differentially Exhibited in Olfactory Cortex. *PLoS One* 2015-01-01 [PMID: 26407299] (IF/IHC, Mouse)

Abdi A, Mallet N, Mohamed FY et al. Prototypic and Arkypallidal Neurons in the Dopamine-Intact External Globus Pallidus. *J Neurosci* 2015-04-29 [PMID: 25926446] (ICC/IF, Rat)

Reimers-Kipping S, Hevers W, Paabo S et al. Humanized Foxp2 specifically affects cortico-basal ganglia circuits. *Neuroscience* 2011-02-01 [PMID: 21111790]

Enard W, Gehre S, Hammerschmidt K et al. A humanized version of Foxp2 affects cortico-basal ganglia circuits in mice. *Cell* 2009-05-01 [PMID: 19490899]



### **Novus Biologicals USA**

10730 E. Briarwood Avenue  
Centennial, CO 80112  
USA  
Phone: 303.730.1950  
Toll Free: 1.888.506.6887  
Fax: 303.730.1966  
nb-customerservice@bio-techne.com

### **Bio-Techne Canada**

21 Canmotor Ave  
Toronto, ON M8Z 4E6  
Canada  
Phone: 905.827.6400  
Toll Free: 855.668.8722  
Fax: 905.827.6402  
canada.inquires@bio-techne.com

### **Bio-Techne Ltd**

19 Barton Lane  
Abingdon Science Park  
Abingdon, OX14 3NB, United Kingdom  
Phone: (44) (0) 1235 529449  
Free Phone: 0800 37 34 15  
Fax: (44) (0) 1235 533420  
info.EMEA@bio-techne.com

### **General Contact Information**

www.novusbio.com  
Technical Support: nb-technical@bio-techne.com  
Orders: nb-customerservice@bio-techne.com  
General: novus@novusbio.com

### **Products Related to NBP1-86671**

---

NBP1-86671PEP	FoxP2 Recombinant Protein Antigen
NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

---

### **Limitations**

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit [www.novusbio.com/guarantee](http://www.novusbio.com/guarantee)

Earn gift cards/discounts by submitting a review: [www.novusbio.com/reviews/submit/NBP1-86671](http://www.novusbio.com/reviews/submit/NBP1-86671)

Earn gift cards/discounts by submitting a publication using this product:  
[www.novusbio.com/publications](http://www.novusbio.com/publications)

