

Product Datasheet

PHGDH Antibody - BSA Free

NBP1-87311

Unit Size: 0.1 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 7

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP1-87311

Updated 12/2/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP1-87311



NBP1-87311

PHGDH Antibody - BSA Free

Product Information	
Unit Size	0.1 ml
Concentration	Concentrations vary lot to lot. See vial label for concentration. If unlisted please contact technical services.
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.02% Sodium Azide
Isotype	IgG
Purity	Affinity purified
Buffer	PBS (pH 7.2) and 40% Glycerol


Product Description	
Description	Novus Biologicals Rabbit PHGDH Antibody - BSA Free (NBP1-87311) is a polyclonal antibody validated for use in IHC, WB, ICC/IF and Simple Western. Anti-PHGDH Antibody: Cited in 7 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	26227
Gene Symbol	PHGDH
Species	Human, Mouse, Rat
Immunogen	This antibody was developed against Recombinant Protein corresponding to amino acids: LEEIWPLCDFITVHTPLLPSTTGLLNDNTFAQCKKGVRV VNCARGGIVDEGALL RALQSGQCAGAALDVFTEEPPRDRALVDHENVISCPHLGASTKEAQSRCGEEI AVQFVDM

Product Application Details	
Applications	Western Blot, Simple Western, Immunohistochemistry-Paraffin, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry
Recommended Dilutions	Western Blot 0.04-0.4 ug/ml, Simple Western 1:20, Immunohistochemistry 1:500 - 1:1000, Immunocytochemistry/ Immunofluorescence 0.25-2 ug/ml, Immunohistochemistry-Paraffin 1:500 - 1:1000
Application Notes	ICC/IF Fixation Permeabilization: Use PFA/Triton X-100. IHC-Paraffin HIER pH6 retrieval is recommended. In Simple Western only 10 - 15 uL of the recommended dilution is used per data point. See Simple Western Antibody Database for Simple Western validation: Tested in RT-4 and U-251MG, separated by Size, antibody dilution of 1:20, apparent MW was 56 kDa. Separated by Size-Wes, Sally Sue/Peggy Sue.




Images


Western Blot: PHGDH Antibody [NBP1-87311] - Analysis in human cell lines SK-MEL-30 and A-549. Corresponding RNA-seq data are presented for the same cell lines. Loading control: Anti-COX411.



Immunocytochemistry/Immunofluorescence: PHGDH Antibody [NBP1-87311] - Staining of human cell line U-2 OS shows localization to plasma membrane & cytosol. Antibody staining is shown in green.




Immunohistochemistry-Paraffin: PHGDH Antibody [NBP1-87311] - Immunohistochemical staining of human cerebral cortex shows moderate to strong cytoplasmic positivity in neuronal and glial cells.




Western Blot: PHGDH Antibody [NBP1-87311] - Analysis in mouse cell line NIH-3T3 and rat cell line NBT-II.




Immunohistochemistry-Paraffin: PHGDH Antibody [NBP1-87311] - Immunohistochemical staining of human skeletal muscle shows no positivity as expected.




Immunohistochemistry-Paraffin: PHGDH Antibody [NBP1-87311] - Immunohistochemical staining of human breast cancer shows moderate to strong cytoplasmic positivity in tumor cells.




Immunohistochemistry-Paraffin: PHGDH Antibody [NBP1-87311] - Immunohistochemical staining of human skin shows moderate cytoplasmic positivity in epidermal cells.




Simple Western: PHGDH Antibody [NBP1-87311] - Simple Western lane view shows a specific band for PHGDH in 0.2 mg/ml of RT-4 (Left) and U-251MG (Right) lysate. This experiment was performed under reducing conditions using the 12-230 kDa separation system.




Simple Western: PHGDH Antibody [NBP1-87311] - Electropherogram image(s) of corresponding Simple Western lane view. PHGDH antibody was used at 1:20 dilution on RT-4 and U-251MG lysate(s).



Knockdown of PNKP or TDP1 leads to increase in expression of PSAT1 and PHGDH. Western blot analysis of XRCC1, PHGDH and PSAT1 expression upon KD of PNKP (A) or TDP1 (B). β -Actin served as a loading control. (C) TDP1 mRNA levels after KD of TDP1, quantified by qPCR and normalised to GAPDH.



Knockdown of PNKP or TDP1 leads to increase in expression of PSAT1 and PHGDH. Western blot analysis of XRCC1, PHGDH and PSAT1 expression upon KD of PNKP (A) or TDP1 (B). β -Actin served as a loading control. (C) TDP1 mRNA levels after KD of TDP1, quantified by qPCR and normalised to GAPDH. Image collected and cropped by CiteAb from the following open publication (<https://pubmed.ncbi.nlm.nih.gov/25800737>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Publications

Ishida CT, Zhang Y, Bianchetti E et al. Metabolic Reprogramming by Dual Akt/ERK Inhibition Through Imipridones Elicits Unique Vulnerabilities in Glioblastoma. *Clin. Cancer Res.* 2018-07-23 [PMID: 30037819] (WB, Human)

Gina M DeNicola, Pei-Hsuan Chen, Edouard Mullarky et al. NRF2 regulates serine biosynthesis in non-small cell lung cancer. *Nature Genetics* 2015-10-19 [PMID: 26482881] (WB, Human)

Mattaini KR, Brignole EJ, Kini M et al. An epitope tag alters phosphoglycerate dehydrogenase structure and impairs ability to support cell proliferation. *Cancer Metab* 2015-01-01 [PMID: 25926973] (WB, ICC/IF, Human)

Markkanen E, Fischer R, Ledentcova M et al. Cells deficient in base-excision repair reveal cancer hallmarks originating from adjustments to genetic instability. *Nucleic Acids Res* 2015-03-23 [PMID: 25800737] (WB, Human)

Maddocks OD, Berkers CR, Mason SM et al. Serine starvation induces stress and p53-dependent metabolic remodelling in cancer cells. *Nature* 2013-01-24 [PMID: 23242140]

Nilsson LM, Forshell TZ, Rimpi S et al. Mouse genetics suggests cell-context dependency for Myc-regulated metabolic enzymes during tumorigenesis. *PLoS Genet* 2012-01-01 [PMID: 22438825]

Possemato R, Marks KM, Shaul YD et al. Functional genomics reveal that the serine synthesis pathway is essential in breast cancer. *Nature* 2011-08-01 [PMID: 21760589]



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP1-87311

NBP1-87311PEP	PHGDH Recombinant Protein Antigen
NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP1-87311

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications



