

Product Datasheet

Glutaminase Antibody - BSA Free NBP1-89766

Unit Size: 0.1 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 5

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP1-89766

Updated 12/2/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP1-89766



NBP1-89766


Glutaminase Antibody - BSA Free

Product Information	
Unit Size	0.1 ml
Concentration	Concentrations vary lot to lot. See vial label for concentration. If unlisted please contact technical services.
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.02% Sodium Azide
Isotype	IgG
Purity	Affinity purified
Buffer	PBS (pH 7.2) and 40% Glycerol
Product Description	
Description	Novus Biologicals Rabbit Glutaminase Antibody - BSA Free (NBP1-89766) is a polyclonal antibody validated for use in IHC, WB and ICC/IF. Anti-Glutaminase Antibody: Cited in 5 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	2744
Gene Symbol	GLS
Species	Human, Mouse, Rat
Reactivity Notes	Mouse reactivity reported in scientific literature (PMID: 25798620). Rat reactivity reported in scientific literature (PMID: 31265816).
Immunogen	This antibody was developed against Recombinant Protein corresponding to amino acids: DRWNNTPMDEALHFGHHDVFKILQEYQVQYTPQGSDNGKENQTVHKNLDG L
Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry
Recommended Dilutions	Western Blot 0.04-0.4 ug/ml, Immunohistochemistry 1:50 - 1:200, Immunocytochemistry/ Immunofluorescence 0.25-2 ug/ml, Immunohistochemistry-Paraffin 1:50 - 1:200
Application Notes	For IHC-Paraffin, HIER pH 6 retrieval is recommended. ICC/IF Fixation Permeabilization: Use PFA/Triton X-100.




Images


Immunohistochemistry-Paraffin: Glutaminase Antibody [NBP1-89766] - Staining in human kidney and skeletal muscle tissues using NBP1-89766 antibody. Corresponding GLS RNA-seq data are presented for the same tissues.




Immunocytochemistry/Immunofluorescence: Glutaminase Antibody [NBP1-89766] - Staining of human cell line SiHa shows localization to mitochondria. Antibody staining is shown in green.




Immunohistochemistry-Paraffin: Glutaminase Antibody [NBP1-89766] - Staining of human cerebral cortex shows moderate to strong granular cytoplasmic positivity in neurons.




Immunohistochemistry-Paraffin: Glutaminase Antibody [NBP1-89766] - Staining of human duodenum shows moderate granular cytoplasmic positivity in glandular cells.




Immunohistochemistry-Paraffin: Glutaminase Antibody [NBP1-89766] - Staining of human kidney shows strong granular cytoplasmic positivity in cells in tubules.




Immunohistochemistry-Paraffin: Glutaminase Antibody [NBP1-89766] - Staining of human skeletal muscle shows no positivity in myocytes as expected.




Analysis in human cell line EFO-21.




Glutaminase-Antibody-Western-Blot-NBP1-89766-img0016.jpg




Western Blot: Glutaminase Antibody [NBP1-89766] - Region-based expression of the pyruvate recycling pathway marker protein glutaminase (GLT) the VMN of icv Lz-pretreated male & female rats. GLT protein was measured by Western blot in the rostral (a; $F(7,40) = 12.20$; $p < 0.0001$), middle (b; $F(7,40) = 17.39$; $p < 0.0001$), & caudal (c; $F(7,40) = 11.67$; $p < 0.0001$) VMN of V/V, V/INS, Lz/V, & Lz/INS groups of male (M; left-hand side) & female (F; right-hand side) rats. * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$; **** $p < 0.0001$. For each VMN segment, data depict mean normalized GAD O.D. values \pm S.E.M. Results are summarized in the table at bottom; \uparrow & \downarrow denote a relative increase or decrease, respectively, between treatment groups; no change (NC) between groups is also indicated Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/33238883>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Western Blot: Glutaminase Antibody [NBP1-89766] - Region-based expression of the pyruvate recycling pathway marker protein glutaminase (GLT) the VMN of icv Lz-pretreated male & female rats. GLT protein was measured by Western blot in the rostral (a; $F(7,40) = 12.20$; $p < 0.0001$), middle (b; $F(7,40) = 17.39$; $p < 0.0001$), & caudal (c; $F(7,40) = 11.67$; $p < 0.0001$) VMN of V/V, V/INS, Lz/V, & Lz/INS groups of male (M; left-hand side) & female (F; right-hand side) rats. * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$; **** $p < 0.0001$. For each VMN segment, data depict mean normalized GAD O.D. values \pm S.E.M. Results are summarized in the table at bottom; \uparrow & \downarrow denote a relative increase or decrease, respectively, between treatment groups; no change (NC) between groups is also indicated Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/33238883>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Western Blot: Glutaminase Antibody [NBP1-89766] - Region-based expression of the pyruvate recycling pathway marker protein glutaminase (GLT) the VMN of icv Lz-pretreated male & female rats. GLT protein was measured by Western blot in the rostral (a; $F(7,40) = 12.20$; $p < 0.0001$), middle (b; $F(7,40) = 17.39$; $p < 0.0001$), & caudal (c; $F(7,40) = 11.67$; $p < 0.0001$) VMN of V/V, V/INS, Lz/V, & Lz/INS groups of male (M; left-hand side) & female (F; right-hand side) rats. * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$; **** $p < 0.0001$. For each VMN segment, data depict mean normalized GAD O.D. values \pm S.E.M. Results are summarized in the table at bottom; \uparrow & \downarrow denote a relative increase or decrease, respectively, between treatment groups; no change (NC) between groups is also indicated Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/33238883>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Publications

Uddin MM, Ibrahim MMH, Briski KP Sex-dimorphic neuroestradiol regulation of ventromedial hypothalamic nucleus glucoregulatory transmitter and glycogen metabolism enzyme protein expression in the rat BMC Neurosci 2020-11-25 [PMID: 33238883] (Immunohistochemistry, Mouse)

Emmanuel Benichou, Bolaji Seffou, Selin Topçu, Ophélie Renoult, Véronique Lenoir, Julien Planchais, Caroline Bonner, Catherine Postic, Carina Prip-Buus, Claire Pecqueur, Sandra Guilmeau, Marie-Clotilde Alves-Guerra, Renaud Dentin The transcription factor ChREBP Orchestrates liver carcinogenesis by coordinating the PI3K/AKT signaling and cancer metabolism Nature Communications 2024-02-29 [PMID: 38424041]

Briski KP, Napit PR, Alhamyani A et al. Sex-Dimorphic Octadecaneuropeptide (ODN) Regulation of Ventromedial Hypothalamic Nucleus Glucoregulatory Neuron Function and Counterregulatory Hormone Secretion ASN neuro 2023 -05-17 [PMID: 37194319] (WB, Rat)

Uddin MM, Mahmood ASMH, Ibrahim MMH, Briski KP Sex-Dimorphic Estrogen Receptor Regulation of Ventromedial Hypothalamic Nucleus Glucoregulatory Neuron Adrenergic Receptor Expression in Hypoglycemic Male and Female Rats Brain Res. 2019-06-29 [PMID: 31265816] (WB, Rat)

Tanaka K, Sasayama T, Irino Y et al. Compensatory glutamine metabolism promotes glioblastoma resistance to mTOR inhibitor treatment. J Clin Invest 2015-04-01 [PMID: 25798620] (IF/IHC, Mouse)





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP1-89766

NBP1-89766PEP	Glutaminase Recombinant Protein Antigen
NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP1-89766

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

