

# Product Datasheet

## A2BP1 Antibody (D8H8) - BSA Free NBP2-13169

Unit Size: 0.1 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

[www.novusbio.com](http://www.novusbio.com)



[technical@novusbio.com](mailto:technical@novusbio.com)

### Publications: 4

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:  
[www.novusbio.com/NBP2-13169](http://www.novusbio.com/NBP2-13169)

Updated 9/9/2025 v.20.1

Earn rewards for product  
reviews and publications.

Submit a publication at [www.novusbio.com/publications](http://www.novusbio.com/publications)

Submit a review at [www.novusbio.com/reviews/destination/NBP2-13169](http://www.novusbio.com/reviews/destination/NBP2-13169)



**NBP2-13169**


A2BP1 Antibody (D8H8) - BSA Free

Product Information	
Unit Size	0.1 ml
Concentration	1.0 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	D8H8
Preservative	0.05% Sodium Azide
Isotype	IgG1 Kappa
Purity	Protein G purified
Buffer	Tris-Glycine (pH 7.5) and 0.15M NaCl
Product Description	
Description	Novus Biologicals Mouse A2BP1 Antibody (D8H8) - BSA Free (NBP2-13169) is a monoclonal antibody validated for use in IHC, WB, ICC/IF, Simple Western and IP. Anti-A2BP1 Antibody: Cited in 4 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Mouse
Gene ID	54715
Gene Symbol	RBFOX1
Species	Human, Mouse
Specificity/Sensitivity	Does not cross react with paralogue FOX2
Immunogen	Mouse recombinant A2BP1 [Swiss-Prot# Q9JJ43]
Product Application Details	
Applications	Western Blot, Simple Western, Immunohistochemistry-Paraffin, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Frozen, Immunoprecipitation
Recommended Dilutions	Western Blot 1:1000, Simple Western 1:20, Immunohistochemistry 1:200, Immunocytochemistry/ Immunofluorescence 1:50-1:100, Immunoprecipitation 1:10-1:100, Immunohistochemistry-Paraffin 1:200, Immunohistochemistry-Frozen reported in scientific literature (PMID 30001398)
Application Notes	<p>In Western Blot, a band is seen 50-55 kDa representing A2BP1. Prior to immunostaining paraffin tissues, antigen retrieval with sodium citrate buffer (pH 6.0) is recommended.</p> <p>In Simple Western only 10 - 15 uL of the recommended dilution is used per data point.</p> <p>See <a href="#">Simple Western Antibody Database</a> for Simple Western validation: Tested in Human Brain lysate 0.5 mg/mL, separated by Size, antibody dilution of 1:20, apparent MW was 54 kDa. Separated by Size-Wes, Sally Sue/Peggy Sue.</p>




## Images


Western Blot: A2BP1 Antibody (D8H8) [NBP2-13169] - Western blot analysis of A2BP1 expression in 1) human brain and 2) mouse brain tissue lysates.



Immunocytochemistry/Immunofluorescence: A2BP1 Antibody (D8H8) [NBP2-13169] - A2BP1 antibody was tested in Neuro-2a cells with Dylight 488 (green). Nuclei and alpha-tubulin were counterstained with DAPI (blue) and Dylight 550 (red).



Immunohistochemistry: A2BP1 Antibody (D8H8) [NBP2-13169] - IHC staining of A2BP1 in mouse brain using DAB with hematoxylin counterstain.



Simple Western: A2BP1 Antibody (D8H8) [NBP2-13169] - Simple Western lane view shows a specific band for A2BP1 in 0.5 mg/ml of Human Brain lysate. This experiment was performed under reducing conditions using the 12-230 kDa separation system.



## Publications

Gu L, Caprioli J, Piri N Rbfox1 expression in amacrine cells is restricted to GABAergic and VGlut3 glycinergic cells Bioscience reports 2022-06-22 [PMID: 35730583] (IF/IHC, Mouse)

Gu L, Kawaguchi R, Caprioli J, Piri N The effect of Rbfox2 modulation on retinal transcriptome and visual function J Mol Med (Berl) 2020-11-06 [PMID: 33184471]

Lund C, Yellapragada V, Vuoristo S et al. Characterization of the human GnRH neuron developmental transcriptome using a GNRH1-TdTomato reporter line in human pluripotent stem cells Dis Model Mech 2020-01-29 [PMID: 31996360] (PCR, Human)

Gu L, Bok D, Yu F et al. Downregulation of splicing regulator RBFOX1 compromises visual depth perception. PLoS ONE. 2018-07-12 [PMID: 30001398] (IHC-Fr, Mouse)



## Procedures

### Western blot Protocol Specific for A2BP1 Antibody (D8H8) [NBP2-13169]

A2BP1 Antibody (D8H8):

Western Blot Protocol

1. Perform SDS-PAGE on samples to be analyzed, loading 40 ug of total protein per lane.
  2. Transfer proteins to membrane according to the instructions provided by the manufacturer of the membrane and transfer apparatus.
  3. Stain according to standard Ponceau S procedure (or similar product) to assess transfer success, and mark molecular weight standards where appropriate.
  4. Rinse the blot.
  5. Block the membrane using standard blocking buffer for at least 1 hour.
  6. Wash the membrane in wash buffer three times for 10 minutes each.
  7. Dilute primary antibody in blocking buffer and incubate 1 hour at room temperature.
  8. Wash the membrane in wash buffer three times for 10 minutes each.
  9. Apply the diluted HRP conjugated secondary antibody in blocking buffer (as per manufacturers instructions) and incubate 1 hour at room temperature.
  10. Wash the blot in wash buffer three times for 10 minutes each (this step can be repeated as required to reduce background).
  11. Apply the detection reagent of choice in accordance with the manufacturers instructions.
- Note: Tween-20 can be added to the blocking or antibody dilution buffer at a final concentration of 0.05-0.2%.

\*The above information is only intended as a guide. The researcher should determine what protocol best meets their needs. Please follow safe laboratory procedures.

### Immunocytochemistry/Immunofluorescence protocol for A2BP1 Antibody (NBP2-13169)

A2BP1 Antibody (D8H8):

Immunocytochemistry Protocol

Culture cells to appropriate density in 35 mm culture dishes or 6-well plates.

1. Remove culture medium and add 10% formalin to the dish. Fix at room temperature for 30 minutes.
2. Remove the formalin and add ice cold methanol. Incubate for 5-10 minutes.
3. Remove methanol and add washing solution (i.e. PBS). Be sure to not let the specimen dry out. Wash three times for 10 minutes.
4. To block nonspecific antibody binding incubate in 10% normal goat serum from 1 hour to overnight at room temperature.
5. Add primary antibody at appropriate dilution and incubate at room temperature from 2 hours to overnight at room temperature.
6. Remove primary antibody and replace with washing solution. Wash three times for 10 minutes.
7. Add secondary antibody at appropriate dilution. Incubate for 1 hour at room temperature.
8. Remove antibody and replace with wash solution, then wash for 10 minutes. Add Hoechst 33258 to wash solution at 1:25,000 and incubate for 10 minutes. Wash a third time for 10 minutes.
9. Cells can be viewed directly after washing. The plates can also be stored in PBS containing Azide covered in Parafilm (TM). Cells can also be cover-slipped using Fluoromount, with appropriate sealing.

\*The above information is only intended as a guide. The researcher should determine what protocol best meets their needs. Please follow safe laboratory procedures.



**Immunohistochemistry-Paraffin Embedded Sections Protocol Specific for A2BP1 Antibody (D8H8) [NBP2-13169]**

A2BP1 Antibody (D8H8):

Immunohistochemistry-Paraffin Embedded Sections

**Antigen Unmasking:**

Bring slides to a boil in 10 mM sodium citrate buffer (pH 6.0) then maintain at a sub-boiling temperature for 10 minutes. Cool slides on bench-top for 30 minutes.

**Staining:**

1. Wash sections in deionized water three times for 5 minutes each.
2. Wash sections in wash buffer for 5 minutes.
3. Block each section with 100-400 ul blocking solution for 1 hour at room temperature.
4. Remove blocking solution and add 100-400 ul diluted primary antibody. Incubate overnight at 4 C.
5. Remove antibody solution and wash sections in wash buffer three times for 5 minutes each.
6. Add 100-400 ul biotinylated diluted secondary antibody. Incubate 30 minutes at room temperature.
7. Remove secondary antibody solution and wash sections three times with wash buffer for 5 minutes each.
8. Add 100-400 ul Streptavidin-HRP reagent to each section and incubate for 30 minutes at room temperature.
9. Wash sections three times in wash buffer for 5 minutes each.
10. Add 100-400 ul DAB substrate to each section and monitor staining closely.
11. As soon as the sections develop, immerse slides in deionized water.
12. Counterstain sections in hematoxylin.
13. Wash sections in deionized water two times for 5 minutes each.
14. Dehydrate sections.
15. Mount coverslips.

\*The above information is only intended as a guide. The researcher should determine what protocol best meets their needs. Please follow safe laboratory procedures.





### **Novus Biologicals USA**

10730 E. Briarwood Avenue  
Centennial, CO 80112  
USA  
Phone: 303.730.1950  
Toll Free: 1.888.506.6887  
Fax: 303.730.1966  
nb-customerservice@bio-techne.com

### **Bio-Techne Canada**

21 Canmotor Ave  
Toronto, ON M8Z 4E6  
Canada  
Phone: 905.827.6400  
Toll Free: 855.668.8722  
Fax: 905.827.6402  
canada.inquires@bio-techne.com

### **Bio-Techne Ltd**

19 Barton Lane  
Abingdon Science Park  
Abingdon, OX14 3NB, United Kingdom  
Phone: (44) (0) 1235 529449  
Free Phone: 0800 37 34 15  
Fax: (44) (0) 1235 533420  
info.EMEA@bio-techne.com

### **General Contact Information**

www.novusbio.com  
Technical Support: nb-technical@bio-techne.com  
Orders: nb-customerservice@bio-techne.com  
General: novus@novusbio.com

### **Products Related to NBP2-13169**

---

NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB7539	Goat anti-Mouse IgG (H+L) Secondary Antibody [HRP]
NBP1-43319-0.5mg	Mouse IgG1 Kappa Isotype Control (P3.6.2.8.1)

---

### **Limitations**

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit [www.novusbio.com/guarantee](http://www.novusbio.com/guarantee)

Earn gift cards/discounts by submitting a review: [www.novusbio.com/reviews/submit/NBP2-13169](http://www.novusbio.com/reviews/submit/NBP2-13169)

Earn gift cards/discounts by submitting a publication using this product:  
[www.novusbio.com/publications](http://www.novusbio.com/publications)

