

Product Datasheet

GADD153/CHOP Antibody - BSA Free NBP2-13172

Unit Size: 0.1 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Reviews: 1 **Publications: 27**

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-13172

Updated 9/9/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-13172



NBP2-13172

GADD153/CHOP Antibody - BSA Free


Product Information	
Unit Size	0.1 ml
Concentration	1 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.02% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	PBS

Product Description	
Description	Novus Biologicals Rabbit GADD153/CHOP Antibody - BSA Free (NBP2-13172) is a polyclonal antibody validated for use in IHC, WB, Flow and ICC/IF. Anti-GADD153/CHOP Antibody: Cited in 27 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	1649
Gene Symbol	DDIT3
Species	Human, Mouse, Rat, Primate
Reactivity Notes	Mouse reactivity reported in scientific literature (PMID:32814096).
Marker	ER Stress Marker
Immunogen	A synthetic peptide made to an internal portion of the human CHOP/GADD153 protein (between residues 100-150) [UniProt P35638]


Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, Flow Cytometry, Flow (Intracellular), Immunocytochemistry/ Immunofluorescence, Immunohistochemistry
Recommended Dilutions	Western Blot 1:1000, Flow Cytometry 2-5 ug/million cells, Immunohistochemistry 1:400, Immunocytochemistry/ Immunofluorescence 1-5 ug/ml, Immunohistochemistry-Paraffin 1:400, Flow (Intracellular) 2-5 ug/million cells

Images


Western Blot: GADD153/CHOP Antibody - BSA Free [NBP2-13172] - GADD153/CHOP Antibody [NBP2-13172] - NASTRp induces ER stress and eventually leads to cell death with Bim up-regulation. NASTRp induced ER stress and activated UPR led to apoptosis with Bim induction in NSCLC cell lines. Cells were treated with the indicated concentrations of NASTRp for 24 hours and cell lysates were subjected to SDS-PAGE/Western blot analysis using the indicated antibodies. Image collected and cropped by CiteAb from the following publication ([//doi.org/10.1371/journal.pone.0122628](https://doi.org/10.1371/journal.pone.0122628)) licensed under a CC-BY license.




Immunocytochemistry/Immunofluorescence: GADD153/CHOP Antibody [NBP2-13172] - HeLa cells were fixed in 4% paraformaldehyde for 10 minutes and permeabilized in 0.5% Triton X-100 in PBS for 5 minutes. The cells were incubated with anti-GADD153/CHOP Antibody NBP2-13172 at 1 ug/ml overnight at 4C and detected with an anti-rabbit Dylight 488 (Green) at a 1:1000 dilution for 60 minutes. Nuclei were counterstained with DAPI (Blue). Cells were imaged using a 100X objective and digitally deconvolved.




Western Blot: GADD153/CHOP Antibody - BSA Free [NBP2-13172] - GADD153/CHOP Antibody [NBP2-13172] - Effects of psoralen on ER-stress related protein expression in SMMC7721. Western blot assays showing the effects of psoralen and thapsigargin on ER-stress related protein expression. Image collected and cropped by CiteAb from the following publication (<https://biolres.biomedcentral.com/articles/10.1186/s40659-019-0241-8>), licensed under a CC-BY license.




Immunohistochemistry: GADD153/CHOP Antibody [NBP2-13172] - Mouse liver tissue. Antigen retrieval: Citrate buffer pH=6.0, Blocking: 10% BSA in 0.1% Tween in TBS, Primary antibody: 1:100 in 1% BSA in 0.1% Tween in TBS, Incubation: Overnight at 4C, Secondary antibody: 1:200 incubation for 1hr at room temperature, AB complex: 1:100 for 1hr at room temperature Development DAB. IHC-P image submitted by a verified customer review.




Flow Cytometry: GADD153/CHOP Antibody [NBP2-13172] - An intracellular stain was performed on SK-MEL-28 cells with GADD153/CHOP Antibody NBP2-13172F (blue) and a matched isotype control (orange). Cells were fixed with 4% PFA and then permeabilized with 0.1% saponin. Cells were incubated in an antibody dilution of 5 ug/mL for 30 minutes at room temperature. Both antibodies were conjugated to FITC.




Western Blot: GADD153/CHOP Antibody [NBP2-13172] - CHOP/GADD153 Antibody [NBP2-13172] - WB analysis of CHOP/GADD153 in tunicamycin treated (+) and untreated (-) HeLa cell lysate.




Immunohistochemistry: GADD153/CHOP Antibody [NBP2-13172] - IHC staining of CHOP/GADD153 in mouse colon using DAB with hematoxylin counterstain.




Western Blot: GADD153/CHOP Antibody - BSA Free [NBP2-13172] - Effects of psoralen on ER-stress related protein expression in SMMC7721. a Western blot assays showing the effects of psoralen & thapsigargin on ER-stress related protein expression. b Western blot assays showing the effects of TUDC on the expression of ER-stress related protein induced by PSO. c The mRNA levels of GRP78 & DDIT3 with TUDC under PSO & TG treatment. Values are mean \pm SD (n = 3) & * is means compared to the Con group, **P < 0.01 Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/31277690>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.




Western Blot: GADD153/CHOP Antibody - BSA Free [NBP2-13172] - Reduction of fibronectin prevents Dex-induced ER stress in human GTM-3 cells. (A) GTM3 cells were transfected with a plasmid expressing CRISPR-Cas9 targeting fibronectin (CR-FN) & then treated with Veh or Dex for 48 hours. Fibronectin knockdown partially reduced ER stress as evident from reduced GRP78 & CHOP levels after Dex treatment (n = 2). (B) Human GTM3 cells were treated with Dex (100 nM) with or without CR-FN for 48 hours. Fixed cells were stained for fibronectin & KDEL to examine Dex-induced fibronectin & ER stress. Fibronectin knockdown reduced fibronectin & ER stress in Dex-treated TM cells (n = 2). Image collected & cropped by CiteAb from the following publication (<https://www.nature.com/articles/s41598-017-14938-0>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.




Western Blot: GADD153/CHOP Antibody - BSA Free [NBP2-13172] - TM cells are sensitive to ER stress from Dex-induced abnormal ECM compared to other ocular cells: Primary human TM cells (A), RPE (B) & corneal fibroblasts (C) were treated with Veh or Dex for 3 days & immunostained for fibronectin & KDEL. Although fibronectin was increased in all of these cell types, ER stress was only observed in TM cells. (D) Human GTM3 or ARPE-19 cells were treated with Veh or Dex for 3 days. Fibronectin & chronic ER stress markers ATF4 & CHOP were examined in cellular lysates (n = 2). Although Dex increased fibronectin in both GTM3 & ARPE-19 cells, chronic ER stress was only observed in GTM3 cells indicating that TM cells are more prone to ER stress from Dex-induced ECM accumulation. (E) GTM-3 or ARPE-19 cells were treated with cellular fibronectin (cFn) for 2 days & ER stress was examined by Western blot analysis. Image collected & cropped by CiteAb from the following publication (<https://www.nature.com/articles/s41598-017-14938-0>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Western Blot: GADD153/CHOP Antibody - BSA Free [NBP2-13172] - Overexpression of extracellular fibronectin induces ER stress in primary human TM cells. (A) Primary human TM cells (n = 2 strains) were treated with cellular fibronectin (10 ug/ml) for 48 hours & cellular lysates were analyzed for ER stress markers. (B) Primary human TM cells (n = 2 strains) were treated with cellular fibronectin (10 ug/ml) for 48 hours & fixed cells were stained with fibronectin & KDEL antibodies. Cellular fibronectin increased extracellular fibronectin & ER stress in GTM3 cells (n = 2). Scale bar = 50 μ m. (C) GTM3 cells were transduced with fibronectin activation lentiviral particles for 48 hours & stained with fibronectin & KDEL antibodies. Increased fibronectin expression alone is sufficient to induce ER stress in TM cells (n = 2). Scale bar = 50 μ m. Image collected & cropped by CiteAb from the following publication (<https://www.nature.com/articles/s41598-017-14938-0>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.




Western Blot: GADD153/CHOP Antibody - BSA Free [NBP2-13172] - NASTRp induces ER stress & eventually leads to cell death with Bim up-regulation. (A) qRT-PCR analysis of biomarkers of ER stress/UPR related genes in NCI-H441 treated with 20 μ M NASTRp in time course manner. (B) NASTRp induced ER stress & activated UPR led to apoptosis with Bim induction in NSCLC cell lines. Cells were treated with the indicated concentrations of NASTRp for 24 hours & cell lysates were subjected to SDS-PAGE/Western blot analysis using the indicated antibodies. (C) Upregulation of CHOP by NASTRp. NCI-H441 cells were treated with 20 NASTRp for 0–24 hours. At the indicated time points, cells were fixed & subjected to immunofluorescence assay with anti-CHOP antibody. Cell images were microphotographed using fluorescence microscopy at 60X magnification. Scale bars: white; 10 μ m. (D) TUNEL-positive cells in NASTRp-treated (for 48 hours) A549 cells were counted & analyzed as relative % of TUNEL-positive cells. *P < 0.05 versus vehicle. Image collected & cropped by CiteAb from the following publication (<https://dx.plos.org/10.1371/journal.pone.0122628>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Immunocytochemistry/ Immunofluorescence: GADD153/CHOP Antibody - BSA Free [NBP2-13172] - NASTRp induces ER stress & eventually leads to cell death with Bim up-regulation. (A) qRT-PCR analysis of biomarkers of ER stress/UPR related genes in NCI-H441 treated with 20 μ M NASTRp in time course manner. (B) NASTRp induced ER stress & activated UPR led to apoptosis with Bim induction in NSCLC cell lines. Cells were treated with the indicated concentrations of NASTRp for 24 hours & cell lysates were subjected to SDS-PAGE/Western blot analysis using the indicated antibodies. (C) Upregulation of CHOP by NASTRp. NCI-H441 cells were treated with 20 NASTRp for 0–24 hours. At the indicated time points, cells were fixed & subjected to immunofluorescence assay with anti-CHOP antibody. Cell images were microphotographed using fluorescence microscopy at 60X magnification. Scale bars: white; 10 μ m. (D) TUNEL-positive cells in NASTRp-treated (for 48 hours) A549 cells were counted & analyzed as relative % of TUNEL-positive cells. *P < 0.05 versus vehicle. Image collected & cropped by CiteAb from the following publication (<https://dx.plos.org/10.1371/journal.pone.0122628>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.

C



Publications

Torrens JN, Hetzer SM, Evanson NK. Brief Oxygen Exposure after Traumatic Brain Injury Hastens Recovery and Promotes Adaptive Chronic Endoplasmic Reticulum Stress Responses International Journal of Molecular Sciences 2023-06-06 [PMID: 37372978] (Western Blot, Mouse)

Fielder E, Wan T, Alimohammadiha G et al. Short senolytic or senostatic interventions rescue progression of radiation-induced frailty and premature ageing in mice eLife 2022-05-04 [PMID: 35507395] (Western Blot, Mouse)

Shastri S, Shinde T, Perera AP et al. Idebenone Protects against Spontaneous Chronic Murine Colitis by Alleviating Endoplasmic Reticulum Stress and Inflammatory Response Biomedicines 2020-09-28 [PMID: 32998266] (Western Blot, Mouse)

Shastri S, Shinde T, Woolley KL et al. Short-Chain Naphthoquinone Protects Against Both Acute and Spontaneous Chronic Murine Colitis by Alleviating Inflammatory Responses Frontiers in Pharmacology 2021-08-23 [PMID: 34497514] (Western Blot, Mouse)

Criado-Marrero M, Blazier DM, Gould LA et al. Evidence against a contribution of the CCAAT-enhancer binding protein homologous protein (CHOP) in mediating neurotoxicity in rTg4510 mice Scientific Reports 2022-05-05 [PMID: 35513476] (Western Blot, Mouse)

Weaver FE, White E, Peek AM et al. 4-Phenylbutyric acid mitigates ER stress-induced neurodegeneration in the spinal cords of a GM2 gangliosidosis mouse model Hum Mol Genet 2024-11-12 [PMID: 39530163]

Brueck L, Roocke S, Matschke V et al. FGF23 and Cell Stress in SaOS-2 Cells-A Model Reflecting X-Linked Hypophosphatemia Dynamics Cells 2024-09-10 [PMID: 39329699]

Salvador-Barbero B, Alvarez-Fernandez M, Zapatero-Solana E et al. CDK4/6 Inhibitors Impair Recovery from Cytotoxic Chemotherapy in Pancreatic Adenocarcinoma Cancer Cell 2020-03-16 [PMID: 32109375]

Espina M, Di Franco N, Brañas-Navarro M et al. The GRP78-PERK axis contributes to memory and synaptic impairments in Huntington's disease R6/1 mice Neurobiology of disease 2023-07-11 [PMID: 37442396] (WB, Mouse)

Details:

Dilution: 1:1000

Toyama T, Kudryashova TV, Ichihara A et al. GATA6 coordinates cross-talk between BMP10 and oxidative stress axis in pulmonary arterial hypertension Scientific reports 2023-04-22 [PMID: 37087509] (WB, Mouse)

Sozen E, Demirel-Yalciner T, Sari D, Ozer NK Cholesterol accumulation in hepatocytes mediates IRE1/p38 branch of endoplasmic reticulum stress to promote nonalcoholic steatohepatitis Free radical biology & medicine 2022-08-19 [PMID: 35995397] (ICC/IF, Mouse)

Details:

AML12 cells (mouse hepatocytes)

Ou C, Xie W, Jiang P et al. Lycium barbarum L. and Salvia miltiorrhiza Bunge protect retinal pigment epithelial cells through endoplasmic reticulum stress Journal of ethnopharmacology 2022-10-05 [PMID: 35792279]

More publications at <http://www.novusbio.com/NBP2-13172>

Procedures

Western Blot Protocol specific for CHOP/GADD153 antibody (NBP2-13172)

Western Blot Protocol

1. Perform SDS-PAGE on samples to be analyzed, loading 40 ug of total protein per lane.
 2. Transfer proteins to membrane according to the instructions provided by the manufacturer of the membrane and transfer apparatus.
 3. Stain according to standard Ponceau S procedure (or similar product) to assess transfer success, and mark molecular weight standards where appropriate.
 4. Rinse the blot.
 5. Block the membrane using standard blocking buffer for at least 1 hour.
 6. Wash the membrane in wash buffer three times for 10 minutes each.
 7. Dilute primary antibody in blocking buffer and incubate 1 hour at room temperature.
 8. Wash the membrane in wash buffer three times for 10 minutes each.
 9. Apply the diluted HRP conjugated secondary antibody in blocking buffer (as per manufacturers instructions) and incubate 1 hour at room temperature.
 10. Wash the blot in wash buffer three times for 10 minutes each (this step can be repeated as required to reduce background).
 11. Apply the detection reagent of choice in accordance with the manufacturers instructions.
- Note: Tween-20 can be added to the blocking or antibody dilution buffer at a final concentration of 0.05-0.2%.

*The above information is only intended as a guide. The researcher should determine what protocol best meets their needs. Please follow safe laboratory procedures.

Immunohistochemistry-Paraffin Embedded Sections Protocol specific for CHOP/GADD153 antibody (NBP2-13172)

Immunohistochemistry-Paraffin Embedded Sections Protocol

Antigen Unmasking:

Bring slides to a boil in 10 mM sodium citrate buffer (pH 6.0) then maintain at a sub-boiling temperature for 10 minutes. Cool slides on bench-top for 30 minutes.

Staining:

1. Wash sections in deionized water three times for 5 minutes each.
2. Wash sections in wash buffer for 5 minutes.
3. Block each section with 100-400 ul blocking solution for 1 hour at room temperature.
4. Remove blocking solution and add 100-400 ul diluted primary antibody. Incubate overnight at 4 C.
5. Remove antibody solution and wash sections in wash buffer three times for 5 minutes each.
6. Add 100-400 ul biotinylated diluted secondary antibody. Incubate 30 minutes at room temperature.
7. Remove secondary antibody solution and wash sections three times with wash buffer for 5 minutes each.
8. Add 100-400 ul Streptavidin-HRP reagent to each section and incubate for 30 minutes at room temperature.
9. Wash sections three times in wash buffer for 5 minutes each.
10. Add 100-400 ul DAB substrate to each section and monitor staining closely.
11. As soon as the sections develop, immerse slides in deionized water.
12. Counterstain sections in hematoxylin.
13. Wash sections in deionized water two times for 5 minutes each.
14. Dehydrate sections.
15. Mount coverslips.

*The above information is only intended as a guide. The researcher should determine what protocol best meets their needs. Please follow safe laboratory procedures.

Immunocytochemistry/Immunofluorescence Protocol for GADD153/CHOP Antibody (NBP2-13172)

Immunocytochemistry Protocol

Culture cells to appropriate density in 35 mm culture dishes or 6-well plates.

1. Remove culture medium and wash the cells briefly in PBS. Add 10% formalin to the dish and fix at room temperature for 10 minutes.
2. Remove the formalin and wash the cells in PBS.
3. Permeabilize the cells with 0.1% Triton X100 or other suitable detergent for 10 min.
4. Remove the permeabilization buffer and wash three times for 10 minutes each in PBS. Be sure to not let the specimen dry out.
5. To block nonspecific antibody binding, incubate in 10% normal goat serum from 1 hour to overnight at room temperature.
6. Add primary antibody at appropriate dilution and incubate overnight at 4C.
7. Remove primary antibody and replace with PBS. Wash three times for 10 minutes each.
8. Add secondary antibody at appropriate dilution. Incubate for 1 hour at room temperature.
9. Remove secondary antibody and replace with PBS. Wash three times for 10 minutes each.
10. Counter stain DNA with DAPI if required.



Flow (Intracellular) Protocol for GADD153/CHOP Antibody (NBP2-13172)

Protocol for Flow Cytometry Intracellular Staining

Sample Preparation.

1. Grow cells to 60-85% confluency. Flow cytometry requires between 2×10^5 and 1×10^6 cells for optimal performance.
2. If cells are adherent, harvest gently by washing once with staining buffer and then scraping. Avoid using trypsin as this can disrupt certain epitopes of interest. If enzymatic harvest is required, use Accutase, Collagenase, or TrypLE Express for a less damaging option.
3. Reserve 100 μ L for counting, then transfer cell volume into a 50 mL conical tube and centrifuge for 8 minutes at 400 RCF.
 - a. Count cells using a hemocytometer and a 1:1 trypan blue exclusion stain to determine cell viability before starting the flow protocol. If cells appear blue, do not proceed.
4. Re-suspend cells to a concentration of 1×10^6 cells/mL in staining buffer (NBP2-26247).
5. Aliquot out 100 μ L samples in accordance with your experimental samples.

Tip: When cell surface and intracellular staining are required in the same sample, it is advisable that the cell surface staining be performed first since the fixation and permeabilization steps might reduce the availability of surface antigens.

Intracellular Staining.

Tip: When performing intracellular staining, it is important to use appropriate fixation and permeabilization reagents based upon the target and its subcellular location. Generally, our Intracellular Flow Assay Kit (NBP2-29450) is a good place to start as it contains an optimized combination of reagents for intracellular staining as well as an inhibitor of intracellular protein transport (necessary if staining secreted proteins). Certain targets may require more gentle or transient permeabilization protocols such as the commonly employed methanol or saponin-based methods.

Protocol for Cytoplasmic Targets:

1. Fix the cells by adding 100 μ L fixation solution (such as 4% PFA) to each sample for 10-15 minutes.
2. Permeabilize cells by adding 100 μ L of a permeabilization buffer to every 1×10^6 cells present in the sample. Mix well and incubate at room temperature for 15 minutes.
 - a. For cytoplasmic targets, use a gentle permeabilization solution such as 1X PBS + 0.5% Saponin or 1X PBS + 0.5% Tween-20.
 - b. To maintain the permeabilized state throughout your experiment, use staining buffer + 0.1% of the permeabilization reagent (i.e. 0.1% Tween-20 or 0.1% Saponin).
3. Following the 15 minute incubation, add 2 mL of the staining buffer + 0.1% permeabilizer to each sample.
4. Centrifuge for 1 minute at 400 RCF.
5. Discard supernatant and re-suspend in 100 μ L of staining buffer + 0.1% permeabilizer.
6. Add appropriate amount of each antibody (eg. 1 test or 1 μ g per sample, as experimentally determined).
7. Mix well and incubate at room temperature for 30 minutes- 1 hour. Gently mix samples every 10-15 minutes.
8. Following the primary/conjugate incubation, add 1-2 mL/sample of staining buffer +0.1% permeabilizer and centrifuge for 1 minute at 400 RCF.
9. Wash twice by re-suspending cells in staining buffer (2 mL for tubes or 200 μ L for wells) and centrifuging at 400 RCF for 5 minutes. Discard supernatant.
10. Add appropriate amount of secondary antibody (as experimentally determined) to each sample.
11. Incubate at room temperature in dark for 20 minutes.
12. Add 1-2 mL of staining buffer and centrifuge at 400 RCF for 1 minute and discard supernatant.
13. Wash twice by re-suspending cells in staining buffer (2 mL for tubes or 200 μ L for wells) and centrifuging at 400 RCF for 5 minutes. Discard supernatant.
14. Resuspend in an appropriate volume of staining buffer (usually 500 μ L per sample) and proceed with analysis on your flow cytometer.



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP2-13172

NB800-PC1	HeLa Whole Cell Lysate
NBP2-13172PEP	GADD153/CHOP Antibody Blocking Peptide
NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-13172

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications



