

Product Datasheet

APP Antibody - BSA Free

NBP2-15575

Unit Size: 0.1 ml

Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Reviews: 1 Publications: 2

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:

www.novusbio.com/NBP2-15575

Updated 9/9/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-15575



NBP2-15575

APP Antibody - BSA Free

Product Information

Unit Size	0.1 ml
Concentration	Concentrations vary lot to lot. See vial label for concentration. If unlisted please contact technical services.
Storage	Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.01% Thimerosal
Isotype	IgG
Purity	Antigen Affinity-purified
Buffer	0.1M Tris (pH 7), 0.1M Glycine, 20% Glycerol
Target Molecular Weight	87 kDa

Product Description


Description	Novus Biologicals Rabbit APP Antibody - BSA Free (NBP2-15575) is a polyclonal antibody validated for use in IHC, WB, ELISA and ICC/IF. Anti-APP Antibody: Cited in 2 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	351
Gene Symbol	APP
Species	Human, Mouse, Rat
Reactivity Notes	Xenopus laevis (83%).
Immunogen	Recombinant protein encompassing a sequence within the N-terminus region of human APP. The exact sequence is proprietary.

Product Application Details


Applications	Western Blot, Immunohistochemistry-Paraffin, ELISA, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry
Recommended Dilutions	Western Blot 1:500-1:3000, ELISA 1:1000-1:10000, Immunohistochemistry 1:100-1:1000, Immunocytochemistry/ Immunofluorescence 1:100-1:1000, Immunohistochemistry-Paraffin 1:100-1:1000

Images


Western Blot: APP Antibody [NBP2-15575] - Rat tissue extract (50 ug) was separated by 7.5% SDS-PAGE, and the membrane was blotted with APP antibody diluted at 1:500.




Immunocytochemistry/Immunofluorescence: APP Antibody [NBP2-15575] - SH-SY5Y cells were fixed in 4% paraformaldehyde at RT for 15 min. Green: APP protein stained by APP antibody diluted at 1:100. Red: beta Tubulin 3/ TUJ1 protein stained by beta Tubulin 3/ TUJ1 antibody diluted at 1:200. Blue: Hoechst 33342 staining.




Immunohistochemistry-Paraffin: APP Antibody [NBP2-15575] - Rat brain. APP antibody diluted at 1:500. Antigen Retrieval: Citrate buffer, pH 6.0, 15 min




Western Blot: APP Antibody [NBP2-15575] - Various whole cell extracts (30 ug) were separated by 7.5% SDS-PAGE, and the membrane was blotted with APP antibody diluted at 1:500.




Western Blot: APP Antibody [NBP2-15575] - Human tissue extract (30 ug) was separated by 7.5% SDS-PAGE, and the membrane was blotted with APP antibody diluted at 1:500. Human tissue extract (30 ug) was separated by 7.5% SDS-PAGE, and the membrane was blotted with APP antibody diluted at 1:500.




Western Blot: APP Antibody [NBP2-15575] - Mouse tissue extract (50 ug) was separated by 7.5% SDS-PAGE, and the membrane was blotted with APP antibody diluted at 1:500.



Immunocytochemistry/Immunofluorescence: APP Antibody [NBP2-15575] - U-87 MG cells were fixed in 4% paraformaldehyde at RT for 15 min. Green: APP protein stained by APP antibody diluted at 1:100. Blue: Hoechst 33342 staining.



ELISA: APP Antibody [NBP2-15575] - Detection of non-transfected and transfected 293T whole cell extracts using NBP2-15575 as capture antibody at concentration of 5 ug/mL and a mouse monoclonal anti-APP antibody as detection antibody at concentration of 1 ug/mL. Mouse IgG antibody (HRP) (NBP2-19382) was diluted at 1:10000 and used to detect the primary antibody.



Publications

Zaidi Y, Corker A, Vasileva VY Et al. Chronic Porphyromonas gingivalis lipopolysaccharide induces adverse myocardial infarction wound healing through activation of CD8+ T cells American journal of physiology. Heart and circulatory physiology 2021-11-01 [PMID: 34597184] (WB, Mouse)

Choi JY, Yeo IJ, Kim KC et al. K284-6111 prevents the amyloid beta-induced neuroinflammation and impairment of recognition memory through inhibition of NF- κ B-mediated CHI3L1 expression J Neuroinflammation 2018-08-11 [PMID: 30098604] (WB, Mouse)



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP2-15575

HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control
NBP1-90246PEP	APP Recombinant Protein Antigen

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-15575

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications