

Product Datasheet

DCK Antibody - BSA Free NBP2-16108

Unit Size: 0.1 ml

Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 1

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-16108

Updated 9/25/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-16108



NBP2-16108

DCK Antibody - BSA Free


Product Information	
Unit Size	0.1 ml
Concentration	Concentrations vary lot to lot. See vial label for concentration. If unlisted please contact technical services.
Storage	Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.025% Proclin 300
Isotype	IgG
Purity	Antigen Affinity-purified
Buffer	PBS, 20% Glycerol
Target Molecular Weight	31 kDa

Product Description	
Description	Novus Biologicals Rabbit DCK Antibody - BSA Free (NBP2-16108) is a polyclonal antibody validated for use in IHC, WB, ICC/IF and IP. Anti-DCK Antibody: Cited in 1 publication. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	1633
Gene Symbol	DCK
Species	Human, Mouse, Rat
Reactivity Notes	Chicken (82%).
Immunogen	Recombinant protein encompassing a sequence within the center region of human DCK. The exact sequence is proprietary.


Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunoprecipitation
Recommended Dilutions	Western Blot 1:1000-1:20000, Immunohistochemistry 1:100-1:1000, Immunocytochemistry/ Immunofluorescence 1:100-1:1000, Immunoprecipitation 1:100-1:500, Immunohistochemistry-Paraffin 1:100-1:1000

Images


Western Blot: DCK Antibody [NBP2-16108] - Non-transfected (-) and transfected (+) 293T whole cell extracts (30 ug) were separated by 12% SDS-PAGE, and the membrane was blotted with DCK antibody. HRP-conjugated anti-rabbit IgG antibody (NBP2-19301) was used to detect the primary antibody.




Western Blot: DCK Antibody [NBP2-16108] - Non-transfected (-) and transfected (+) 293T whole cell extracts (30 ug) were separated by 12% SDS-PAGE, and the membrane was blotted with DCK antibody diluted at 1:5000. HRP-conjugated anti-rabbit IgG antibody was used to detect the primary antibody.




Immunohistochemistry-Paraffin: DCK Antibody [NBP2-16108] - Rat lung. DCK stained by DCK antibody diluted at 1:500. Antigen Retrieval: Citrate buffer, pH 6.0, 15 min.




Western Blot: DCK Antibody [NBP2-16108] - Various whole cell extracts (30 ug) were separated by 12% SDS-PAGE, and the membrane was blotted with DCK antibody diluted at a dilution of 1:5000. The HRP-conjugated anti-rabbit IgG antibody (NBP2-19301) was used to detect the primary antibody.




Western Blot: DCK Antibody [NBP2-16108] - A. 30 ug PC-12 whole cell lysate/extract B. 30 ug Rat2 whole cell lysate/extract 12% SDS-PAGE DCK antibody dilution: 1:1000 The HRP-conjugated anti-rabbit IgG antibody (NBP2-19301) was used to detect the primary antibody.




Immunohistochemistry-Paraffin: DCK Antibody [NBP2-16108] - Mouse hind brain. DCK antibody dilution: 1:500. Antigen Retrieval: Trilogy™ (EDTA based, pH 8.0) buffer, 15min.



Immunoprecipitation: DCK Antibody [NBP2-16108] - Immunoprecipitation of DCK protein from 293T whole cell extracts using 5 ug of DCK antibody. Western blot analysis was performed using DCK antibody. EasyBlot anti-Rabbit IgG was used as a secondary reagent.



Western Blot: DCK Antibody [NBP2-16108] - Various whole cell extracts (30 ug) were separated by 12% SDS-PAGE, and the membrane was blotted with DCK antibody (NBP2-16108) diluted at 1:3000. The HRP-conjugated anti-rabbit IgG antibody was used to detect the primary antibody.




DCK antibody detects DCK protein at nucleus by immunofluorescent analysis.

Sample: HeLa cells were fixed in 4% paraformaldehyde at RT for 15 min.

Green: DCK protein stained by DCK antibody (NBP2-16108) diluted at 1:500.

Red: Phalloidin, a cytoskeleton marker, diluted at 1:100.

Scale bar = 10 um.



Publications

Chen YH, Chen YC, Lin CC et al. Synergistic Anticancer Effects of Gemcitabine with Pitavastatin on Pancreatic Cancer Cell Line MIA PaCa-2 in vitro and in vivo Cancer Manag Res 2020-06-17 [PMID: 32606957]



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP2-16108

NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-16108

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

