

Product Datasheet

RAP1A Antibody (1D2-1C64)

NBP2-22526

Unit Size: 100 ug

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-22526

Updated 9/9/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-22526



NBP2-22526**RAP1A Antibody (1D2-1C64)**


Product Information	
Unit Size	100 ug
Concentration	1 mg/ml
Storage	Store at -20C. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	1D2-1C64
Preservative	0.05% Sodium Azide
Isotype	IgG2a
Purity	Protein A purified
Buffer	PBS with 1 mg/ml BSA and 30% glycerol
Target Molecular Weight	21 kDa

Product Description	
Description	Novus Biologicals Mouse RAP1A Antibody (1D2-1C64) (NBP2-22526) is a monoclonal antibody validated for use in IHC, WB, ICC/IF and IP. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Mouse
Gene ID	5906
Gene Symbol	RAP1A
Species	Human, Mouse
Reactivity Notes	Please note that this antibody is reactive to Mouse and derived from the same host, Mouse. Additional Mouse on Mouse blocking steps may be required for IHC and ICC experiments. Please contact Technical Support for more information.
Specificity/Sensitivity	This product was made specifically to the RAP1A human protein (P62834). It has not yet been tested for cross reactivity to other related proteins.
Immunogen	Recombinant full-length human Rap1


Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunoprecipitation
Recommended Dilutions	Western Blot 1:500 - 1:1000, Immunohistochemistry 1:100 - 1:1000, Immunocytochemistry/ Immunofluorescence 1:10 - 1:200, Immunoprecipitation 2 ug, Immunohistochemistry-Paraffin 1:100 - 1:1000

Images


Western Blot: RAP1A Antibody (1D2-1C64) [NBP2-22526] - Analysis of 25 ug of various whole cell lysates per well.




Immunocytochemistry/Immunofluorescence: RAP1A Antibody (1D2-1C64) [NBP2-22526] - Analysis of Rap1 (green) showing staining in the cytoplasm of C2C12 cells (right) compared to a negative control without primary antibody (left).




Immunohistochemistry: RAP1A Antibody (1D2-1C64) [NBP2-22526] - Analysis of Rap1 showing staining in the membrane of human kidney tissue (right) compared with a negative control without primary antibody (left).



Immunocytochemistry/Immunofluorescence: RAP1A Antibody (1D2-1C64) [NBP2-22526] - Analysis of Rap1 (green) in HeLa cells. Formalin fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 10 minutes at room temperature and blocked with 1% BSA/TBST for 15 minutes at room temperature. Cells were probed with a Rap1 monoclonal antibody at a dilution of 1:100 for at least 1 hour at room temperature, washed with PBS, and incubated with DyLight 488-conjugated goat anti-mouse IgG secondary antibody at a dilution of 1:400 for 30 minutes at room temperature. Nuclei (blue) were stained with Hoechst 33342 dye.



Immunoprecipitation: RAP1A Antibody (1D2-1C64) [NBP2-22526] - Analysis of Rap1 was performed on NIH3T3 cells. Antigen-antibody complexes were formed by incubating 750ug of NIH3T3 cell lysate with 2ug of a Rap1 monoclonal antibody overnight on a rocking platform at 4C. The immune complexes were captured on 50ul Protein A/G Plus Agarose, washed extensively, and eluted with 5X Lane Marker Reducing Sample Buffer. Eluted sample (right lane) and 25ug of NIH3T3 control lysate (left lane) were resolved on a 4-20% Tris-HCl polyacrylamide gel, transferred to a PVDF membrane, and blocked with 5% BSA/TBS-0.1%Tween for at least 1 hour. The membrane was probed with a Rap1 monoclonal antibody at a dilution of 1:500 overnight rotating at 4C, washed in TBST, and probed with Clean-blot IP Detection Reagent at a dilution of 1:2500 for at least 1 hour. Chemiluminescent detection was performed using SuperSignal West Dura.





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP2-22526

NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB7539	Goat anti-Mouse IgG (H+L) Secondary Antibody [HRP]
NBP1-96778	Mouse IgG2a Isotype Control (M2A)

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-22526

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

