

Product Datasheet

c-Maf Antibody - BSA Free NBP2-24551

Unit Size: 0.1 mg

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Reviews: 3 Publications: 2

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-24551

Updated 9/9/2025 v.20.1

**Earn rewards for product
reviews and publications.**

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-24551



NBP2-24551

c-Maf Antibody - BSA Free


Product Information	
Unit Size	0.1 mg
Concentration	1.0 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.02% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	PBS

Product Description	
Description	Novus Biologicals Rabbit c-Maf Antibody - BSA Free (NBP2-24551) is a polyclonal antibody validated for use in IHC and WB. Anti-c-Maf Antibody: Cited in 2 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	4094
Gene Symbol	MAF
Species	Human, Mouse, Rat, Bovine
Reactivity Notes	Chimpanzee (87%), Bovine, Equine (81%).
Specificity/Sensitivity	100% homologous in human isoforms CRA_a, CRA_b, and CRA_c).
Immunogen	A synthetic peptide corresponding to amino acids 75-110 of human c-maf was used as the immunogen, GenBank no. NP_001026974.1 .


Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, Immunohistochemistry
Recommended Dilutions	Western Blot 2-4 ug/ml, Immunohistochemistry, Immunohistochemistry-Paraffin 1-2 ug/ml

Images


Western Blot: c-Maf Antibody [NBP2-24551] - Total protein from Human Pancreas and Human Small Intestine was separated on a 7.5% gel by SDS-PAGE, transferred to PVDF membrane and blocked in 5% non-fat milk in TBST. The membrane was probed with 2.0 ug/ml anti-cMaf in 1% block buffer and detected with an anti-rabbit HRP secondary antibody using chemiluminescence.




Western Blot: c-Maf Antibody [NBP2-24551] - SUM159 and HCC1954 whole cell lysates were loaded with 50 ug/lane. 10% SDS-PAGE. c-Maf Antibody (NBP2-24551) was used for primary antibody: 1:500, 4C, overnight. WB image submitted by a verified customer review.



Immunohistochemistry-Paraffin: c-Maf Antibody [NBP2-24551] - IHC analysis of a formalin fixed paraffin embedded (FFPE) tissue section of human liver cancer using c-MAF antibody at 2ug/ml concentration (1:500 dilution). The primary antibody binding to c-MAF protein was detected using HRP conjugated anti-rabbit secondary antibody with DAB reagent, and the sections were further counterstained with hematoxylin for labeling cellular nuclei. This c-MAF antibody showed a very strong nuclear and moderate to strong staining in the endothelial cells (blood vessels) and in a subset of apparently infiltrating inflammatory cells. Weak to moderate cytoplasmic and nuclear staining was observed in liver cancer cells and the cells of tumor stroma including fibroblasts.




Western Blot: c-Maf Antibody [NBP2-24551] - Analysis of Lysate from human brain in the 1) absence, 2) presence of immunizing peptide, 3) mouse brain and 4) rat brain probed at 3 ug/ml.



Immunohistochemistry-Paraffin: c-Maf Antibody [NBP2-24551] - Staining of Human kidney probed with c-maf antibody at 10 ug/ml.



Immunohistochemistry-Paraffin: c-Maf Antibody [NBP2-24551] - Staining of Human placenta probed with c-maf antibody at 10 ug/ml.



Publications

Aoki H, Ogino H, Tomita H et al. Disruption of Rest Leads to the Early Onset of Cataracts with the Aberrant Terminal Differentiation of Lens Fiber Cells. PLoS ONE. 2016-09-16 [PMID: 27631609] (IF/IHC, Mouse)

Hansen A, Henderson S, Lagos D et al. KSHV-encoded miRNAs target MAF to induce endothelial cell reprogramming. Genes Dev. 2010-01-15 [PMID: 20080955] (WB, Human)

Details:

WB (Primary human lymphatic epithelial cells): Figs 3C, 4C, 6B.

Procedures

Western Blot Protocol for c-Maf Antibody (NBP2-24551)

Western Blot Protocol

1. Perform SDS-PAGE on samples to be analyzed, loading 10-25 ug of total protein per lane.
2. Transfer proteins to PVDF membrane according to the instructions provided by the manufacturer of the membrane and transfer apparatus.
3. Stain the membrane with Ponceau S (or similar product) to assess transfer success, and mark molecular weight standards where appropriate.
4. Rinse the blot TBS -0.05% Tween 20 (TBST).
5. Block the membrane in 5% Non-fat milk in TBST (blocking buffer) for at least 1 hour.
6. Wash the membrane in TBST three times for 10 minutes each.
7. Dilute primary antibody in blocking buffer and incubate overnight at 4C with gentle rocking.
8. Wash the membrane in TBST three times for 10 minutes each.
9. Incubate the membrane in diluted HRP conjugated secondary antibody in blocking buffer (as per manufacturer's instructions) for 1 hour at room temperature.
10. Wash the blot in TBST three times for 10 minutes each (this step can be repeated as required to reduce background).
11. Apply the detection reagent of choice in accordance with the manufacturer's instructions.





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP2-24551

NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-24551

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

