

Product Datasheet

TLR4 Antibody (MTS510) - Azide and BSA Free NBP2-24865

Unit Size: 0.1 mg

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 13

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-24865

Updated 9/9/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-24865



NBP2-24865

TLR4 Antibody (MTS510) - Azide and BSA Free

Product Information	
Unit Size	0.1 mg
Concentration	1.0 mg/ml
Storage	Store at -20C. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	MTS510
Preservative	No Preservative
Isotype	IgG2a Kappa
Purity	Protein G purified
Buffer	PBS
Target Molecular Weight	95.7 kDa


Product Description	
Description	Novus Biologicals Rat TLR4 Antibody (MTS510) - Azide and BSA Free (NB100-56560) is a monoclonal antibody validated for use in WB, Flow and ICC/IF. Anti-TLR4 Antibody: Cited in 12 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rat
Gene ID	7099
Gene Symbol	TLR4
Species	Human, Mouse
Specificity/Sensitivity	This antibody preferentially recognizes TLR4-MD-2 complex than of TLR4 alone.
Immunogen	This TLR4 Antibody (MTS510) was developed by immunizing rats with the Ba/F3 cell line expressing mouse TLR4 and MD-2 (Akashi et al, 2000).

Product Application Details	
Applications	Western Blot, Flow Cytometry, Flow (Cell Surface), Immunocytochemistry/Immunofluorescence, Block/Neutralize, CyTOF-ready
Recommended Dilutions	Western Blot reported in scientific literature (PMID 23629653), Flow Cytometry 1ul/1 million cells, Immunocytochemistry/ Immunofluorescence reported in scientific literature (PMID 22573811), Flow (Cell Surface) reported in scientific literature (PMID 15879150), CyTOF-ready, Block/Neutralize
Application Notes	In FA, the antibody blocks activation of monocytes with LPS (Akashi et al. 2000). The optimal condition has to be determined for individual experiments.




Images


Flow Cytometry: TLR4 Antibody (MTS510) - Azide Free [NBP2-24865] - Analysis using Azide/BSA FREE version of NBP2-24865. Cell surface analysis of TLR4 on mouse RAW cells using 0.5 ug/10⁶ cells of TLR4 antibody, anti-rat APC conjugated secondary this antibody and TLR cell surface flow kit this antibody (green represents isotype control; red represents TLR4 antibody).




Flow Cytometry: TLR4 Antibody (MTS510) - Azide Free [NBP2-24865] - Cell surface analysis of TLR4 in mouse peritoneal cells using this antibody. The green histogram represents the isotype control and red represents anti-TLR4 antibody.




Flow Cytometry: TLR4 Antibody (MTS510) - Azide Free [NBP2-24865] - Analysis using the FITC conjugate of NBP2-24865. Staining of TLR4 in mouse peritoneal cells using 1 ug of this antibody. Shaded histogram represents cells without antibody; green represents isotype control; red represents anti-TLR4 antibody.




Flow Cytometry: TLR4 Antibody (MTS510) - Azide Free [NBP2-24865] - Analysis using the FITC conjugate of NBP2-24865. Staining of TLR4 in mouse peritoneal cells using 1 ug of this antibody. Shaded histogram represents cells without antibody; green represents isotype control; red represents anti-TLR4 antibody.




Flow Cytometry: TLR4 Antibody (MTS510) - Azide Free [NBP2-24865] - Analysis using the FITC conjugate of NBP2-24865. Staining of TLR4 in RAW cells using 0.5 ug of this antibody. Black histogram represents cells without antibody; green represents isotype control; red represents anti-TLR4 antibody.



Flow (Cell Surface): TLR4 Antibody (MTS510) - Azide Free [NBP2-24865] - Analysis using the PE conjugate of NBP2-24865. Staining of cell surface TLR4 using this antibody at 0.625 ug/10⁶ Raw cells (pretreated with anti-CD16/32). The shaded histogram represents the isotype control.



Flow Cytometry: TLR4 Antibody (MTS510) - Azide Free [NBP2-24865] - Analysis using the PE conjugate of NBP2-24865. Staining of TLR4 in mouse peritoneal cells using 0.5 ug of this antibody. Shaded histogram represents cells without antibody; green represents isotype control; red represents anti-TLR4 antibody.



Publications

Jung DY, Lee H, Jung BY et al. TLR4, but not TLR2, signals autoregulatory apoptosis of cultured microglia: a critical role of IFN-beta as a decision maker J Immunol 2005-05-10 [PMID: 15879150]

Jung DY, Lee H, Jung BY et al. TLR4, but not TLR2, signals autoregulatory apoptosis of cultured microglia: a critical role of IFN-beta as a decision maker J Immunol (Flow-CS)

Details:

This publication used the FITC conjugated form of this antibody (Cat# NBP2-24450) and the PE conjugated form of this antibody (Cat# NBP2-24741).

Tsukamoto H, Fukudome K, Takao S et al. Lipopolysaccharide-binding protein-mediated Toll-like receptor 4 dimerization enables rapid signal transduction against lipopolysaccharide stimulation on membrane-associated CD14-expressing cells. Int Immunol. 2010-04-01 [PMID: 20133493] (WB)

Details:

TLR4 (IMG-578): WB, TLR4 stably transfected Ba/F3 cells. Note: These cells were immunoprecipitated with a TLR4 mAb (clone UT41) then western blotted with the IMG-578A TLR4 pAb Fig 3B. 3. TLR4 (IMG-428A): Flow (cell surface), RAW264 and TLR4 stably transfected Ba/F3 cells. Figs 1B, 1E.

Schaefer L, Babelova A, Kiss E et al. The matrix component biglycan is proinflammatory and signals through Toll-like receptors 4 and 2 in macrophages J Clin Invest 2005-08-01 [PMID: 16025156]

Details:

This reference used the Azide Free version of NB100-56560.

Park HJ, Hong JH, Kwon HJ et al. TLR4-mediated activation of mouse macrophages by Korean mistletoe lectin-C (KML-C). Biochem Biophys Res Commun. 2010-06-04 [PMID: 20450885] (WB, Mouse)

Details:

The following antibodies were used: TLR4/CD284 (IMG-428E), TLR4/CD284 (IMG-577), and TLR4 (IMG-6307A). IMG-577 & IMG-6307A were used in Fig 1 (WB): Peritoneal primary mouse (BALB/C) macrophages stimulated with LPS and then treated with KML-C (Korean mistl

EI Shikh ME, El Sayed RM, Wu Y et al. TLR4 on follicular dendritic cells: an activation pathway that promotes accessory activity. J Immunol. 2007-10-01 [PMID: 17878340]

Details:

Products cited: 1. TLR4-FITC (IMG-428C): Flow (Cell Surface), Fig 1C, D [primary follicular dendritic (FDC) and B cells isolated from BALB/c mouse splenic leukocytes]. FDC cells expressed TLR4, but the B cells did not. 2. TLR4 (IMG-579A): IHC (F), Fig 1A

Tsai CC, Lin CR, Tsai HY et al. The immunologically active oligosaccharides isolated from wheatgrass modulate monocytes via Toll-like receptor-2 signaling J Biol Chem 2013-05-01 [PMID: 23629653] (WB)

Ma CY, Shi GY, Shi CS et al. Monocytic thrombomodulin triggers LPS- and gram-negative bacteria-induced inflammatory response. J Immunol. 2012-06-15 [PMID: 22573811] (ICC/IF, Mouse)

Details:

TLR4 (IMG-428A). IF: mouse peritoneal macrophages (Fig 5D).

Corbucci C, Cenci E, Skrzypek F et al. Immune response to *Candida albicans* is preserved despite defect in O-mannosylation of secretory proteins. Med Mycol. 2007-12-01 [PMID: 17885949] (Flow-CS)

Details:

Products cited (mouse macrophages): 1. IMG-428C (TLR4): Flow (Cell Surface), Fig. 5A 2. IMG-428E (TLR4): FA, Fig. 5B.

Kim MY, Shu Y, Carsillo T et al. hsp70 and a novel axis of type I interferon-dependent antiviral immunity in the measles virus-infected brain. J Virol. 2013-01-01 [PMID: 23135720]

Macedo GC, Magnani DM, Carvalho NB et al. Central role of MyD88-dependent dendritic cell maturation and proinflammatory cytokine production to control *Brucella abortus* infection. J Immunol. 2008-01-15 [PMID: 18178848] (FLOW, Mouse)

Details:

IMG-428E: FA (mouse bone marrow derived and splenic dendritic cells), Fig. 7C,D.

Bruscia EM, Zhang PX, Satoh A et al. Abnormal trafficking and degradation of TLR4 underlie the elevated inflammatory response in cystic fibrosis. J Immunol. 2011-06-15 [PMID: 21593379]

Details:

1. TLR4/CD284 (IMG-428C) & TLR2/CD282 (IMG-6320C) : WB, IF/ICC, Flow (cell surface): Human macrophages with cystic fibrosis (CF) and wild type (WT) mouse macrophages, Fig 1 & S4. 2. TLR4/CD284 WB: CF and wild type macrophages treated with protein synthesis

More publications at <http://www.novusbio.com/NBP2-24865>



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP2-24865

NBP2-25295-1.0mg	LPS from E. Coli, TLR4 ligand
NBP2-26244	TLR4 Inhibitor Peptide Set
NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF005	Goat anti-Rat IgG Secondary Antibody [HRP]
NB7115	Goat anti-Rat IgG (H+L) Secondary Antibody [HRP]
NBP1-43321-0.5mg	Rat IgG2a Kappa Light Chain Isotype Control (R2a)

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-24865

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications



