

Product Datasheet

HMGB1/HMG-1 Antibody (1F3) - BSA Free NBP2-25148

Unit Size: 0.1 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 2

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-25148

Updated 9/9/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-25148




NBP2-25148**HMGB1/HMG-1 Antibody (1F3) - BSA Free**


Product Information	
Unit Size	0.1 ml
Concentration	1 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	1F3
Preservative	5mM Sodium Azide
Isotype	IgG2b
Purity	Immunogen affinity purified
Buffer	50% PBS, 50% glycerol
Target Molecular Weight	25 kDa
Product Description	
Description	Novus Biologicals Mouse HMGB1/HMG-1 Antibody (1F3) - BSA Free (NBP2-25148) is a monoclonal antibody validated for use in IHC, WB, Flow, ICC/IF and Simple Western. Anti-HMGB1/HMG-1 Antibody: Cited in 2 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Mouse
Gene ID	3146
Gene Symbol	HMGB1
Species	Human, Mouse, Rat
Immunogen	Full length recombinant human HMGB1 protein expressed in and purified from E. coli. [UniProt# P09429]
Product Application Details	
Applications	Western Blot, Simple Western, Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, CyTOF-ready
Recommended Dilutions	Western Blot 1:1000-1:2000, Simple Western 1:500, Flow Cytometry 1 ug per million cells, Immunohistochemistry 1:1000, Immunocytochemistry/Immunofluorescence 1:1000, CyTOF-ready
Application Notes	<p>This HMGB1 (1F3) antibody is useful for Immunocytochemistry/Immunofluorescence, Immunohistochemistry, and Western Blot, where a band is seen at approximately 25 kDa. This antibody is CyTOF ready.</p> <p>In Simple Western only 10 - 15 ul of the recommended dilution is used per data point.</p> <p>See Simple Western Antibody Database for Simple Western validation: Tested in Jurkat lysate 0.05 mg/mL, separated by Size, antibody dilution of 1:500, apparent MW was 36 kDa. Separated by Size-Wes, Sally Sue/Peggy Sue.</p>

Images

Western Blot: HMGB1/HMG-1 Antibody (1F3) [NBP2-25148] - Analysis of lysates from different cell lines probed with mouse mAb to high mobility group protein B1, (HMGB1), NBP2-25148, dilution 1:2,000: [1] protein standard, [2] NIH-3T3, [3] C6, [4] HEK293, [5] HeLa, and [6] SH-SY5Y. The 25kDa band revealed by MCA-1F3 antibody, corresponds to HMGB1 protein.




Immunocytochemistry/Immunofluorescence: HMGB1/HMG-1 Antibody (1F3) [NBP2-25148] - Analysis of HeLa cells stained with mouse mAb to HMGB1, NBP2-25148, dilution 1:2,000 in red and costained with rabbit pAb to GAPDH, dilution 1:2,000 in green. MCA-1F3 antibody stains the chromatin binding protein HMGB1, which is localized in the nuclei.



Flow Cytometry: HMGB1/HMG-1 Antibody (1F3) [NBP2-25148] - Intracellular flow cytometric staining of 1×10^6 MCF-7 cells using HMGB1 antibody (dark blue). Isotype control shown in orange. An antibody concentration of $1 \mu\text{g}/1 \times 10^6$ cells was used.



Immunocytochemistry/Immunofluorescence: HMGB1/HMG-1 Antibody (1F3) [NBP2-25148] - HeLa cells grown in tissue culture and stained with NBP2-25148 (red), chicken polyclonal antibody to Vimentin NB300-223 (green) and DNA (blue). NBP2-25148 reveals strong nuclear staining which overlaps with the DNA stain.



Simple Western: HMGB1/HMG-1 Antibody (1F3) [NBP2-25148] - Simple Western lane view shows a specific band for HMGB1 in 0.05 mg/ml of Jurkat lysate. This experiment was performed under reducing conditions using the 12-230 kDa separation system.



Publications

Tonner H, Hunn S, Auler N et al. A Monoclonal Anti-HMGB1 Antibody Attenuates Neurodegeneration in an Experimental Animal Model of Glaucoma International journal of molecular sciences 2022-04-07 [PMID: 35456925] (In Vivo, Rat)

Nestor J, Arinuma Y, Huerta TS et al. Lupus antibodies induce behavioral changes mediated by microglia and blocked by ACE inhibitors. J. Exp. Med. 2018-09-05 [PMID: 30185634] (ICC/IF, Mouse)



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP2-25148

NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB7539	Goat anti-Mouse IgG (H+L) Secondary Antibody [HRP]
NBP2-27231	Mouse IgG2b Isotype Control (MPC-11)

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-25148

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

