

Product Datasheet

CCDC124 Antibody - BSA Free NBP2-48858

Unit Size: 0.1 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 1

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-48858

Updated 12/2/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-48858



NBP2-48858

CCDC124 Antibody - BSA Free

Product Information	
Unit Size	0.1 ml
Concentration	Concentrations vary lot to lot. See vial label for concentration. If unlisted please contact technical services.
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.02% Sodium Azide
Isotype	IgG
Purity	Affinity purified
Buffer	PBS (pH 7.2) and 40% Glycerol


Product Description	
Description	Novus Biologicals Rabbit CCDC124 Antibody - BSA Free (NBP2-48858) is a polyclonal antibody validated for use in IHC, WB and ICC/IF. Anti-CCDC124 Antibody: Cited in 1 publication. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	115098
Gene Symbol	CCDC124
Species	Human
Reactivity Notes	Mouse (84%), Rat (82%)
Immunogen	This antibody was developed against a recombinant protein corresponding to amino acids: LEDAYWKDDDKHVMRKEQRKEEKEKRRLDQLERKKETQRLLEEEDSKLKGGK APRVATSSKVTRAQIEDTLRRDHQLRE

Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, Immunocytochemistry/Immunofluorescence, Immunohistochemistry
Recommended Dilutions	Western Blot Reported in scientific literature (PMID: 31044552), Immunohistochemistry 1:2500 - 1:5000, Immunocytochemistry/Immunofluorescence 0.25-2 ug/ml, Immunohistochemistry-Paraffin 1:2500 - 1:5000
Application Notes	For IHC-Paraffin, HIER pH 6 retrieval is recommended. ICC/IF, Fixation Permeabilization: Use PFA/Triton X-100.




Images


Immunocytochemistry/Immunofluorescence: CCDC124 Antibody [NBP2-48858] - Immunofluorescent staining of human cell line U-2 OS shows localization to plasma membrane & cytosol.




Immunohistochemistry-Paraffin: CCDC124 Antibody [NBP2-48858] - Staining of human hippocampus shows strong cytoplasmic positivity in neuronal cells.




Western Blot: CCDC124 Antibody [NBP2-48858] - The expression of IL-25 affects the cisplatin resistance of lung cancer cells. A, The expression of IL-25 in A549 & A549/CDDP cells. B, IL-25 overexpressed in A549 cells. C, silencing of IL-25 expression by siRNA in A549/CDDP cells. D, IL-25 overexpression desensitizes A549 cells to cisplatin treatment. E, Silencing of IL-25 expression sensitizes A549/CDDP cells to cisplatin treatment. *P < 0.01 vs indicated controls, all figures represent the average of three sets of independent experiments Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/31044552>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.




Western Blot: CCDC124 Antibody [NBP2-48858] - The expression of IL-25 affects the cisplatin resistance of lung cancer cells. A, The expression of IL-25 in A549 & A549/CDDP cells. B, IL-25 overexpressed in A549 cells. C, silencing of IL-25 expression by siRNA in A549/CDDP cells. D, IL-25 overexpression desensitizes A549 cells to cisplatin treatment. E, Silencing of IL-25 expression sensitizes A549/CDDP cells to cisplatin treatment. *P < 0.01 vs indicated controls, all figures represent the average of three sets of independent experiments Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/31044552>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.




Western Blot: CCDC124 Antibody [NBP2-48858] - IL-25 regulates the NF- κ B signaling pathway activity. A, The change of NF- κ B signaling pathway in IL-25 overexpressed A549 cells & IL-25 silenced A549/CDDP cells. B, The effect of exogenous IL-25 on the activity of NF- κ B signaling pathway in A549 cells. C, QNZ blocks the effect of exogenous IL-25 on the expression of major vault protein (MVP) & the activity of NF- κ B signaling pathway in A549 cells. *P < 0.01 vs indicated controls, all figures represent the average of three sets of independent experiments Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/31044552>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Western Blot: CCDC124 Antibody [NBP2-48858] - The expression of IL-25 affects the cisplatin resistance of lung cancer cells. A, The expression of IL-25 in A549 & A549/CDDP cells. B, IL-25 overexpressed in A549 cells. C, silence of IL-25 expression by siRNA in A549/CDDP cells. D, IL-25 overexpression desensitizes A549 cells to cisplatin treatment. E, Silence of IL-25 expression sensitizes A549/CDDP cells to cisplatin treatment. *P < 0.01 vs indicated controls, all figures represent the average of three sets of independent experiments Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/31044552>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Western Blot: CCDC124 Antibody [NBP2-48858] - The expression of IL-25 affects the cisplatin resistance of lung cancer cells. A, The expression of IL-25 in A549 & A549/CDDP cells. B, IL-25 overexpressed in A549 cells. C, silence of IL-25 expression by siRNA in A549/CDDP cells. D, IL-25 overexpression desensitizes A549 cells to cisplatin treatment. E, Silence of IL-25 expression sensitizes A549/CDDP cells to cisplatin treatment. *P < 0.01 vs indicated controls, all figures represent the average of three sets of independent experiments Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/31044552>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Publications

Shen W, Qiu Y, Li J, Wu C et al. IL-25 promotes cisplatin resistance of lung cancer cells by activating NF- κ B signaling pathway to increase of major vault protein Cancer Med 2019-05-01 [PMID: 31044552] (WB, Human)



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP2-48858

NBP2-48858PEP	CCDC124 Recombinant Protein Antigen
NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-48858

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

