

Product Datasheet

Parvalbumin Antibody (3C9) - BSA Free NBP2-50038

Unit Size: 0.1 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 3

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-50038

Updated 9/9/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-50038



NBP2-50038

Parvalbumin Antibody (3C9) - BSA Free

Product Information	
Unit Size	0.1 ml
Concentration	1 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	3C9
Preservative	5mM Sodium Azide
Isotype	IgG1
Purity	Immunogen affinity purified
Buffer	50% PBS, 50% glycerol
Target Molecular Weight	12 kDa


Product Description	
Description	Novus Biologicals Mouse Parvalbumin Antibody (3C9) - BSA Free (NBP2-50038) is a monoclonal antibody validated for use in IHC, WB and ICC/IF. Anti-Parvalbumin Antibody: Cited in 1 publication. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Mouse
Gene ID	5816
Gene Symbol	PVALB
Species	Human, Mouse, Rat, Porcine, Bovine, Equine
Immunogen	Full-length recombinant human Parvalbumin expressed in and purified from E. coli. [UniProt# P20472]

Product Application Details	
Applications	Western Blot, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry
Recommended Dilutions	Western Blot 1:1000 - 1:5000, Immunohistochemistry 1:1000 - 1:5000, Immunocytochemistry/ Immunofluorescence 1:1000 - 1:5000
Application Notes	In WB, this antibody recognizes a band at 12 kDa in skeletal muscle lysates, which represents Parvalbumin. Note, this antibody is not cross-reactive with either calbindin or calretinin despite their related amino acid sequences.




Images


Western Blot: Parvalbumin Antibody (3C9) [NBP2-50038] - Analysis of skeletal muscle lysates and His-tagged recombinant proteins using mouse parvalbumin mAb, dilution 1:1,000 (Green): [1] protein standard (Red), [2] mouse muscle, [3] full length human parvalbumin, [4] full length human calretinin, and [5] full length human calbindin. A band at 12kDa is detected in in muscle lysate and one at 18kDa in the His-tagged recombinant parvalbumin protein lane as expected since the His-tag and other vector derived sequence adds about 6kDa to the molecule. Note that the parvalbumin antibody is not cross-reactive with either calbindin or calretinin despite their related amino acid sequences.



Immunocytochemistry/Immunofluorescence: Parvalbumin Antibody (3C9) [NBP2-50038] - Analysis of rat cerebellum section stained with mouse mAb to parvalbumin, NBP2-50038, dilution 1:1,000, in green, and costained with chicken pAb to calbindin, dilution 1:2,000 in red. The blue is DAPI staining of nuclear DNA. Following transcardial perfusion of rat with 4% paraformaldehyde, brain was post fixed for 24 hours, cut to 45 uM, and free-floating sections were stained with above antibodies. Most Purkinje cells strongly express both parvalbumin and calbindin and so appear yellow, whereas basket, stellate and Golgi cells express parvalbumin alone and so appear are green.



Immunocytochemistry/Immunofluorescence: Parvalbumin Antibody (3C9) [NBP2-50038] - Adult rat cerebellum floating section was stained with NBP2-50038 at 1:1,000 (green). Parvalbumin is prominently expressed in the dendrites and perikarya of Purkinje cells and some interneurons in the molecular layer. Blue is a DNA stain.



Publications

Wang L, Xu W, Wang K et al. Chronic 40 Hz light flicker mitigates epileptogenesis through a visual pathway associated with the dorsal lateral geniculate nucleus shell Nature Communications 2025-10-17 [PMID: 41107227]

Sun XY, Liu L, Song YT et al. Two parallel medial prefrontal cortex-amygdala pathways mediate memory deficits via glutamatergic projection in surgery mice Cell reports 2023-06-30 [PMID: 37392387] (Mouse)

Sun X, Liu L, Wu T et al. Two Parallel Medial Prefrontal Cortex-Amygdala Circuits Mediate Memory Deficit via Glutamatergic Projection in Surgery Mice SSRN Electronic Journal 2022-10-12 (IF/IHC, Mouse)



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP2-50038

NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB7539	Goat anti-Mouse IgG (H+L) Secondary Antibody [HRP]
NBP1-97005-0.5mg	Mouse IgG1 Isotype Control (MG1)

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-50038

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

