

Product Datasheet

Apc4 Antibody (CIV1.1) - BSA Free NBP2-50105

Unit Size: 0.1 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 1

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-50105

Updated 9/9/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-50105



NBP2-50105

Apc4 Antibody (CIV1.1) - BSA Free

Product Information	
Unit Size	0.1 ml
Concentration	1 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	CIV1.1
Preservative	0.02% Sodium Azide
Isotype	IgG
Purity	Protein A purified
Buffer	PBS
Target Molecular Weight	92 kDa

Product Description	
Description	Novus Biologicals Mouse Apc4 Antibody (CIV1.1) - BSA Free (NBP2-50105) is a monoclonal antibody validated for use in WB and IP. Anti-Apc4 Antibody: Cited in 1 publication. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Mouse
Gene ID	29945
Gene Symbol	ANAPC4
Species	Human
Specificity/Sensitivity	CIV1.1 may be used for the affinity purification of the Anaphase Promoting Complex (APC/C) by peptide elution. Suggested Dilution for western blot: 1:500. It is suggested to blot for at least 2 hours (wet transfer) to ensure that APC4 is transferred efficiently. Mass spectrometry was used to confirm that the whole APC/C is co-precipitated. To establish the APC4 specific bands, it is suggested to do an APC4 immunoprecipitation assay showing input, IP and FT and blot for at least 2 hours.
Immunogen	C-terminus of Apc4

Product Application Details	
Applications	Western Blot, Immunoprecipitation
Recommended Dilutions	Western Blot 1:500, Immunoprecipitation 1:10 - 1:500
Application Notes	WB - It is suggested to blot for at least 2 hours (wet transfer) to ensure that APC4 is transferred efficiently. Mass spectrometry was used to confirm that the whole APC/C is co-precipitated. To establish the APC4 specific bands, it is suggested to do an Apc4 immunoprecipitation assay showing input, IP and FT and blot for at least 2 hours. Positive control(s): Purified APC/C by silver and mass spectrometry.

Publications

De K, Grubb TM, Zalenski AA et al. Hyperphosphorylation of CDH1 in glioblastoma cancer stem cells attenuates APC/CCDH1 activity and pharmacological inhibition of APC/CCDH1/CDC20 compromises viability Mol. Cancer Res. 2019-04-29 [PMID: 31036696] (WB, Human)





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP2-50105

NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB7539	Goat anti-Mouse IgG (H+L) Secondary Antibody [HRP]
NBP1-97019-5mg	Mouse IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-50105

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

