

Product Datasheet

CXCL16 Antibody (4.4) - BSA Free NBP2-50187

Unit Size: 0.1 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-50187

Updated 9/9/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-50187



NBP2-50187**CXCL16 Antibody (4.4) - BSA Free**


Product Information	
Unit Size	0.1 ml
Concentration	1 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	4.4
Preservative	0.02% Sodium Azide
Isotype	IgG1
Purity	Protein A purified
Buffer	PBS

Product Description	
Description	Novus Biologicals Mouse CXCL16 Antibody (4.4) - BSA Free (NBP2-50187) is a monoclonal antibody validated for use in WB and ICC/IF. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Mouse
Gene ID	58191
Gene Symbol	CXCL16
Species	Human
Immunogen	GST-CXCL16 (143-224aa) fusion protein


Product Application Details	
Applications	Western Blot, Immunocytochemistry/ Immunofluorescence
Recommended Dilutions	Western Blot 1:100 - 1:2000, Immunocytochemistry/ Immunofluorescence 1:10 - 1:500

Images

Western Blot: CXCL16 Antibody (4.4) [NBP2-50187] - Analysis of lysates prepared from HEK293 FT cells with a plasmid expressing the full length CXCL16 open reading frame. Lane 1: 293FT transfected cell lysate probed with anti-CXCL16 [4.4], Lane 2: 293FT mock cell lysate probed with the same antibody.



Immunocytochemistry/Immunofluorescence: CXCL16 Antibody (4.4) [NBP2-50187] - Staining of cos7 cells transfected with a plasmid containing a full-length CXCL16 open reading frame (a). 48 hours post transfection cells were fixed with 4% paraformaldehyde permeabilised with 0.1% Triton X-100 and stained using Anti-CXCL16 [4.4] and followed by secondary goat anti-mouse FITC (1:100). Nucleic acid DAPI counterstain is shown in blue. A negative control mock is also shown (b).





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP2-50187

NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB7539	Goat anti-Mouse IgG (H+L) Secondary Antibody [HRP]
NBP1-97005-0.5mg	Mouse IgG1 Isotype Control (MG1)

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-50187

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

