

# Product Datasheet

## CRISPR-Cas9 Antibody (6G12) - C-terminus - BSA Free NBP2-52398

Unit Size: 0.1 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

[www.novusbio.com](http://www.novusbio.com)



[technical@novusbio.com](mailto:technical@novusbio.com)

### Publications: 2

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:  
[www.novusbio.com/NBP2-52398](http://www.novusbio.com/NBP2-52398)

Updated 9/9/2025 v.20.1

Earn rewards for product  
reviews and publications.

Submit a publication at [www.novusbio.com/publications](http://www.novusbio.com/publications)

Submit a review at [www.novusbio.com/reviews/destination/NBP2-52398](http://www.novusbio.com/reviews/destination/NBP2-52398)



**NBP2-52398**

CRISPR-Cas9 Antibody (6G12) - C-terminus - BSA Free

Product Information	
Unit Size	0.1 ml
Concentration	1.0 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	6G12
Preservative	0.02% Sodium Azide
Isotype	IgG1 Kappa
Purity	Protein G purified
Buffer	PBS
Target Molecular Weight	158.4 kDa


Product Description	
Description	Novus Biologicals Mouse CRISPR-Cas9 Antibody (6G12) - C-terminus - BSA Free (NBP2-52398) is a monoclonal antibody validated for use in WB, ICC/IF, Simple Western, IP and ChIP. Anti-CRISPR-Cas9 Antibody: Cited in 2 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Mouse
Gene ID	901176
Species	Bacteria
Specificity/Sensitivity	This CRISPR-Cas9 antibody (6G12) - C-terminus is specific to Cas9 from <i>Streptococcus pyogenes</i> .
Immunogen	This CRISPR-Cas9 antibody (6G12) - C-terminus was raised against recombinant C-terminal fragment of <i>S.pyogenes</i> CRISPR/Cas9 (between amino acids 1139 to 1368) [UniProt# Q99ZW2]

Product Application Details	
Applications	Western Blot, Simple Western, Immunocytochemistry/ Immunofluorescence, Immunoprecipitation, Chromatin Immunoprecipitation (ChIP)
Recommended Dilutions	Western Blot 1:1000, Simple Western 10-20 ug/ml, Immunocytochemistry/ Immunofluorescence 1:500, Immunoprecipitation, Chromatin Immunoprecipitation (ChIP)




## Images

**Western Blot:** CRISPR-Cas9 Antibody (6G12) - C-terminus [NBP2-52398] - CRISPR-Cas9 Antibody (6G12) - C-Terminus [NBP2-52398] - Control HeLa cells (un-transfected) and HeLa cells expressing Flag-tagged *S. pyogenes*'s CRISPR-Cas9 under the control of PTight (Tet-ON) promoter. Samples were treated for 24 hours with 1ug/uL of Doxycyclin and lysed under native conditions. 30 ug of the whole cell lysate from each sample type per lane was separated by 7.5% SDS-PAGE. Nitrocellulose membrane was incubated with CRISPR-Cas9 antibody clone 6G12 (hybridoma supernatant diluted 1:100 at 4C O/N). After washing, the membranes were incubated with secondary HRP-coupled antibody and bands were visualized by ECL and exposure of X-ray films. Prestained marker bands were visualized with Blue Marker Antibody (NBP2-33376). The image shown is from 1 minute exposure time. Observed molecular weight is ~158 kDa.




**Immunocytochemistry/Immunofluorescence:** CRISPR-Cas9 Antibody (6G12) - C-terminus [NBP2-52398] - CRISPR-Cas9 Antibody (6G12) - C-Terminus [NBP2-52398] - HeLa cells or HeLa cells expressing Flag-tagged SpCas9 under the control of the PTight (Tet-ON) promoter were treated for 24h with 1ug/uL Doxycyclin, fixed and permeabilized with Methanol/Acetone and blocked in 2% BSA in PBS for 2 hours at RT. Cells were stained with 6G12 hybridoma supernatant at 1:10 at 4C O/N, followed by incubation with anti mouse-Alexa Fluor 488 coupled secondary antibody for 1h at RT. Nuclei were counter-stained with Hoechst 33342.




**Simple Western:** CRISPR-Cas9 Antibody (6G12) - C-terminus [NBP2-52398] - Image shows a specific band for Cas9 in 0.2 mg/mL of HeLa Cas9 lysate but not in HeLa WT lysate. This experiment was performed under reducing conditions using the 12-230 kDa separation system. Observed molecular weight is ~158 kDa.




**Chromatin Immunoprecipitation:** CRISPR-Cas9 Antibody (6G12) - C-terminus [NBP2-52398] - CRISPR-Cas9 Antibody (6G12) - C-Terminus [NBP2-52398] - CRISPR-Cas9 Antibody (6G12) [NBP2-52398] - NIH3T3 cells stably expressing GFP-H2B, nuclease dead Cas9, and a GFP-targeting gRNA were fixed with formaldehyde, harvested and sonicated to get 200-500bp DNA fragments. 50ug chromatin was incubated over night at 4C with the indicated antibodies (200ul hybridoma SN, 5ug anti-Flag [M2, Sigma]) followed by incubation with protein G beads for 3h at 4C. After washing chromatin was eluted from the beads and crosslinking was reversed over night at 65C. After a proteinase K digestion step, DNA was separated using phenol/chloroform/isoamyl alcohol, precipitated with ethanol/sodium acetate and dissolved in water. For qPCR, primers either targeting the GFP gene or as negative control non-targeted regions (Ppap2c +7122 and Prkcd +24069 from transcription start) were used.



Western Blot: CRISPR-Cas9 Antibody (6G12) - C-terminus [NBP2-52398] - CRISPR-Cas9 Antibody (6G12) - C-Terminus [NBP2-52398] - Whole cell protein from 293T cells transfected with Cas9-Flag (~150 kDa) was separated on a 7.5% gel by SDS-PAGE, transferred to PVDF membrane and blocked in 5% non-fat milk in TBST. The membrane was probed with 2 ug/mL anti-Cas9 (6G12) in 1% milk, and detected with an anti-mouse HRP secondary antibody using chemiluminescence.



Immunoprecipitation: CRISPR-Cas9 Antibody (6G12) - C-terminus [NBP2-52398] - CRISPR-Cas9 Antibody (6G12) - C-Terminus [NBP2-52398] - HEK293 cells expressing Flag-SpCas9 were lysed under native conditions. SpCas9 was immunoprecipitated at 4C from 300 ug of whole cell lysate with the 6G12 antibody and a 1:1 mixture of protein A/G sepharose. After 4x washing, the bound proteins were boiled off the beads, separated by 7.5% SDS-PAGE and transferred to nitrocellulose membranes, and SpCas9 was detected with a rabbit polyclonal Cas9 antibody. After washing, the membranes were incubated with secondary HRP-coupled antibody and bands were visualized by ECL and exposure of X-ray films.



## Publications

Johnston R, Seamon K, Saada E et al. Use of anti-CRISPR protein AcrIIA4 as a capture ligand for CRISPR/Cas9 detection Biosens Bioelectron 2019-06-18 [PMID: 31207570]

Giehl-Schwab J, Giesert F, Rauser B et al. Parkinson's disease motor symptoms rescue by CRISPRa-reprogramming astrocytes into GABAergic neurons EMBO molecular medicine 2022-04-04 [PMID: 35373464] (WB)



### **Novus Biologicals USA**

10730 E. Briarwood Avenue  
Centennial, CO 80112  
USA  
Phone: 303.730.1950  
Toll Free: 1.888.506.6887  
Fax: 303.730.1966  
nb-customerservice@bio-techne.com

### **Bio-Techne Canada**

21 Canmotor Ave  
Toronto, ON M8Z 4E6  
Canada  
Phone: 905.827.6400  
Toll Free: 855.668.8722  
Fax: 905.827.6402  
canada.inquires@bio-techne.com

### **Bio-Techne Ltd**

19 Barton Lane  
Abingdon Science Park  
Abingdon, OX14 3NB, United Kingdom  
Phone: (44) (0) 1235 529449  
Free Phone: 0800 37 34 15  
Fax: (44) (0) 1235 533420  
info.EMEA@bio-techne.com

### **General Contact Information**

www.novusbio.com  
Technical Support: nb-technical@bio-techne.com  
Orders: nb-customerservice@bio-techne.com  
General: novus@novusbio.com

### **Products Related to NBP2-52398**

---

NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB7539	Goat anti-Mouse IgG (H+L) Secondary Antibody [HRP]
NBP1-43319-0.5mg	Mouse IgG1 Kappa Isotype Control (P3.6.2.8.1)

---

### **Limitations**

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit [www.novusbio.com/guarantee](http://www.novusbio.com/guarantee)

Earn gift cards/discounts by submitting a review: [www.novusbio.com/reviews/submit/NBP2-52398](http://www.novusbio.com/reviews/submit/NBP2-52398)

Earn gift cards/discounts by submitting a publication using this product:  
[www.novusbio.com/publications](http://www.novusbio.com/publications)

