

Product Datasheet

CD81 Antibody (1.3.3.22) - Azide and BSA Free NBP2-54548-100ug

Unit Size: 100 ug

Store at -20 to -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-54548

Updated 10/23/2024 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-54548



NBP2-54548-100ug

CD81 Antibody (1.3.3.22) - Azide and BSA Free

Product Information	
Unit Size	100 ug
Concentration	1 mg/ml
Storage	Store at -20 to -80C. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	1.3.3.22
Preservative	No Preservative
Isotype	IgG1 Kappa
Purity	Protein A or G purified
Buffer	10 mM PBS
Target Molecular Weight	26 kDa

Product Description	
Description	1.0 mg/ml of antibody purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/ml WITH BSA & azide (NBP2-53318). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C.
Host	Mouse
Gene ID	975
Gene Symbol	CD81
Species	Human, Mouse, Rat
Specificity/Sensitivity	Recognizes a protein of 26kDa, identified as CD81 (Workshop VI; Code CD81.1)). CD81 has a very broad cellular distribution, being expressed on T- and B-lymphocytes, NK cells, thymocytes, eosinophils, fibroblasts, epithelial and endothelial cells. Neutrophils, erythrocytes and platelets are negative, while monocytes are variably positive. CD81 is a member of a family of tetraspans transmembrane proteins, including CD9, CD37, CD53, CD63, and CD82. It associates with CD19, CD21, Leu 13, and integrins on cell membrane and is involved in signal transduction in B lymphocyte development and cell adhesion. CD81 also acts as a receptor for the envelope protein E2 of chronic hepatitis C virus. Antibodies to CD81 have anti-proliferative effects on different lymphoid cell lines, particularly those derived from large cell lymphomas.
Immunogen	Recombinant full-length human CD81 protein (Uniprot: P60033)


Product Application Details	
Applications	Western Blot, Flow Cytometry, Flow (Cell Surface), Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready, Knockout Validated, Single Cell Western
Recommended Dilutions	Western Blot 0.5 - 1.0 ug/ml, Flow Cytometry 0.5 - 1 ug/million cells, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 0.5 - 1 ug/ml, Immunohistochemistry-Paraffin 0.5 - 1 ug/ml, Flow (Cell Surface), CyTOF-ready, Knockout Validated, Single Cell Western

Application Notes


Functional Studies: Order Ab without Azide.
 Immunohistochemistry (Formalin-fixed): 2-4ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes. Optimal dilution for a specific application should be determined.
 Western Blot: 1-2ug/ml for 2 hours at RT.

Images


Immunohistochemistry-Paraffin: CD81 Antibody (1.3.3.22) - Azide and BSA Free [NBP2-54548] - Formalin-fixed, paraffin-embedded human Lymph Node stained with CD81 Mouse Monoclonal Antibody (1.3.3.22).




Flow Cytometry: CD81 Antibody (1.3.3.22) - Azide and BSA Free [NBP2-54548] - Flow cytometry analysis of bead-bound exosomes derived from MCF-7 cells. Unstained exosomes (gray) or exosomes stained with CF568-labeled CD81 monoclonal antibody (1.3.3.22) (orange).




Immunohistochemistry-Paraffin: CD81 Antibody (1.3.3.22) - Azide and BSA Free [NBP2-54548] - Formalin-fixed, paraffin-embedded human Tonsil stained with CD81 Mouse Monoclonal Antibody (1.3.3.22).




Flow Cytometry: CD81 Antibody (1.3.3.22) - Azide and BSA Free [NBP2-54548] - Flow cytometry analysis of MCF-7 cells unstained (gray) or stained with CD81 mouse monoclonal antibody (1.3.3.22) and goat anti-mouse CF647 (red).



Knockout Validated: CD81 Antibody (1.3.3.22) - Azide and BSA Free [NBP2-54548] - CD81 was detected in immersion fixed HeLa human cervical epithelial carcinoma cell line but is not detected in CD81 knockout (KO) HeLa cell line using Mouse Anti-Human CD81 Monoclonal Antibody (Catalog # NBP2-53318) at 1 ug/mL for 3 hours at room temperature. Cells were stained using the NorthernLights™ 557-conjugated Anti-Mouse IgG Secondary Antibody (red; Catalog # NL007) and counterstained with DAPI (blue). Specific staining was localized to the plasma membrane.



Positive (HeLa cells)



Negative (HeLa KO cells)



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP2-54548-100ug

HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB720-B	Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]
NBP1-43319-0.5mg	Mouse IgG1 Kappa Isotype Control (P3.6.2.8.1)
NBP2-60215-50ug	Recombinant Human CD81 hIgG-His Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-54548

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

